

GEOMETRIC RIGIDITY OF $\times m$ INVARIANT MEASURES

MICHAEL HOCHMAN

ABSTRACT. Let μ be a probability measure on $[0, 1]$ which is invariant and ergodic for $T_a(x) = ax \bmod 1$, and $0 < \dim \mu < 1$. Let f be a local diffeomorphism on some open set. We show that if $E \subseteq \mathbb{R}$ and $(f\mu)|_E \sim \mu|_E$, then $f'(x) \in \{\pm a^r : r \in \mathbb{Q}\}$ at μ -a.e. point $x \in f^{-1}E$. In particular, if g is a piecewise-analytic map preserving μ then there is an open g -invariant set U containing $\text{supp } \mu$ such that $g|_U$ is piecewise-linear with slopes which are rational powers of a .

In a similar vein, for μ as above, if b is another integer and a, b are not powers of a common integer, and if ν is a T_b -invariant measure, then $f\mu \perp \nu$ for all local diffeomorphisms f of class C^2 . This generalizes the Rudolph-Johnson Theorem and shows that measure rigidity of T_a, T_b is a result not of the structure of the abelian action, but rather of their smooth conjugacy classes: if U, V are maps of \mathbb{R}/\mathbb{Z} which are C^2 -conjugate to T_a, T_b then they have no common measures of positive dimension which are ergodic for both.

1. INTRODUCTION

1.1. Background. The motivating problem of this paper is to understand, for “structured” Borel probability measures on \mathbb{R} , which transformations can map one measure to another, in whole or in part, and how the structure of the measures determines this. More precisely, writing $f\mu$ for the measure $f\mu(A) = \mu(f^{-1}A)$, we ask what one can say about maps f for which $f\mu$ is non-singular with respect to μ , or with respect to some other measure ν . The expectation is that highly structured measures should be preserved by a small number of maps whose structure reflects that of the measure.

There are a few elementary things one can say. Since any two non-atomic probability measures on \mathbb{R} can be mapped to each other by a continuous function, one must impose some regularity assumption for the question to make sense. In this paper we consider diffeomorphisms of the line, and denote the set of such maps by $\text{diff}(\mathbb{R})$. Write $\text{diff}^k(\mathbb{R})$ for the set of diffeomorphisms of class C^k . Also, there are some trivial cases of measures μ such that $\mu, f\mu$ are non-singular for many

2000 *Mathematics Subject Classification.* 37A10, 37E10, 37A45, 37C45, 28A80.

Key words and phrases. Measure rigidity, invariant measure, interval map, fractal geometry, geometric measure theory, scenery flow.

Research supported by NSF grant 0901534.

$f \in \text{diff}(\mathbb{R})$, namely if μ has a Lebesgue component or atoms. One must therefore consider measures without such components. Recall that the local dimension of a measure μ at x is the limit

$$D(\mu, x) = \lim_{r \rightarrow 0} \frac{\log \mu(B_r(x))}{\log r}$$

where $B_r(x)$ is the ball of radius r around x . In general the local dimension may not exist but it exists in the cases which will interest us. We will work with measures of *intermediate dimension*, meaning that $0 < D(\mu, x) < 1$ at μ -a.e. x . This rules out Lebesgue component, and also any zero-dimensional component, including atoms.

In this work we consider probability measures which are invariant under the maps $T_a : [0, 1] \rightarrow [0, 1]$ defined for integers $a \geq 2$ by

$$T_a x = bx \bmod 1$$

i.e. measures such that $T_a \mu = \mu$. We sometimes view T_a as a map of the torus \mathbb{R}/\mathbb{Z} . We note that there is a very rich supply of T_a -invariant measures, including some self-similar measures but also many (most) which are not.

For T_a -invariant measures most work to date has focused on their behavior for maps which are related to the group structure of \mathbb{R}/\mathbb{Z} , i.e. when f is one of the maps T_b (an endomorphism of \mathbb{R}/\mathbb{Z}) or translation by an element of \mathbb{R}/\mathbb{Z} . The principal result for endomorphisms is the measure rigidity theorem of Rudolph [17] and Johnson [11]. Write $a \sim b$ if a, b are powers of a common integer, i.e. $a = c^k$ and $b = c^m$ for some $c, k, m \in \mathbb{N}$, and otherwise write $a \not\sim b$. The Rudolph-Johnson Theorem states that if μ is a T_a -invariant measure whose ergodic components all have entropy strictly between 0 and $\log a$, then it is not preserved by T_b for any $b \not\sim a$. The result can be slightly improved, using a later result of Rudolph-Johnson [10], to conclude that $\mu \perp \nu$ for every T_b -invariant measure ν of intermediate dimension. To date this is essentially the best result towards Furstenberg's $\times 2, \times 3$ conjecture, which predicts that there should be no non-atomic measures except Lebesgue which are jointly invariant under T_a and T_b for $a \not\sim b$. See [13] for a survey of related conjectures and results.

For translations we are aware only of the work of Host [8]. Define μ to be conservative for a subgroup $\Lambda \subseteq \mathbb{R}/\mathbb{Z}$ if for every set A with $\mu(A) > 0$ there is some $0 \neq r \in \Lambda$ such that $\mu(A \cap (A+r)) > 0$. For the groups $\Lambda = \mathbb{Z}[\frac{1}{b}]$ of b -adic rationals when $\text{gcd}(a, b) = 1$, or for the cyclic subgroup generated by an element $r \in \mathbb{R}/\mathbb{Z}$ such that $\{T_a^n r\}_{n \in \mathbb{N}}$ is dense in \mathbb{R}/\mathbb{Z} , Host showed, using methods of harmonic analysis, that the only conservative T_a -invariant measure is Lebesgue measure. Note that these results do not require any assumption about the dimension of the measure, but also they do not directly relate to our question, since they do not say anything about preservation of μ under a particular rotation.

In another direction, the question we are interested in has been studied in the fractal geometry literature for rather general functions f , e.g. C^1 or bi-Lipschitz, but for restricted classes of measures such as self-similar measures on attractors of iterated function systems. In situations like these it has been shown that measures arising from different parameters are mutually singular, and cannot be easily deformed into each other, see for example [3]. Related questions for certain classes of Cantor sets are known, for instance see [2, 1, 19].

1.2. Statement of results. Write $\mu \sim \nu$ and $\mu \perp \nu$ to indicate that the measures are equivalent or singular, respectively, and let $\mu|_E$ denote the restricted measure $\mu|_E(A) = \mu(A \cap E)$. Thus μ, ν are non-singular if and only if $\mu|_E \sim \nu|_E$ for some set E with $\nu(E) > 0$.

Theorem 1.1. *Let μ be a T_a -ergodic measure of intermediate dimension. Then there exists $n \in \mathbb{N}$ such that if $f \in \text{diff}(\mathbb{R})$ and $(f\mu)|_E \sim \mu|_E$, then*

$$f'(x) \in \{\pm a^{k/n} : k \in \mathbb{Z}\} \quad \text{for } \mu\text{-a.e. } x \in f^{-1}E$$

More generally, if ν is another T_a -ergodic measure and $(f\mu)|_E \sim \nu|_E$ then there exists $t \in \mathbb{R}$ such that

$$f'(x) \in \{\pm t \cdot a^{k/n} : k \in \mathbb{Z}\} \quad \text{for } \mu\text{-a.e. } x \in f^{-1}E$$

While we have stated the result for diffeomorphisms of \mathbb{R} , the result is of a local nature and immediately applies to partially defined or piecewise diffeomorphisms.

Theorem 1.1 is close to optimal. One clearly cannot hope to get information about f except on the support of $\mu|_E$. On the other hand, a T_a -invariant measure μ is also invariant for T_{a^n} for every $n \in \mathbb{N}$ and sometimes also for $n = 1/m$ when $a^{1/m} \in \mathbb{N}$. Thus one cannot expect that some intrinsic property of μ will encode a , and the best one can hope for is a power of a . The ergodicity assumption is necessary also: for example fix a T_2 -invariant measure μ of intermediate dimension and form $\nu = \frac{1}{2}\mu + \frac{1}{2}T_3\mu$, which is also T_2 -invariant. It is easy to see that there is a piecewise linear map f with $f\nu, \nu$ non-singular, and with slopes 2 and 3 on sets of positive ν -measure.

We do not know whether it can happen in the theorem that $a^{1/n}$ is not an integer. We also do not know whether a version of the theorem is true under Lipschitz (rather than differentiability) conditions on f .

Under mild additional assumptions, Theorem 1.1 implies that very few maps can preserve a T_a -ergodic measure of intermediate dimension.

Corollary 1.2. *Let μ and n be as in the theorem.. Then every piecewise analytic map of $[0, 1]$ which preserves μ is piecewise linear on an open set U containing the support of μ and on U has slopes of the form $\pm a^{k/n}$, $k \in \mathbb{N}$.*

Corollary 1.3. *If μ is a T_a -ergodic measure on \mathbb{R}/\mathbb{Z} which has intermediate dimension and is globally supported, then every C^1 -map which preserves μ has the form $T_{a'}$ for an integer $a' \sim a$.*

Easy examples show that there can be piecewise linear maps other than T_a which preserve μ , for example one can easily construct them by hand for the "uniform" measure on the middle third Cantor set (i.e. Hausdorff measure at the appropriate dimension, normalized to mass 1), which is T_3 -invariant.

A special case of the above we recover the Rudolph-Johnson Theorem [17, 11]. One may speculate that Theorem 1.1 holds without the entropy assumption, but proving this would imply the full $\times 2, \times 3$ conjecture.

Our methods also allow us to generalize the Rudolph-Johnson theorem in other ways.:

Theorem 1.4. *If $a \not\sim b$ and μ, ν are respectively T_a and T_b -ergodic measures of intermediate dimension, then $f\mu \perp \nu$ for every $f \in \text{diff}^2(\mathbb{R})$.*

We can eliminate the ergodicity and regularity assumptions under some (rather weak) additional hypotheses, for example no ergodicity is needed if f is affine, or when when $f \in \text{diff}(\mathbb{R})$ but the ergodic components of ν under T_b do not have spectrum of the form $\frac{n}{\log a}$.

An interesting consequence of the theorem above is that the measure rigidity phenomenon in the Rudolph-Johnson Theorem is not a consequence of properties of the abelian action generated by T_a and T_b , but rather of the smooth conjugacy classes of the individual maps T_a, T_b :

Corollary 1.5. *Let $a \not\sim b$ and let f, g be self-maps of \mathbb{R}/\mathbb{Z} which are (separately) C^2 -conjugate to T_a, T_b , respectively. Then there is no measure of positive Hausdorff dimension which is ergodic for both f and g , except possibly one which is equivalent to Lebesgue, and this occurs precisely when the conjugating maps differ by a rotation.*

Note that for f, g as above there will generally be no invariant measures at all, but it is hard to verify this for any particular pair of conjugates. It is known that if f, g commute then they are simultaneously C^0 -conjugate to T_a, T_b [12], and then Corollary 1.5 follows from the Rudolph-Johnson Theorem.

After this paper was completed P. Shmerkin suggested another approach which proves Theorem 1.4 for $f \in \text{diff}(\mathbb{R})$ and non-ergodic measures, but which does not give any version of Theorem 1.1. This will appear elsewhere.

1.3. Methods. To arrive at these results we study measures on \mathbb{R}^d through the dynamics of the 1-parameter, measure-valued family obtained by "zooming in" on typical points for the measure. These families are called sceneries, and for

the measures we are considering they behave like generic orbits in an appropriate dynamical system. Many variants of the notion of a scenery have appeared in the fractal geometry literature, see e.g. Bedford and Fisher [1], and have been used as a technical tool in the study of T_m -invariant measures, in disguised form in Furstenberg's paper [4] and more recently in [7]. Our definition of the scenery flow follows that of Gavish [5]. A systematic study of this notion and related ones can be found in [6].

Briefly, we show that for a T_a -invariant measure the sceneries equidistribute for an ergodic flow whose pure-point spectrum contains a rational multiple of $\frac{1}{\log a}$, and the remaining spectrum comes from the original dynamics of T_a . Since these flows are associated to measures in a geometric way, they are invariants of the measure under the application of differentiable, locally bijective maps. Furthermore, the flows derived at different points of a T_a -invariant measure may have different phases with respect eigenvalues of the form $\frac{n}{\log a}$, and applying a smooth map shifts the phase by the logarithm of the derivative. The behavior of these phases underlies the proof of Theorems 1.1 and 1.4.

This method of proof gives an effective condition for a measure μ to be a smooth image of a T_a -invariant measure, namely the spectrum of the associated flow must contain a rational multiple of $\log a$. In a sense, this explains how μ "encodes" the arithmetic class of the dynamics which generated it.

1.4. Related questions. We end this introduction with some open questions. Let us begin by pointing out a connection between Theorem 1.4 and another conjecture of Furstenberg [4]: If $X, Y \subseteq [0, 1]$ are closed and invariant, respectively, under T_a and T_b for $a \not\sim b$, then for every affine map $f(x) = ux + v$,

$$\dim(X \cap fY) \leq \max\{0, \dim X + \dim Y - 1\}$$

This says that all affine images of Y should intersect X in as small a set as possible. Theorem 1.4 gives an analog of this for measures, though of course singularity of measures implies nothing about the intersection of their topological supports. On the other hand, note that Theorem 1.4 has content even when both μ, ν are globally supported.

Returning to the $\times 2, \times 3$ conjecture, the topological version was proved by Furstenberg with no entropy assumptions: any closed infinite subset of $[0, 1]$ which is invariant under T_a, T_b for $a \not\sim b$ is the entire interval. One may similarly ask for topological versions of our results:

Problem 1.6. Suppose $A \subseteq [0, 1]$ is an infinite, proper closed T_a -invariant subset. If $f \in \text{diff}^1(\mathbb{R})$ preserves A , must $|f'(x)| \in \{a^r : r \in \mathbb{Q}\}$ for all non-isolated $x \in A$?

Problem 1.7. Let $a \not\sim b$ and let f, g be maps of \mathbb{R}/\mathbb{Z} which are (separately) conjugate, respectively, to T_a, T_b . If the conjugating maps are sufficiently smooth, can we conclude that there are no infinite, closed proper subsets of \mathbb{R}/\mathbb{Z} which are jointly f - and g -invariant?

We do not have answers except when for some $s > 0$ the s -dimensional Hausdorff measure is positive and finite on A . Then one can apply our results to this measure.

In another direction, there is a strengthening of the Rudolph-Johnson Theorem due Host [8], which asserts that for $\gcd(a, b) = 1$, if μ is T_a -invariant of intermediate dimension, then μ -a.e. point x equidistributes for Lebesgue measure under the action of T_b (in this case x is said to be *normal* in base b). It is natural to ask whether the same is true when the measure is distorted by a nice enough map. Let us formulate this question in the simplest and most plausible case:

Problem 1.8. Let μ be a T_a -invariant measure and of intermediate dimension, and $\gcd(a, b) = 1$. If $f(x) = ux + v$, $u \neq 0$, is $f\mu$ -a.e. point normal in base b ?

Finally, it is very likely that analogs of Theorems 1.1 and 1.4 ours hold for more general interval maps and in higher dimensional settings, for instance for measures on the torus which are invariant under hyperbolic automorphisms under suitable assumptions. However, in neither case does it appear that our methods apply directly.

1.5. Organization. In the next section we describe our methods in more detail. Proofs are given in the following sections. We assume familiarity with basic notions in ergodic theory, recalling some definitions as we go; for an introduction see [18]. For background on geometric measure theory see [15].

1.6. Acknowledgment. I am grateful to Jean Bourgain, Elon Lindenstrauss and Pablo Shmerkin for their comments.

2. MAIN ELEMENTS OF THE PROOFS

In this section we give our main definitions and technical results, and derive the main theorems from them. The remaining proofs are provided in the next section.

2.1. The scenery flow. Let $\mathcal{M} = \mathcal{M}_d$ denote the space of Radon measures on \mathbb{R}^d , endowed with the weak topology. We use the term *measure* for Radon measures on \mathbb{R}^d , and denote measures by μ, ν, σ, τ etc. We reserve the term *distribution* for Borel probability measures on \mathcal{M} , which we denote by P, Q, R etc. The space of distributions carries a measurable structure defined by declaring the map $\mu \mapsto \mu(A)$ to be measurable for all Borel sets $A \subseteq \mathcal{M}$. Write λ for Lebesgue measure and δ_z for the point mass at z , which is a measure when $z \in \mathbb{R}^d$ and a distribution when $z \in \mathcal{M}$. Let $\text{supp } \mu$ denote the topological support of a measure, that is,

the complement of the union of all open sets of μ -measure zero. We write \sim for equivalence of measures or distributions, and also write $z \sim \mu$ to indicate that z is distributed according to μ ; which is intended will be clear from the context.

For $x \in \mathbb{R}^d$ let $U_x : \mathbb{R}^d \rightarrow \mathbb{R}^d$ denote the translation map

$$U_x(y) = y - x$$

and for $t \in \mathbb{R}$ let $S_t : \mathbb{R}^d \rightarrow \mathbb{R}^d$ denote the scaling map

$$S_t(x) = e^t x$$

Note the exponential time scale, which makes $S = (S_t)_{t \in \mathbb{R}}$ into an action of the additive group \mathbb{R} on \mathbb{R}^d . These operations induce maps on \mathcal{M} : for $\mu \in \mathcal{M}$ we have $U_x \mu(A) = \mu(A + x)$ and $S_t \mu(A) = \mu(e^{-t} A)$.

Let $\mu \mapsto \mu^\square$ denote the operation of normalizing a measure to have unit mass on $[-1, 1]^d$ and restricting it to this cube, i.e.

$$\mu^\square = \frac{1}{\mu([-1, 1]^d)} \mu|_{[-1, 1]^d}$$

Let

$$\mathcal{M}^\square = \{\text{Probability measures on } \mathbb{R} \text{ supported on } [-1, 1]^d\}$$

and define $S_t^\square : \mathcal{M}^\square \rightarrow \mathcal{M}^\square$ by

$$S_t^\square \mu = (S_t \mu)^\square$$

so that $S^\square = (S_t^\square)_{t \geq 0}$ is a semigroup acting on the set of $\mu \in \mathcal{M}^\square$ with $0 \in \text{supp } \mu$. This is a Borel subset of \mathcal{M} , and the action is Borel, though not continuous (it is discontinuous at measures which give positive mass to the boundary of $[-1, 1]^d$).

Definition 2.1. Let $\mu \in \mathcal{M}$ and $x \in \text{supp } \mu$. The *scenery* of μ at x is the orbit of $U_x \mu$ under S^\square , i.e. the one-parameter family

$$\mu_{x,t} = S_t^\square(U_x \mu) \quad t \in \mathbb{R}^+$$

In other words, the scenery is what one sees when “zooming in” to μ at x , restricting and normalizing the measure as we go.

In order to discuss the limiting behavior of the scenery, note that \mathcal{M}^\square may be identified with the weak-* compact set of probability measures on $[-1, 1]^d$. Thus we may speak of convergence of distributions on \mathcal{M}^\square .

Definition 2.2. A measure $\mu \in \mathcal{M}$ *generates* a distribution P at $x \in \text{supp } \mu$ if the scenery $(\mu_{x,t})_{t \geq 0}$ equidistributes for P , that is, if the uniform measure on the path $(\mu_{x,t})_{0 \leq t \leq T}$ converges weak-* to P as $T \rightarrow \infty$. Equivalently, for every $f \in C(\mathcal{M}^\square)$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(\mu_{x,t}) dt = \int f dP$$

In this case $(\mathcal{M}^\square, P, S^\square)$ is called the *scenery flow* of μ at x .

For a discussion of the properties of distributions generated in this way see [6]. We mention a few basic facts. First, if there is a positive μ -measure of points at which μ generates some distribution (which may vary from point to point), then μ -a.e. one of the distributions is S^\square -invariant.¹ Second, standard density arguments show that if $\nu \ll \mu$ then a ν -typical point generates a distribution for ν if and only if it does for μ , and in this case the distributions are the same. This applies in particular when $\nu = \mu|_A$.

The following simple observation is a key ingredient in our arguments. Let $\text{diff}_+(\mathbb{R}) \subseteq \text{diff}(\mathbb{R})$ denote the subgroup of orientation-preserving maps.

Lemma 2.3. *Let $\mu \in \mathcal{M}(\mathbb{R})$, $x \in \text{supp } \mu$ and $f \in \text{diff}_+^1(\mathbb{R})$. Then after a time-shift of $s = \log f'(x)$ the sceneries $\mu_{x,t}$ and $(f\mu)_{f(x),t}$ are asymptotic, i.e.*

$$\lim_{t \rightarrow \infty} (\mu_{x,t} - (f\mu)_{f(x),t-s}) = 0 \quad (\text{weak-}^*)$$

In particular μ generates P at x if and only if $f\mu$ generates P at $f(x)$.

The proof is immediate from the fact that, locally, f acts like S_s near x .

The assumption that f preserve orientation is necessary for the conclusion that the scenery flows are the same, but if f is orientation reversing then the scenery flows are isomorphic as measure preserving flows by way of the map induced on \mathcal{M}^\square from $x \mapsto -x$. We omit the details.

2.2. The scenery flow of T_a -invariant measures and its spectral properties.

Next we describe the scenery flow of a T_a -invariant measure and, more generally, products of such measures. Recall that the diagonal action of T_a on $[0, 1]^d$ is given by $T_a(x) = (T_a x_1, T_a x_2, \dots, T_a x_d)$. A product of T_a -invariant measures is invariant under the diagonal action.

We require two more standard constructions, which we recall briefly. First, the natural extension of an ergodic system (Ω, ν, T) is an invertible ergodic system $(\tilde{\Omega}, \tilde{\nu}, \tilde{T})$ factoring onto (Ω, ν, T) and characterized by the property that every factor map from an invertible system to (Ω, ν, T) factors through $(\tilde{\Omega}, \tilde{\nu}, \tilde{T})$. The natural extension may be realized as the inverse limit of the diagram of factor maps

$$\dots \rightarrow (\Omega, \nu, T) \xrightarrow{T} (\Omega, \nu, T) \xrightarrow{T} (\Omega, \nu, T)$$

See also Section 3.

Second, the t_0 -suspension of the discrete time system (Ω, ν, T) is the flow defined on $\Omega \times [0, t_0]$ by

$$T_t(\omega, s) = (T^{\lfloor s/t_0 \rfloor} \omega, \{ \frac{s+t}{t_0} \} t_0)$$

¹Note that S^\square acts discontinuously, so this is not a complete triviality.

where $[r]$ and $\{r\}$ are the integer and fractional parts of r , respectively. This flow preserves the product measure $\nu \times \frac{1}{t_0} \lambda|_{[0, t_0]}$.

Proposition 2.4. *Let μ_1, \dots, μ_d be T_a -invariant measures and $\mu = \times_{i=1}^d \mu_i$. Then μ generates an S^\square -ergodic distribution P_x at a.e. point x , and the system $(\mathcal{M}^\square, P_x, S^\square)$ arises as a factor of the $\log b$ -suspension of the ergodic component $\mu^{(x)}$ of x in $([0, 1]^d, \mu, T_a)$. In particular P_x depends only on the ergodic component $\mu^{(x)}$ of x . Furthermore, if $\mu^{(x)}$ has intermediate dimension then P_x is supported on measures of intermediate dimension.*

This is proved in Section 3.

Write $e(t) = \exp(2\pi it)$. Recall that $\alpha \in \mathbb{R}$ is an eigenvalue of an ergodic measure preserving system $(\Omega, \mathcal{B}, \nu, T)$ if there is a complex function $\varphi \in L^2$ such that $\varphi \circ T = e(\alpha)\varphi$, and α is an eigenvalue of a measure-preserving flow $(\Omega, \mathcal{B}, \nu, (T_t)_{t \in \mathbb{R}})$ if there is a function $\varphi \in L^2$ such that $\varphi \circ T_t = e(\alpha t)\varphi$ for all $t \in \mathbb{R}$. Such φ are called eigenfunctions, and for ergodic transformations and flows they a.s. have constant modulus, which we shall always assume has been normalized to modulus 1. We denote the set of eigenvalues by Σ , with subscripts to indicate the system in question.

Theorem 2.5. *Let μ be T_a -invariant with intermediate entropy. Let $\Sigma_{\mu^{(x)}}$ denote the spectrum of the ergodic component $\mu^{(x)}$ of $([0, 1], \mu, T_a)$ to which x belongs. Let P_x be the distribution generated at x and Σ_{P_x} the spectrum of $(\mathcal{M}^\square, P_x, S^\square)$. Then there is an $n \in \mathbb{N}$ such that*

$$\frac{n}{\log a} \mathbb{Z} \subseteq \Sigma_{P_x} \subseteq \frac{1}{\log a} \Sigma_{\mu^{(x)}} \cup \frac{n}{\log a} \mathbb{Z}$$

One cannot assume that $n = 1$ since if μ is T_a invariant then for every n it is also T_{a^n} invariant, since $T_{a^n} = T_a^n$.

Proof of Theorem 1.4 under spectral assumptions. Suppose $f \in \text{diff}^1$ and μ, ν are respectively T_a, T_b invariant, have intermediate entropy, and the ergodic components of ν do not have pure point spectrum of the form $\frac{n}{\log a}$, $n \in \mathbb{Z}$. By the theorem above the scenery flows generated a.e. by μ have pure point spectrum of this form. If $f\mu \not\sim \nu$ then by Lemma 2.3, with positive μ -probability the scenery flow generated by μ at x is isomorphic to the one generated by ν at $f(x)$. These possibilities are incompatible. \square

2.3. The distribution of phases. More refined information can be obtained from the distribution of phases of the eigenfunctions of the scenery flow. That is, for a T_a -ergodic measure and typical points x, y , we may consider the sceneries at x and y and compare the relative phase of the eigenfunctions corresponding to $\alpha = k/\log a$.

Recall that a joining of S^\square -invariant distributions P_1, P_2 is a distribution P on $\mathcal{M}^\square \times \mathcal{M}^\square$ which projects to P_i on the i -th coordinate, and which is invariant under the diagonal flow S^\square given by $S_t^\square(\mu, \nu) = (S_t^\square \mu, S_t^\square \nu)$. When P_1, P_2 are ergodic, the ergodic components of a joining of P_1, P_2 are also joinings of P_1, P_2 . A P -joining is a joining of P with itself.

Let μ be a measure generating and ergodic distribution P_x at x . Note that there is a bijective correspondence between pairs $(\sigma, \tau) \in \mathcal{M}_1^\square \times \mathcal{M}_1^\square$ and product measures $\sigma \times \tau \in \mathcal{M}_2^\square$, and that the set of product measures is closed in $\mathcal{M}^\square \times \mathcal{M}^\square$. Therefore the accumulation points of the sceneries of $\mu \times \mu$ are product measures, and if $\mu \times \mu$ generates a scenery flow it is supported on product measures. Furthermore, if $\mu \times \mu$ generates a scenery flow $P_{x,y}$ at (x, y) one may verify that, making the identification between product measures and pairs, $P_{x,y}$ is a joining of the scenery flows P_x, P_y generated by μ at x and y . Note that by Proposition 3.7, if μ is T_α -invariant then $\mu \times \mu$ indeed generates sceneries a.e.

Let P be an S^\square -invariant and ergodic distribution with an eigenvalue α and corresponding eigenfunction φ . Then the function $p_\alpha : \mathcal{M}^\square \times \mathcal{M}^\square \rightarrow \mathbb{C}$, defined by

$$p_\alpha(\sigma, \tau) = \frac{\varphi(\sigma)}{\varphi(\tau)}$$

is a.e. invariant on any ergodic P -joining, since

$$p_\alpha(S_t^\square \sigma, S_t^\square \tau) = \frac{\varphi(S_t^\square \sigma)}{\varphi(S_t^\square \tau)} = \frac{e(\alpha t)\varphi(\sigma)}{e(\alpha t)\varphi(\tau)} = p_\alpha(\sigma, \tau)$$

Therefore if R is a P -joining then p_α is constant on a.e. ergodic component of R , and if R is ergodic then we may define

$$p_\alpha(R) = R\text{-a.s. value of } p_\alpha(\cdot, \cdot)$$

Let μ generate some distribution P at a.e. point and assume that $\mu \times \mu$ generates an ergodic P -joining $P_{x,y}$ at $\mu \times \mu$ -a.e. point. Fix a μ -typical x_0 so that $P_{x_0,y}$ is defined for μ -a.e. y and for such y let

$$p_\alpha(\mu, x_0, y) = p_\alpha(P_{x_0,y})$$

Definition 2.6. Let μ be a measure which a.e. generates P , and such that $\mu \times \mu$ generates an ergodic distribution at a.e. point. Let $\alpha \in \Sigma_P$. For a μ -typical point x_0 the *phase measure* $\theta_\alpha = \theta_\alpha(\mu, x_0)$ is the push-forward of μ under the map $y \mapsto p_\alpha(\mu, x_0, y)$.

Under the further assumption that $\mu \times \mu \times \mu$ generates an ergodic distribution at a.e. point, the dependence of the phase measure on x_0 is very mild. Indeed, if we choose another point x_1 then, by considering the threefold joining Q generated

by $\mu \times \mu \times \mu$ at (x_0, y, x_1) we find that, writing $(\sigma_1, \sigma_2, \sigma_3)$ for a Q -typical element,

$$\begin{aligned} p_\alpha(\mu, x_1, y) &= \frac{\varphi(\sigma_3)}{\varphi(\sigma_2)} \\ &= \frac{\varphi(\sigma_3)}{\varphi(\sigma_1)} \cdot \frac{\varphi(\sigma_1)}{\varphi(\sigma_2)} \\ &= c \cdot p_\alpha(\mu, x_0, y) \end{aligned}$$

where $c = p_\alpha(\mu, x_1, x_0)$ does not depend on y . Thus $p_\alpha(\mu, x_0, \cdot)$ depends on x_0 only up to a rotation, and the measures $\theta_\alpha(\mu, x_0)$ and $\theta_\alpha(\mu, x_1)$ are rotations of one another. Since we shall be interested in properties which are independent of rotation, we often suppress the dependence on x_0 and write $\theta_\alpha(\mu)$.

Theorem 2.7. *Let μ be a T_a -invariant measure of intermediate entropy whose ergodic components a.s. generate the same distribution P . Let $\alpha \in \Sigma_P$ and $\theta_\alpha = \theta_\alpha(\mu)$. Then*

- (1) *If $\alpha \in \Sigma_P \setminus \frac{1}{\log a} \mathbb{Q}$ then θ_α is Lebesgue measure.*
- (2) *If $\alpha \in \Sigma_P \cap \frac{1}{\log a} \mathbb{Q}$ then θ_α is singular with respect to Lebesgue measure.*
- (3) *If $\alpha \in \Sigma_P \cap \frac{1}{\log a} \mathbb{Z}$ and μ is ergodic then θ_α consists of a single atom.*

We do not know how large the phase measure can be for non-ergodic μ . With our methods we can go a little further and show that $\underline{\dim} \theta_\alpha(\mu) \leq 1 - \dim \mu$, where $\underline{\dim}$ is the lower Hausdorff dimension of μ . On the other hand, starting with a T_a -ergodic measure μ with $\alpha = \frac{n}{\log a}$ in the spectrum of the scenery flow, the measure $\nu = \sum_{b=1}^{\infty} 2^{-b} T_b \mu$ is again T_a -invariant and its phase measure consists of atoms at $e(\log b / \log a)$, $b \in \mathbb{N}$. We suspect that the phase measure is always atomic or at least of dimension zero, but we have not resolved this.

We next examine how the phase distribution changes when a smooth map is applied to a measure. First, suppose that μ is a measure satisfying the conditions in Definition 2.6 and the discussion following it. In particular for $\mu \times \mu$ -a.e. (x, y) the scenery $(\mu_{x,t} \times \mu_{y,t})_{t \geq 0}$ equidistributes for some P -joining $P_{x,y}$. Now let $s \in \mathbb{R}$ and consider the family $(\mu_{x,t} \times \mu_{y,t+s})_{t \geq 0}$, in which we have shifted the second component by S_s^\square . This family equidistributes for the P -joining $Q = (\text{id} \times S_s^\square) P_{x,y}$ obtained as the push-forward of $P_{x,y}$ through the map $(\sigma, \tau) \mapsto (\sigma, S_s^\square \tau)$.² For $\alpha \in \Sigma_P$ and corresponding eigenfunction φ , let (σ, τ) be a $P_{x,y}$ -typical pair such

²This requires a short argument since S^\square is not continuous, but we omit it.

that $(\sigma, S_s^\square \tau)$ is Q -typical. Then

$$\begin{aligned} p_\alpha(Q) &= p_\alpha(\sigma, S_s^\square \tau) \\ &= \frac{\varphi(\sigma)}{e(\alpha s)\varphi(\tau)} \\ &= e(-\alpha s) \cdot p_\alpha(\sigma, \tau) \\ &= e(-\alpha s) \cdot p_\alpha(P_{x,y}) \end{aligned}$$

Together with Lemma 2.3, this leads to the following result:

Proposition 2.8. *Let μ be a T_a -invariant measure of intermediate dimension which a.e. generates a distribution P , let $\alpha \in \Sigma_P$ and let $f \in \text{diff}^1(\mathbb{R})$. Then $\theta_\alpha(f\mu)$ is well defined, and, fixing a μ -typical x_0 , is given up to rotation by*

$$\theta_\alpha(f\mu) = \int \delta_{e(-\alpha \log f'(y)) \cdot p_\alpha(P_{x_0,y})} d\mu(y)$$

The discussion above proves the proposition when f preserves orientation. An obvious modification of the statement and proof is needed when f is orientation-reversing. See the remark after Lemma 2.3.

2.4. Proof of the main results.

Proof (of Theorem 1.1). Let μ, ν be T_a -ergodic measures of intermediate dimension. Suppose $f \in \text{diff}^1(\mathbb{R})$ and $f\mu|_E \sim \nu|_E$ for some set E with $\nu(E) > 0$. Then $\theta_\alpha(f\mu|_E)$ and $\theta_\alpha(\nu|_E)$ are equivalent for every $\alpha \in \Sigma_P$. Choosing $n \in \mathbb{N}$ and $\alpha = \frac{n}{\log a} \in \Sigma_P$, as we may by Theorem 2.5, it follows from Theorem 2.7 that $\theta_\alpha(\nu)$ is a point mass. Therefore $\theta_\alpha(f\mu|_E)$ is a point mass, and by Proposition 2.8 this implies that $e(\alpha \cdot \log f'(\cdot))$ is μ -a.s. constant on $f^{-1}E$, giving the result.

In the case $\nu = \mu$ we have $f\mu|_E \ll \mu|_E$ so $\theta_\alpha(f\mu|_E) \ll \theta_\alpha(\mu)$, and since both consist of a single atom we have equality. Hence $e(\alpha \cdot \log f'(\cdot)) = 1$ at μ -a.e. point of $f^{-1}E$, so $f'|_{f^{-1}E}$ is μ -a.e. an integer power of $a^{1/n}$. \square

Before proving the next theorems we require one more technical result:

Proposition 2.9. *Let μ be a T_a -invariant measure of intermediate entropy generating P , and $\alpha \in \Sigma_P \cap \frac{1}{\log a}\mathbb{Z}$. Suppose one of the following holds:*

- (1) μ is ergodic and $f \in \text{diff}^2(\mathbb{R})$,
- (2) f is affine.

Then $\theta_\alpha(f\mu)$ is singular with respect to Lebesgue.

Proof. In the first case $\theta_\lambda(\mu, x_0)$ consists of a single atom (Theorem 2.7), and $p_\lambda(x_0, \cdot)$ is independent of y . Then by Proposition 2.8, up to rotation $\theta_\alpha(f\mu)$ is the image of μ under $g : x \mapsto e(f'(x))$. Since $f \in C^2$ we have $f' \in C^1$ and in

particular f' , and hence g , is Lipschitz. Since μ has intermediate dimension it is singular with respect to Lebesgue, so this is also true of $g\mu$, as desired.

In the second case $f(x) = ux + v$ and we need only consider the case $u \neq 0$. Since $f'(x)$ does not depend on x , we find by Proposition 2.8 that $\theta_\alpha(f\mu)$ is a rotation of $\theta_\alpha(\mu)$ by $e(\log u)$, and so, since θ_λ is singular by Theorem 2.7, so is $\theta_\alpha(f\mu)$. \square

Proof (of Theorem 1.4 and variants). Let μ be a T_a -invariant measure and ν a T_b -invariant measure, both of intermediate entropy. Suppose that $f \in \text{diff}^2(\mathbb{R})$ and that $(f\mu)|_E \sim \nu|_E$ for some E with $\nu(E) > 0$. Then $\theta_\lambda(f\mu|_E)$ is equivalent to $\theta_\lambda(f\nu|_E)$ and it suffices to show that this is impossible.

Assume that μ is ergodic, let P denote the distribution generated by μ , and choose $\alpha \in \Sigma_P \cap \frac{1}{\log a}\mathbb{Z}$, which is possible by Theorem 3.12. Then by Proposition 2.9, $\theta_\alpha(f\mu)$ is singular with respect to Lebesgue, while by Theorem 2.7 $\theta_\alpha(\nu)$ is absolutely continuous. Thus the two are not equivalent. Note that for this argument we did not require ergodicity of ν .

Assume instead that f is affine (but μ, ν need not be ergodic). We first disintegrate μ according to the partition of $[0, 1]$ determined by the level sets of $x \mapsto P_x$, where P_x is the distribution generated by μ at x . Since P_x depends only on $\mu^{(x)}$, this is a coarsening of the ergodic decomposition. Decompose ν similarly. By Lemma 2.3 f respects these partitions, so it suffices to prove the result for the corresponding conditional measures, which in the case of μ are T_a -invariant, and T_b -invariant in the case of ν . Hence we may assume from the start that μ, ν generate a single distribution P a.e. The result now follows as above from the second part of Corollary 2.9.

The case of C^1 -maps when μ, ν satisfy some spectral assumptions was sketched after Theorem 3.12. \square

Proof. Proof (of Corollary 1.5) Suppose $\varphi f \varphi^{-1} = T_a$ and $\psi g \psi^{-1} = T_b$ for $\varphi, \psi \in \text{diff}^2(\mathbb{R}/\mathbb{Z})$. Let μ be a common measure of positive dimension. The measures $\varphi\mu, \psi\mu$ are invariant, respectively, for T_a and T_b , and the dimension hypothesis implies that they have no ergodic component of entropy zero. Now $\psi\varphi^{-1}(\varphi\mu) = \varphi\mu$ and $\psi\varphi^{-1} \in \text{diff}^2(\mathbb{R}/\mathbb{Z})$, so Theorem 1.4 implies that $\varphi\mu$ is Lebesgue. Therefore $\psi\varphi^{-1}(\varphi\mu)$ is a T_b -invariant measure equivalent to Lebesgue, so it must be Lebesgue measure as well, and so $\psi\varphi^{-1}$ preserves Lebesgue measure. The only diffeomorphism on \mathbb{R}/\mathbb{Z} with this property is a rotation. \square

3. CONSTRUCTION AND PROPERTIES OF THE SCENERY FLOW

Throughout this section we fix an integer $b > 1$ and a non-atomic probability measure μ on $[0, 1]$ which is invariant under T_b . We write $[u; v] = [u, v] \cap \mathbb{Z}$, and similarly $[u; v]$ etc. Our convention is $\mathbb{N} = \{1, 2, 3, \dots\}$.

3.1. The extended scenery flow. For the moment fix the dimension $d = 1$ and consider measures on \mathbb{R} . We use $*$ to denote the operation of normalizing a measure on $[-1, 1]^d$, that is, if τ is a Radon measure on \mathbb{R}^d and $\tau([-1, 1]) > 0$, then

$$\tau^* = \frac{1}{\tau([-1, 1])} \tau$$

Thus $\tau^\square = (\tau^*)|_{[-1, 1]}$. Let $\mathcal{M}^* \subseteq \mathcal{M}$ denote the set of measures giving unit mass to $[-1, 1]$. Write $S_t^* : \mathcal{M}^* \rightarrow \mathcal{M}^*$ for the partially defined map

$$S_t^* \mu = (S_t \mu)^*$$

Thus $S^* = (S_t^*)_{t \in \mathbb{R}}$ is a measurable flow on the Borel subset of measures $\mu \in \mathcal{M}^*$ with 0 in their support.

While working with S^* is more natural than S^\square , we used the latter in the definition of the scenery flow because $\mathcal{P}(\mathcal{M}^*)$ does not carry a nice topology with which to define equidistribution of S^* -orbits. However there is a simple way to move between invariant distributions of the two flows. First, one may verify that $\tau \mapsto \tau^\square$ is a factor map from the measurable flow (\mathcal{M}^*, S^*) to the semi-flow $(\mathcal{M}^\square, S^\square)$, i.e. $S_t^\square(\mu^\square) = (S_t^* \mu)^\square$, and so an S^* -invariant distribution Q is pushed via $\mu \mapsto \mu^\square$ to an S^\square -invariant distribution $P = Q^\square$ called the restricted version of Q . Conversely, if P is an S^\square -invariant distribution then there is a unique S^* -invariant distribution Q on \mathcal{M}^* , called the *extended version* of P , satisfying $Q^\square = P$. The extended version may be obtained as the inverse limit of the diagram

$$\dots \xrightarrow{S_1^\square} \mathcal{M}^\square \xrightarrow{S_1^\square} \mathcal{M}^\square \xrightarrow{S_1^\square} \mathcal{M}^\square$$

(so dynamically Q is the natural extension of P). Indeed, starting from a left-infinite sequence $(\dots, \mu_{-2}, \mu_{-1}, \mu_0)$ with $S_1^\square \mu_{i+1} = \mu_i$, there is a unique measure $\mu_\infty \in \mathcal{M}^*$ such that $\mu_\infty|_{[-b^n, b^n]} = S_n^* \mu_{-n}$, and the induced distribution Q on these measures is seen to be S^* -invariant and satisfy $Q^\square = P$.

We shall usually not make the distinction between the extended and restricted versions of these flows. We note for later use that they have the same pure point spectrum.

3.2. The scenery flow of a T_a -invariant measure. In this section we construct a flow associated to a T_b -invariant measure, and study its properties. Let $\Omega = \Omega_b = \{0, \dots, b-1\}^{\mathbb{Z}}$, and denote the shift map on Ω by T , i.e. $(T\omega)_i = \omega_{i+1}$. We write ω_I for the subsequence $(\omega_i)_{i \in I}$.

Let $\xi_k : \Omega \rightarrow [0, b^{-k}]$ denote the map

$$\xi_k(\omega) = \sum_{i=k+1}^{\infty} b^{-i} \omega_i$$

In particular we write

$$\xi = \xi_0$$

which is the base- b coding map taking $\omega \in \Omega$ to the point $x \in [0, 1]$ whose base- b expansion is $0.\omega_1\omega_2\dots$ (note that this map is everywhere uncountable-to-one since it discards the non-positive coordinates of ω).

Every non-atomic T_b -invariant measure μ lifts to a unique T -invariant measure $\tilde{\mu}$ on Ω such that $\xi\tilde{\mu} = \mu$; the system $(\tilde{\mu}, T)$ is a realization of the natural extension of (μ, T_b) . For $x \in [0, 1]$ we denote by $\mu^{(x)}$ the ergodic component of x in μ , and write $\tilde{\mu}^{(x)}$ for the unique ergodic component of $\tilde{\mu}$ which maps under ξ_0 to $\mu^{(x)}$. We also write $\tilde{\mu}^{(\omega)}$ for the ergodic component of $\omega \in \Omega$.

Recall that a measure on \mathbb{R} has exact dimension α if its local dimension exists a.e. and is a.e. equal to α .

Below we construct a map $\pi : \Omega \rightarrow \mathcal{M}$, usually denoted $\omega \mapsto \mu_\omega$, defined a.e. for every T -invariant measure on Ω , and satisfying the following properties.

Proposition 3.1. *For every T_b -invariant measure μ , the following hold:*

- (1) $\mu^{(x)}$ can be represented as

$$\mu^{(x)} = \int (U_{-\xi(\omega)}\mu_\omega)|_{[0,1]^d} d\tilde{\mu}^{(x)}(\omega)$$

and in particular

$$\mu = \int (U_{-\xi(\omega)}\mu_\omega)|_{[0,1]^d} d\tilde{\mu}(\omega)$$

- (2) The map $\pi^* : \omega \mapsto \mu_\omega^*$ intertwines the actions of T and $S_{\log b}^*$, i.e.

$$S_{\log b}^* \mu_\omega^* = \mu_{T\omega}^*$$

In particular, the distribution

$$\tilde{P}_x = \pi^* \tilde{\mu}^{(x)}$$

is $S_{\log b}^*$ -invariant, and the map

$$\pi^* : (\Omega, \tilde{\mu}^{(x)}, T) \rightarrow (\mathcal{M}^*, \tilde{P}_x, S_{\log b}^*)$$

is a factor map of discrete-time systems.

- (3) μ_ω has exact dimension $h(\tilde{\mu}^{(\omega)})/\log b$.

We now begin the construction of μ_ω . Let μ be a T_b -invariant measure. Given a left-infinite sequence $\omega_{(-\infty; k]} \in \{0, \dots, b-1\}^{(-\infty; k]}$ let $\tilde{\mu}(\cdot | \omega_{(-\infty; k]})$ be the probability measure obtained by conditioning $\tilde{\mu}$ on the set $\{\eta \in \Omega : \eta_{(-\infty; k]} = \omega_{(-\infty; k]}\}$. Note that this set can be identified, using ξ_k , with $[0, b^{-k}]$. Define the measure $\mu(\cdot | \omega_{(-\infty; k]})$ on $[0, b^{-k}]$ by pushing these conditional measure forward through ξ_k ,

i.e.

$$\mu(\cdot \mid \omega_{(-\infty; k]}) = \tilde{\mu}(\xi_k^{-1}(\cdot) \mid \omega_{(-\infty; k]})$$

Note that this definition depends only on the ergodic component $\tilde{\mu}^{(\omega)}$, rather than the pair $(\tilde{\mu}, \omega)$.

For any measurable $A \subseteq [0, b^{-k}]$ we have the relation

$$\mu(A \mid \omega_{(-\infty; k]}) = c_\omega^k \cdot \mu(A + b^{-k}\omega_k \mid \omega_{(-\infty; k-1]})$$

where c_ω^k is a normalizing constant chosen so that equality holds for $A = [0, b^{-k}]$. More generally, for $k < m$ and $A \subseteq [0, b^{-m+1}]$, we have

$$(3.1) \quad \mu(A \mid \omega_{(-\infty; m]}) = c_\omega^{k,m} \cdot \mu(A + \sum_{k < i \leq m} b^{-i}\omega_i \mid \omega_{(-\infty; k]})$$

where $c_\omega^{k,m} = c_\omega^{k+1} \cdot \dots \cdot c_\omega^m$.

It follows that the sequence of measures $\mu_{\omega, k} \in \mathcal{M}$, defined for $k \leq 0$ by

$$(3.2) \quad \mu_{\omega, k}(A) = c_\omega^{k,0} \cdot \mu(A + \sum_{i=k+1}^{\infty} b^{-i}\omega_i \mid \omega_{(-\infty, k]})$$

agree as $k \rightarrow -\infty$ on the increasing sequence of intervals

$$[-\xi_k(\omega), \xi_k(\omega) + b^{-k}] = [-\sum_{i=k+1}^{\infty} b^{-i}\omega_i, \sum_{i=k+1}^{\infty} b^{-i}\omega_i + b^{-k}]$$

and vanish outside of them. Therefore, as $k \rightarrow -\infty$ the measures $\mu_{\omega, k}$ converge to a Radon measure which we denote

$$\mu_\omega = \lim_{k \rightarrow -\infty} \mu_{\omega, k}$$

Let us now verify the properties stated in Proposition 3.1. First, from equation (3.2) we see that for $k < 0$,

$$\mu_{\omega, k}(A - \xi_0(\omega)) = c_\omega^{k,0} \cdot \mu(A + \sum_{i=k+1}^0 b^{-i}\omega_i \mid \omega_{(-\infty, k]})$$

which, from equation (3.1), implies

$$\mu_\omega(A - \xi_0(\omega)) = \mu(A \mid \omega_{(-\infty, 0]})$$

Integrating over $\omega \sim \tilde{\mu}^{(x)}$ or over $\omega \sim \tilde{\mu}$ gives part (1) of Proposition 3.1.

Note that $0 \in \text{supp } \mu_\omega$ and the relation (3.1) implies

$$(3.3) \quad \mu_{T\omega}(A) = c_\omega \mu_\omega\left(\frac{1}{b}A\right)$$

for some constant c_ω independent of A . Thus by (3.3), the map $\omega \mapsto \mu_\omega^*$ is defined $\tilde{\mu}$ -a.e. and intertwines the shift T and the scaling map $S_{\log b}^*$, i.e.

$$S_{\log b}^* \mu_\omega^* = \mu_{T\omega}^*$$

This establishes part (2) of Proposition 3.1 (the later statements in that part follow from the first).

Let

$$[\omega_1 \dots \omega_n] = \{\eta \in \Omega : \eta_1 \dots \eta_n = \omega_1 \dots \omega_n\}$$

denote the cylinder set corresponding to a finite sequence $\omega_1 \dots \omega_n$. A variant of the Shannon-McMillan-Breiman theorem states that for $\tilde{\mu}$ -a.e. ω ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \tilde{\mu}([\omega_1, \dots, \omega_n] | \omega_{(-\infty; 0]}) = h(\tilde{\mu}^{(\omega)})$$

As a consequence for $\tilde{\mu}$ -a.e. ω the measure $\tau = \mu(\cdot | \omega_{(-\infty, 0]})$ has exact dimension $h(\tilde{\mu}^{(\omega)}) / \log b$, i.e.

$$\lim_{r \searrow 0} \frac{\log \tau(B_r(x))}{\log r} = \frac{h(\tilde{\mu}^{(\omega)})}{\log b} \quad \text{at } \tau\text{-a.e. point } x$$

(what is obvious is that this limit holds when, instead of $B_r(x)$, we consider the mass of b -adic intervals containing x , since these correspond to cylinder sets; the version above follows using e.g. [16, Theorem 15.3]). The same argument also holds for $\mu(\cdot | \omega_{(-\infty, k]})$ and for any $k < 0$, hence for $\mu_{\omega, k}$, and gives the result for μ_{ω} .

As a special case, we remark that when $h(\tilde{\mu}) = 0$ all the conditional measures $\mu(\cdot | \omega_{(-\infty, k]})$ consist of a single atom, and consequently, $\mu_{\omega} = \delta_0$. Likewise, when $\tilde{\mu}$ is λ^* it is easy to verify that $\mu_{\omega} = \lambda^*$ for $\tilde{\mu}$ -a.e. ω . In these cases the flows P_x are trivial, consisting of point masses at the S^* -fixed points δ_0 or λ^* .

3.3. Convergence of the scenery to the scenery flow. We now turn to the sceneries of T_b -invariant measures and their relation to the flow constructed above. We continue to work in dimension $d = 1$ and with the notation of the previous section.

Proposition 3.2. *Let μ be a T_b -invariant measure and let $\tilde{\mu}, \tilde{P}_x$ be as in Proposition 3.1. Then at μ -a.e. x , μ generates the S^* -invariant distribution*

$$P_x = \int_0^1 S_{t \log b}^* \tilde{P}_x dt$$

In particular, the scenery flow generated by μ at x depends only on the ergodic component of x and arises as a factor of the $\log b$ -suspension of $(\Omega, \tilde{\mu}^{(x)}, T)$.

Note that, since by Proposition 2.4 \tilde{P}_x is $S_{\log b}^*$ invariant and ergodic, it follows immediately that P_x , as defined above, is S^* -invariant and ergodic and that $(\mathcal{M}^*, \tilde{P}_x, S_{\log b}^*)$ is a factor of the $\log b$ -suspension of the discrete time system $(\mathcal{M}^*, \tilde{P}_x, S_{\log b}^*)$. Since $(\mathcal{M}^*, \tilde{P}_x, S_{\log b}^*)$ is a factor of $(\Omega, \tilde{\mu}^{(x)}, T)$ by Proposition 3.1, the statement in the second part of the proposition above is automatic, and it remains only to prove the first, i.e. that μ generates P_x at x .

To this end we introduce another sequence of measures $\mu'_{\omega,k}$ on $[0, b^{-k}]$ by

$$\mu'_{\omega,k}(A) = c_{\omega}^{k,0} \cdot \tilde{\mu}(\xi_k^{-1}(A + \sum_{i=k+1}^{\infty} b^{-i}\omega_i))$$

This is the same as the definition of $\mu_{\omega,k}$ except we have not conditioned on $\omega_{(-\infty,k]}$

Proposition 3.3. $\lim_{k \rightarrow -\infty} \mu'_{\omega,k} = \mu_{\omega}$ weak-* on any compact set in \mathbb{R} .

Proof. Let $\tilde{\mu}(\cdot | \omega_{[k+1;\infty)})$ denote the conditional measure $\tilde{\mu}$ on sequences $\eta \in \Omega$ given the “future” $\eta_{[k+1;\infty)} = \omega_{[k+1;\infty)}$. Then

$$\begin{aligned} \mu'_{\omega,k} &= \int \mu_{\eta,k} d\tilde{\mu}(\eta | \omega_{[k+1;\infty)}) \\ &= \int (\mu_{\eta,k} - \mu_{\eta}) d\tilde{\mu}(\eta | \omega_{[k+1;\infty)}) + \int \mu_{\eta} d\tilde{\mu}(\eta | \omega_{[k+1;\infty)}) \end{aligned}$$

The second term in the last expression is a measure-valued martingale in the variable ω with respect to the filtration $\mathcal{F}_k \subseteq \mathcal{F}_{k-1}, \dots$, where \mathcal{F}_k is the σ -algebra generated by coordinates $k+1, k+2, \dots$. Since these algebras generate the Borel algebra on Ω , the term on the right converges $\tilde{\mu}$ -a.e. to μ_{ω} as $k \rightarrow -\infty$.

In order to deal with the first term on the right, note that if we integrate against any compactly supported function f on \mathbb{R} , for $k > k_0(\omega)$ the measures $\mu_{\eta,k}$ and $\mu_{\eta,-\infty}$ will agree on the support of f , and so the integral will vanish. Hence the first term also converges to 0 weak-* on any compact set. \square

Corollary 3.4. $\lim_{k \rightarrow -\infty} (\mu'_{\omega,k})^{\square} = (\mu_{\omega})^{\square}$ in the weak-* sense for $\tilde{\mu}$ -a.e. ω .

For the next step we rely on a classical ergodic theorem due to Maker:

Theorem 3.5 (Maker, [14]). *Let (Ω, ν, T) be a measure preserving system. Let F_n be measurable functions with $\sup_n |F_n| \in L^1$ and suppose that $F_n \rightarrow F$ a.e.. Then*

$$\frac{1}{N} \sum_{n=1}^N T^n F_n \rightarrow \mathbb{E}(F | \mathcal{E})$$

a.e., where \mathcal{E} is the σ -algebra of T -invariant sets.

Proposition 3.6. For μ -a.e. x

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \delta_{\mu_{x, n \log b}} \rightarrow \tilde{P}_x$$

Proof. Define $F_k(\omega) = \delta_{(\mu'_{\omega,-k})^{\square}}$ and $F_k(\omega) = \delta_{(\mu_{\omega})^{\square}}$. From the corollary we know that $F_k \rightarrow F$ a.s. and these distribution-valued functions are uniformly bounded in the space of distributions on $\mathcal{P}(\mathcal{M}^{\square})$. Thus by Maker’s theorem,

$$\frac{1}{N} \sum_{n=1}^N F_k(T^n \omega) \rightarrow \mathbb{E}(F | \mathcal{E})$$

$\tilde{\mu}$ -a.e. One verifies from the definitions that

$$(\mu'_{T^k \omega, k})^\square = \mu_{x, k \log k}$$

for $x = \xi_0(\omega)$. The proposition follows, since $\mathbb{E}(\delta_{(\mu_\omega)^\square} | \mathcal{E}) = \tilde{P}_x$. \square

Finally, applying the operator $\int_0^1 S_{t \log b}^\square dt$ to the limit in the proposition above, we arrive at the conclusion of Proposition 3.1 concerning continuous-time sceneries.

3.4. The multidimensional case. We now turn to the higher dimensional setting. Let $*$ denote the normalization operation $\mu \mapsto \mu^* = \frac{1}{\mu([-1, 1]^d)} \mu$ and define the associated objects as in Section 3.1.

Let μ_1, \dots, μ_d be T_b -invariant measures and write $\mu = \mu_1 \times \dots \times \mu_d$, which is a measure on $[0, 1]^d$ invariant under the diagonal map $T_b(x) = (T_b x_1, \dots, T_b x_d)$. The measure $\tilde{\mu} = \tilde{\mu}_1 \times \dots \times \tilde{\mu}_d$ on $\Omega^d \cong (\{0, \dots, b-1\}^d)^\mathbb{Z}$ is the natural extension of μ . We may define maps $\xi_k : \Omega^d \rightarrow [0, b^{-k}]^d$ by applying ξ_k coordinatewise and define a map $\omega \mapsto \mu_\omega \in \mathcal{M}_d$ using the same procedure as in dimension $d = 1$.

Let φ_i denote projection to the i -th coordinate. Let π^* and π_i^* denote the maps $\omega \mapsto \mu_\omega^*$ and $\eta \mapsto (\mu_i)_\eta^*$ respectively.

Proposition 3.7. *Let μ_1, \dots, μ_d be T_b -invariant measures on $[0, 1]$, let $\mu = \times_{i=1}^d \mu_i$, and let $\omega \mapsto \mu_\omega$ be as above. Then:*

- (1) The analogs of Propositions 3.1 and 3.2 hold.
- (2) $\mu_\omega = \times_{i=1}^d (\mu_i)_{\omega^i}$.
- (3) If μ generates \tilde{P}_x at $x \in [0, 1]^d$ and μ_i generates $\tilde{P}_{i,y}$ at y , and we identify product measures with d -tuples of measures, then the following diagram factor maps commutes:

$$\begin{array}{ccc} ([0, 1]^d, \tilde{\mu}^{(x)}, T) & \xrightarrow{\pi^*} & (\mathcal{M}^d, \tilde{P}_x, S_{\log b}^*) \\ \varphi_i \downarrow & & \downarrow \varphi_i \\ ([0, 1], \tilde{\mu}_i^{(x_i)}, T) & \xrightarrow{\pi_i^*} & (\mathcal{M}, \tilde{P}_{i,x_i}, S_{\log b}^*) \end{array}$$

Proof. The proof of (1) is the same as the 1-dimensional case.

For (2), note that the space of product measures on \mathbb{R}^d is closed and each $\mu_{x,t}$ is a product measure, so the scenery flow of a product measure is supported on product measures. Hence by part (1), μ_ω is a product measure.

In order to see that $\mu_\omega = \times_{i=1}^d \mu_{i,\omega^i}$, let Q be the $S_{\log b}^*$ -invariant distribution on pairs of measures obtained by pushing forward $\tilde{\mu}$ through the map $\omega \mapsto ((\mu_1)_{\omega^1}^*, \mu_\omega^*)$, and let Q_x be the push-forward of $\tilde{\mu}^{(x)}$ by the same map, so that $Q = \int Q_x d\tilde{\mu}(x)$. We wish to show that Q -a.e. pair $(\tau, \nu_1 \times \dots \times \nu_d)$ satisfies $\tau = \nu_1$, so we must show that for μ -a.e. x this holds for Q_x . To see this, consider for μ -typical $x \in [0, 1]^d$

the sequence

$$(\tau_n, \nu_n) = ((\mu_1)_{x_1, n \log b}, \mu_{x, n \log b})$$

$n = 1, 2, 3, \dots$, and repeat the proof of Proposition 3.2 to conclude that $\frac{1}{N} \sum_{n=1}^N \delta_{(\tau_n, \nu_n)} \rightarrow Q_x$. Since the relationship $\tau_n = \pi_1(\nu_n)$ holds for all n and this is a closed condition, it holds also for the limiting distribution Q_x .

Finally, the commutativity of the diagram is a direct result of the relationship $(\mu_i)_{\omega^i} = \pi_i(\mu_\omega)$. \square

3.5. Eigenvalues and ergodicity of flows. We briefly present some technical facts about flows and their spectrum and ergodicity properties. For sake of economy we present the discussion for an invariant distribution P on (\mathcal{M}^*, S^*) .

A function $f \in L^2(P)$ generally is defined only P -a.e. and hence for typical ν it is defined at $S_t^* \nu$ for only Lebesgue-a.e. t . The next lemma says that a function which behaves like an eigenfunction at a.e. point along a.e. orbit may be modified on a set of measure zero to become an eigenfunction.

Lemma 3.8. *Let $\varphi \in L^2(P)$ and suppose that for every $t \in \mathbb{R}$ we have $S_t^* \varphi = e(\alpha t) \varphi$ P -a.e. Then there exists $\bar{\varphi} \in L^2(P)$ which, for P -a.e. ν , satisfies $S_t^* \bar{\varphi}(\nu) = e(\alpha t) \bar{\varphi}(\nu)$ for every $t \in \mathbb{R}$, and $\varphi = \bar{\varphi}$ a.e.*

Proof. Define $\bar{\varphi}(\nu) = \int_0^1 e(-\alpha t) \varphi(S_t^* \nu) dt$. \square

Usually, the ergodic decomposition of a measure is defined only in an a.e. sense. For the decomposition of P with respect to $S_{t_0}^*$ we can give a more canonical description. We say that an $S_{t_0}^*$ -invariant distribution Q is an ergodic component of P (with respect to $S_{t_0}^*$) if it is ergodic for $S_{t_0}^*$ and $\int_0^1 S_{t_0+t}^* Q dt = P$. Note that if $P = \int Q_\nu dP(\nu)$ is an abstract ergodic decomposition of P with respect to $S_{t_0}^*$ then $P = \int (\int_0^1 S_{t_0+t}^* Q_\nu dt) dP(\nu)$, and each of the inner integrals is S^* -invariant. Therefore, ergodicity of P implies that for P -a.e. ν the inner integral is P , so Q_ν is an ergodic component. Hence ergodic components exist.

Lemma 3.9. *If Q and Q' are ergodic components for $S_{t_0}^*$ then $S_{t_0+r}^* Q' = Q$ for some $r \in [0, 1]$. In particular, the representations of P as $\int_0^1 S_{t_0+t}^* Q dt$ does not depend (up to a translation modulo 1 of the parameter space) on the ergodic component Q .*

Proof. Since $\int_0^1 S_{t_0+t}^* Q dt$, $\int_0^1 S_{t_0+t}^* Q' dt$ are both ergodic decompositions of P , by uniqueness of the ergodic decomposition we see that for a.e. $t \in [0, 1]$ there is an $s \in [0, 1]$ with $S_{t_0+t}^* Q = S_{t_0+s}^* Q'$. Then for $r = s - t$ (or $r = 1 + s - t$ if $s < t$), we have $Q = S_{t_0(s-t)}^* Q'$. The second statement is immediate from the first. \square

Lemma 3.10. *Let Q be an ergodic component of P with respect to $S_{t_0}^*$. Then either $Q = P$, i.e. $S_{t_0}^*$ is ergodic, or there is a largest $n \in \mathbb{N}$ such that Q is invariant under $S_{t_0/n}^*$.*

Proof. Consider the map $q : \mathbb{R}/\mathbb{Z} \rightarrow \mathcal{P}(\mathcal{M})$ given by $q(t) = S_{t_0 \cdot t}^* Q$, which is well defined since $S_{t_0}^* Q = Q$. Let $\Lambda \subseteq \mathbb{R}/\mathbb{Z}$ denote the set of periods of q , that is $r \in \Lambda$ if $q(t+r) = q(t)$ for $t \in \mathbb{R}/\mathbb{Z}$, or equivalently, such that $q(r) = q(0)$. Since q is measurable with respect to Lebesgue measure, either Λ is discrete or $\Lambda = \mathbb{R}/\mathbb{Z}$. In the latter case, since $P = \int_{\mathbb{R}/\mathbb{Z}} q(t) dt = q(0) = Q$. In the former case Λ has the form $\{\frac{k}{n} : 0 \leq k < n\}$ for some n , and this is the n we are looking for. \square

Lemma 3.11. *For $t_0 > 0$ the following are equivalent:*

- (1) $\frac{1}{t_0} \in \Sigma_P$.
- (2) *For some (equivalently every) ergodic component Q of $S_{t_0}^*$, the flow (\mathcal{M}^*, P, S^*) is isomorphic to the t_0 -suspension of $(\mathcal{M}^*, Q, S_{t_0}^*)$.*
- (3) *$S_{t_0}^*$ is not ergodic, and its ergodic components are not preserved under $S_{t_0/n}^*$ for any $n \in \mathbb{N}$.*

Proof (sketch). (1) \implies (2): If φ is an eigenfunction for $\frac{1}{t_0}$ then one may verify that the ergodic components of $S_{t_0}^*$ are precisely the conditional distributions of P on the level sets of φ . Fixing an ergodic component Q supported on a level set $\varphi^{-1}(z)$ define $r : \mathcal{M}^* \rightarrow [0, 1)$ by $e(t_0 r(\nu)) = z$. Then $\nu \mapsto (S_{-r(\nu)}^* \nu, \varphi(\nu))$ is an isomorphism of (\mathcal{M}^*, P, S^*) and the t_0 -suspension of $(\mathcal{M}^*, Q, S_{t_0}^*)$.

(2) \implies (3): Trivial since e.g. the subset of \mathcal{M}^* corresponding to $\mathcal{M}^* \times [0, \frac{t_0}{2})$ in the suspension is $S_{t_0}^*$ -invariant, but not $S_{t_0/n}^*$ -invariant for any $1 \neq n \in \mathbb{N}$.

(3) \implies (1): By the previous lemma we find that the action of S^* on the ergodic components for $S_{t_0}^*$ is isomorphic to $[0, t_0)$ with addition modulo t_0 . Since P -a.e. point belongs to a well defined ergodic component, this gives an eigenfunction with eigenvalue $\frac{1}{t_0}$. \square

3.6. The spectrum of $(\mathcal{M}^*, P_x, S^*)$. In this section we prove Theorem 2.5. Let μ be T_b -ergodic with entropy strictly between 0 and $\log b$. Recall the construction and notation from Section 3.2: specifically $(\Omega, \tilde{\mu}, T)$ is the natural extension of $([0, 1], \mu, T_b)$, the image of $\tilde{\mu}$ under $\omega \mapsto \mu_\omega$ is the $S_{\log b}^*$ -ergodic distribution \tilde{P} (it does not depend on x or ω , as it did in previous sections, because μ and $\tilde{\mu}$ are now ergodic), and $P = \int_0^1 S_{t \log b}^* \tilde{P} dt$ is the distribution of the scenery flow of μ .

Recall that the lower Hausdorff dimension of a measure τ is

$$\underline{\dim} \tau = \inf\{\dim A : \tau(A) > 0\}$$

We note that if τ has exact dimension α then $\underline{\dim} \tau = \alpha$ as well.

It is simple to verify that $\tau \ll \tau'$ implies $\underline{\dim} \tau \geq \underline{\dim} \tau'$ and, more generally, if a measure τ can be written as $\tau = \int \tau_i d\sigma(i)$, then $\underline{\dim} \tau \geq \text{essinf}_{i \sim \sigma} \underline{\dim} \tau_i$. Thus if f is a map then $f\tau = \int f\tau_i d\sigma(i)$, and a similar bound applies.

We are out to show that

$$\frac{n}{\log a} \mathbb{Z} \subseteq \Sigma_{P_x} \subseteq \frac{1}{\log a} \Sigma_{\mu^{(x)}} \cup \frac{n}{\log a} \mathbb{Z}$$

The right hand side follows from the fact that $(\mathcal{M}^*, P_x, S^*)$ is a factor of the log b -suspension of $([0, 1], \mu, T_b)$. To establish the left hand inclusion it suffices to prove the following theorem.

Theorem 3.12. *There exists an $n \in \mathbb{N}$ such that $\frac{n}{\log a} \in \Sigma_{(\mathcal{M}^*, P, S^*)}$.*

Proof. Since \tilde{P} is an ergodic component of P with respect to $S_{\log a}^*$, by Lemma 3.10 and 3.11 it suffices to show that \tilde{P} is not S^* -invariant. Suppose that it were S^* -invariant. We claim that this implies that μ is Lebesgue, contradicting the assumption of intermediate dimension.

To this end, choose an integer d such that $d \dim \mu > 1$ and write μ^{*d} for the d -fold convolution of μ , which is the image of the d -fold product $\times_{i=1}^d \mu$ by the map $f(x) = \sum_{i=1}^d x_i$. We first show that μ^{*d} has dimension 1. Recall that by Proposition 3.1,

$$\mu = \int U_{-\xi(\omega)} \mu_\omega d\tilde{\mu}(\omega)$$

Since \tilde{P} is S^* -invariant, there is a function $\xi(x, t) \in [0, 1]$ such that

$$(3.4) \quad \mu = \int_0^1 \int_0^1 (U_{-\xi(\omega, t)} S_{t \log b} \mu_\omega)|_{[0, 1]} dt \tilde{\mu}(\omega)$$

This gives a similar representation of the product measure: write $t = (t_1, \dots, t_d)$ and u^d for uniform measure on $[0, 1]^d$, and likewise write $\omega = (\omega^1, \dots, \omega^d)$ and $\tilde{\mu}^d = \times_{i=1}^d \tilde{\mu}$. Then

$$\times_{i=1}^d \mu = \int_{\Omega^d} \int_{[0, 1]^d} \times_{i=1}^d ((U_{-\xi(\omega^i, t^i)} S_{t_i \log b} \mu_{\omega^i}^*)|_{[0, 1]}) du^d(t) d\tilde{\mu}^d(\omega)$$

Therefore, using the comments preceding the proposition,

$$\underline{\dim} \mu^{*d} \geq \operatorname{essinf}_{\omega \sim \tilde{\mu}^d, t \sim u^d} f(\times_{i=1}^d (U_{-\xi(\omega^i, t^i)} S_{t_i \log b} \mu_{\omega^i}^*)|_{[0, 1]})$$

Since f is linear and $\xi(\omega^i, t^i) \in [0, 1]$, we have

$$\geq \operatorname{essinf}_{\omega \sim \tilde{\mu}^d, t \sim u^d} \underline{\dim} f(\times_{i=1}^d (S_{t_i \log b} \mu_\omega)|_{[-1, 1]})$$

because each of the previous measures is absolutely continuous with respect to the corresponding measure above. Writing $f_t(x) = \sum t_i x_i$, we have by another absolute-continuity argument

$$\geq \operatorname{essinf}_{\omega \sim \tilde{\mu}^d} \left(\operatorname{essinf}_{t \sim u^d} \underline{\dim} f_t(\times_{i=1}^d \mu_{\omega^i})|_{[-b, 2b]} \right)$$

Finally, for fixed typical ω we have

$$\dim \times_{i=1}^d \mu_{\omega^i} = \sum_{i=1}^d \dim \mu_{\omega^i} > 1$$

so the inner essinf (over $t \sim u^d$) in the previous expression is 1 by the following version of Mastrand's classical theorem on projections of measures.

Theorem 3.13 (Hunt-Kaloshin [9]). *Let σ be an exact dimensional probability measure on \mathbb{R}^d with $\dim \sigma = \alpha$. Then for Lebesgue-a.e. $(t_1, \dots, t_d) \in \mathbb{R}^d$, the image of σ under $x \mapsto \sum_{i=1}^d t_i x_i$ is exact dimensional and has dimension $\min\{1, \dim \sigma\}$.*

Thus, we have shown that μ^{*d} has dimension 1. Next, note that convolution in \mathbb{R}/\mathbb{Z} is obtained by taking the convolution in \mathbb{R} modulo 1. Since this is a countable-to-1 local isometry $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z}$ it does not change dimension, so the d -th convolution of μ in \mathbb{R}/\mathbb{Z} has dimension 1. Since this convolved measure is also T_b -invariant it is exact dimensional, and hence its exact dimension is 1, and it must be Lebesgue measure because this is the only measure of dimension 1 invariant under T_b . Finally, by examining the Fourier coefficients and using the elementary relation $(\widehat{\mu^{*d}})(k) = \widehat{\mu}(k)^d$, we conclude that μ is Lebesgue measure. This is the desired contradiction. \square

3.7. The phase. Let μ be a T_b -invariant measure of intermediate dimension generating a.e. the same distribution P . As usual we denote by P_x the distribution generated by μ at x and by $P_{x,y}$ the distribution generated by $\mu \times \mu$ at (x, y) , and by \tilde{P}_x and $\tilde{P}_{x,y}$ distributions obtained from $\tilde{\mu}$ and $\tilde{\mu} \times \tilde{\mu}$ as in Proposition 3.1.

Throughout this section $\alpha \in \Sigma_P$ and φ is the corresponding eigenvalue. Recall that the phase $p_\alpha(P_{x,y})$ is the almost sure value of $\varphi(\sigma)/\varphi(y)$ for $\sigma \times \tau \sim P_{x,y}$, which is the same as for $\sigma \times \tau \sim \tilde{P}_{x,y}$. Therefore for $\mu \times \mu$ -typical (x, y) and corresponding typical $(\omega, \eta) \in \Omega \times \Omega$,

$$p_\alpha(P_{x,y}) = \frac{\varphi(\mu_\omega^*)}{\varphi(\mu_\eta^*)}$$

Fixing a μ -typical x_0 , the phase measure $\theta_\alpha = \theta_\alpha(\mu, x_0)$ is the push-forward of μ via $y \mapsto p_\alpha(P_{x_0,y})$. Thus, for ω_0 corresponding to x_0 , we find that $\theta_\alpha(\mu, x_0)$ is the push-forward of $\tilde{\mu}$ through the map $\eta \mapsto \varphi(\mu_{\omega_0}^*)/\varphi(\mu_\eta^*)$.

Proposition 3.14. *If $\lambda \in \Sigma_P \setminus \frac{1}{\log b} \mathbb{Q}$ then θ_λ is Lebesgue measure.*

Proof. $\eta \mapsto \varphi(\mu_\eta^*)$ is an eigenfunction for the system $(\Omega, \tilde{\mu}, T)$ with eigenvalue $\lambda \log b$. Since $\lambda \log b$ is irrational the distribution of $\varphi(\mu_\eta^*)$ is uniform on the circle, so the same is true for $\varphi(\mu_{\omega_0}^*)/\varphi(\mu_\eta^*)$, and the conclusion follows. \square

Proposition 3.15. *If $\alpha = \frac{n}{m \log b} \in \Sigma_{(\mathcal{M}^*, P, S^*)}$ and μ is ergodic then θ_α is uniform measure on a rotation of the m -th roots of unity. In particular if $m = 1$ then θ_α consists of a single atom.*

Proof. $\eta \mapsto \varphi(\mu_\eta^*)$ is an eigenfunction of $(\Omega, \tilde{\mu}, T)$ with eigenvalue $(\log b)\alpha = n/m$. The distribution of $\varphi(\mu_\eta^*)$ for $\eta \sim \tilde{\mu}$ is just the distribution of this eigenfunction, which is uniform on a rotation of the m -th roots of unity. Therefore the same is true for $\varphi(\mu_{\omega_0}^*)/\varphi(\mu_\eta^*)$, $\eta \sim \tilde{\mu}$, proving the proposition. \square

Now we turn to the non-ergodic case. For $z = e(t)$ with $t \in [0, 1)$ let us denote $L(z) = t/\alpha$.

Lemma 3.16. *The distribution of μ_ω^* , $\omega \sim \tilde{\mu}$ is the same as the distribution of $S_{L(z)}^* \nu$, $(\nu, z) \sim \tilde{P}_{x_0} \times \theta_\alpha$, where $\theta_\alpha = \theta_\alpha(\mu, x_0)$.*

Proof. Write for brevity

$$p(\omega) = \frac{\varphi(\mu_{\omega_0}^*)}{\varphi(\mu_\omega^*)}$$

and consider the map $\Omega \rightarrow \mathcal{M}^* \times \{|z| = 1\}$ defined by

$$\omega \mapsto (S_{-L(p(\omega))}^* \mu_\omega^*, p(\omega))$$

It suffices to show that this map takes $\tilde{\mu}$ to $\tilde{P} \times \theta_\alpha$, and for this we must show that (a) the second component of the image measure is θ_α and (b) conditioned on the value of the second component, the distribution of the first component is \tilde{P}_{x_0} .

For (a), for $x \sim \mu$. Then for $\tilde{\mu}^{(x)}$ -a.e. ω the value of $p(\omega) = \varphi(\mu_{\omega_0}^*)/\varphi(\mu_\omega^*)$ is $p_\alpha(\tilde{P}_{x_0, x})$, because, by definition, $\mu_{\omega_0}^* \times \mu_\omega^*$ is a typical element of $\tilde{P}_{x_0, x}$; and this is the same as $p_\alpha(P_{x_0, x})$, because $P_{x_0, x} = \int_0^1 S_{t \log b}^* \tilde{P}_{x_0, x} dt$. The distribution of $p_\alpha(P_{x_0, x})$ this quantity for $x \sim \mu$ is by definition equal to θ_α .

Next, conditioned on the value $p(\omega) = \varphi(\mu_{\omega_0}^*)/\varphi(\mu_\omega^*)$ we know by Corollary that $S_{-L(z)}^* \tilde{P}_x = \tilde{P}_{x_0}$. This proves the lemma. \square

Proposition 3.17. *If $\alpha \in \Sigma_P \cap \frac{1}{\log b} \mathbb{Q}$ then θ_α is singular with respect to Lebesgue.*

Proof. Our strategy is similar to the proof of Theorem 3.12. Choose an integer d with $d \cdot \dim \mu > 1$; we show that if θ_α were continuous with respect to Lebesgue measure then this would imply that $\dim \mu^{*d} = 1$, which would contradict the to intermediate entropy of μ , as in Theorem 3.12.

We aim to show that $\dim \mu^{*d} = 1$. By Proposition 3.1 we have

$$\mu = \int (U_{\xi(\omega)} \mu_\omega) |_{[0,1]} d\tilde{\mu}(\omega)$$

Write $\omega = (\omega^1, \dots, \omega^d)$ and $\tilde{\mu}^d = \times_{i=1}^d \tilde{\mu}$. Let $f(x) = \sum_{i=1}^d x_i$. Then

$$\begin{aligned} \underline{\dim} \mu^{*d} &= \underline{\dim} f \left(\int \times_{i=1}^d (U_{\xi(\omega^i)} \mu_{\omega^i})|_{[0,1]} d\tilde{\mu}^d(\omega) \right) \\ &\geq \operatorname{ess\,inf}_{\omega \sim \tilde{\mu}^d} \underline{\dim} f \left((\times_{i=1}^d U_{\xi(\omega^i)} \mu_{\omega^i})|_{[0,1]^d} \right) \\ &\geq \operatorname{ess\,inf}_{\omega \sim \tilde{\mu}^d} \underline{\dim} f \left((\times_{i=1}^d \mu_{\omega^i})|_{[-1,1]^d} \right) \end{aligned}$$

where in the last equality we used linearity of f . Writing $\nu = (\nu_1, \dots, \nu_d)$, $z = (z_1, \dots, z_d)$ and $\tilde{P}^d, \theta_\alpha^d$ for the d -fold product measures, we can apply the previous lemma and get

$$= \operatorname{ess\,inf}_{\nu \sim \tilde{P}_{x_0}, t \sim \theta_\alpha^d} \underline{\dim} f \left((\times_{i=1}^d S_{L(z^i)}^* \nu_i)|_{[-1,1]^d} \right)$$

Setting $f_t(x) = \sum_{i=1}^d e^{t_i} x_i$ and $L(z) = (L(z_1), \dots, L(z_d))$,

$$\geq \operatorname{ess\,inf}_{\nu \sim \tilde{P}_{x_0}, t \sim \theta_\alpha^d} \underline{\dim} f_{-L(z)} \left((\times_{i=1}^d \nu_i)|_{[-b,b]^d} \right)$$

Now with ν_1, \dots, ν_d fixed typical measures for \tilde{P}_{x_0} we know that $\times_{i=1}^d \nu_i|_{[-b,b]^d}$ has exact dimension $d \dim \mu > 1$. Also, since L is a piecewise smooth map, if θ_α were absolutely continuous then the distribution of $L(z)$ for $z \sim \theta_\alpha^d$ would be absolutely continuous with respect to d -dimensional Lebesgue measure. Hence, applying Marstrand's theorem again, we find that for $\tilde{P}_{x_0}^d$ -a.e. choice of ν the dimension in the expression above is 1, so the essential infimum is 1. This completes the proof. \square

REFERENCES

- [1] Tim Bedford and Albert M. Fisher. Ratio geometry, rigidity and the scenery process for hyperbolic Cantor sets. *Ergodic Theory Dynam. Systems*, 17(3):531–564, 1997.
- [2] Daryl Cooper and Thea Pignataro. On the shape of Cantor sets. *J. Differential Geom.*, 28(2):203–221, 1988.
- [3] Márton Elekes, Tamás Keleti, and Andra Mathe. Self-similar and self-affine sets: measure of the intersection of two copies. *Ergodic Theory Dynam. Sys.* To appear.
- [4] Harry Furstenberg. Intersections of Cantor sets and transversality of semigroups. In *Problems in analysis (Sympos. Salomon Bochner, Princeton Univ., Princeton, N.J., 1969)*, pages 41–59. Princeton Univ. Press, Princeton, N.J., 1970.
- [5] Matan Gavish. Measures with uniform scaling scenery. *Preprint*, 2009.
- [6] Michael Hochman. Dynamics on fractals and fractal distributions. *Preprint*, 2009.
- [7] Michael Hochman and Pablo Shmerkin. local entropy and dimension of projections. *preprint*.
- [8] Bernard Host. Nombres normaux, entropie, translations. *Israel J. Math.*, 91(1-3):419–428, 1995.
- [9] Brian R. Hunt and Vadim Yu. Kaloshin. How projections affect the dimension spectrum of fractal measures. *Nonlinearity*, 10(5):1031–1046, 1997.
- [10] Aimee Johnson and Daniel J. Rudolph. Convergence under \times_q of \times_p invariant measures on the circle. *Adv. Math.*, 115(1):117–140, 1995.

- [11] Aimee S. A. Johnson. Measures on the circle invariant under multiplication by a nonlacunary subsemigroup of the integers. *Israel J. Math.*, 77(1-2):211–240, 1992.
- [12] Aimee S. A. Johnson and Daniel J. Rudolph. Commuting endomorphisms of the circle. *Ergodic Theory Dynam. Systems*, 12(4):743–748, 1992.
- [13] Elon Lindenstrauss. Rigidity of multiparameter actions. *Israel J. Math.*, 149:199–226, 2005. Probability in mathematics.
- [14] Philip T. Maker. The ergodic theorem for a sequence of functions. *Duke Math. J.*, 6:27–30, 1940.
- [15] Pertti Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [16] Yakov B. Pesin. *Dimension theory in dynamical systems*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1997. Contemporary views and applications.
- [17] Daniel J. Rudolph. $\times 2$ and $\times 3$ invariant measures and entropy. *Ergodic Theory Dynam. Systems*, 10(2):395–406, 1990.
- [18] Peter Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [19] Li-Feng Xi. Lipschitz equivalence of self-conformal sets. *J. London Math. Soc. (2)*, 70(2):369–382, 2004.