

EXAMPLES OF NON-POLYGONAL LIMIT SHAPES IN I.I.D. FIRST-PASSAGE PERCOLATION AND INFINITE COEXISTENCE IN SPATIAL GROWTH MODELS

MICHAEL DAMRON AND MICHAEL HOCHMAN

ABSTRACT. We construct an edge-weight distribution for i.i.d. first-passage percolation on \mathbb{Z}^2 whose limit shape is not a polygon and has extreme points which are arbitrarily dense in the boundary. Consequently, the associated Richardson-type growth model can support coexistence of a countably infinite number of distinct species, and the graph of infection has infinitely many ends.

1. INTRODUCTION

Throughout this note μ denotes a Borel probability measure on \mathbb{R}^+ with finite mean, and \mathcal{M} is the family of such measures. Let \mathbb{E} denote the set of nearest-neighbor edges of the lattice \mathbb{Z}^2 , and let $\{\tau_e : e \in \mathbb{E}\}$ be a family of i.i.d. random variables with marginal μ and joint distribution $\mathbb{P} = \mu^{\mathbb{E}}$. The *passage time* of a path $\gamma = (e_1, \dots, e_n) \in \mathbb{E}^n$ in the graph $(\mathbb{Z}^2, \mathbb{E})$ is $\tau(\gamma) = \sum_{i=1}^n \tau_{e_i}$, and for $x, y \in \mathbb{Z}^2$ the *passage time* from x to y is

$$\tau(x, y) = \min_{\gamma} \tau(\gamma),$$

where the minimum is over all paths γ joining x to y . A minimizing path is called a *geodesic* from x to y .

The theory of first passage percolation (FPP) is concerned with the large-scale geometry of the metric space (\mathbb{Z}^2, τ) . The following fundamental result concerns the asymptotic geometry of balls. Write $B(t) = \{x \in \mathbb{Z}^2 : \tau(0, x) \leq t\}$ for the ball of radius t at the origin, and for $S \subseteq \mathbb{R}^2$ and $a \geq 0$, write $aS = \{ax : x \in S\}$.

Theorem 1.1 (Cox-Durrett [1]). *For each $\mu \in \mathcal{M}$ there exists a deterministic, convex, compact set B_μ such that for any $\varepsilon > 0$,*

$$\mathbb{P} \left((1 - \varepsilon)B_\mu \subseteq \frac{1}{t}B(t) \subseteq (1 + \varepsilon)B_\mu \text{ for all large } t \right) = 1.$$

Little is known about the geometry of B_μ , which is called the *limit shape*. It is conjectured to be strictly convex when μ is non-atomic, and non-polygonal in all but the most degenerate cases, but, in fact, there are currently no known examples of μ for which these properties are verified (see [9]). For a compact, convex set $C \subseteq \mathbb{R}^2$ write $\text{ext}(C)$ for the set of extreme points and $\text{sides}(C) = |\text{ext}(C)|$, so that C is a polygon if and only if $\text{sides}(C) < \infty$. The best result to date, due to Marchand [7], is that under mild

M.D. supported by an NSF Postdoctoral Fellowship. M.H. supported by NSF grant 0901534.

assumptions, $\text{sides}(B_\mu) \geq 8$. Building on results of Marchand, our purpose of this note is to give the first examples of distributions for which the limit shape is not a polygon.

Theorem 1.2. *For every $\varepsilon > 0$ there exists $\mu \in \mathcal{M}$ (with atoms) such that B_μ is not a polygon, i.e., $\text{sides}(B_\mu) = \infty$, and $\text{ext}(B_\mu)$ is ε -dense in ∂B_μ . There exist continuous μ such that $\text{sides}(B_\mu) > 1/\varepsilon$ and $\text{ext}(B_\mu)$ is ε -dense in ∂B_μ .*

It is tempting to try to obtain a strictly convex limit shape by taking a limit of measures μ_n such that B_{μ_n} have progressively denser sets of extreme points, but unfortunately the limit one gets in our example is the unit ball of ℓ^1 .

We also obtain examples of measures μ such that, at the points $v \in \text{ext}(B_\mu)$ which lie on the boundary of the ℓ^1 -unit ball, ∂B_μ is infinitely differentiable. This should be compared with the work of Zhang [10], where such behavior was ruled out for certain μ .

Theorem 1.2 has implications for the Richardson growth model, whose definition we recall next. Fix $x_1, \dots, x_k \in \mathbb{Z}^2$ and imagine that at time 0 the site x_i is inhabited by a species of type i . Each species spreads at unit speed, taking time τ_e to cross an edge $e \in \mathbb{E}$. An uninhabited site is exclusively and permanently colonized by the first species that reaches it, i.e., $y \in \mathbb{Z}^2$ is occupied at time t by the i -th species if $\tau(y, x_i) \leq t$ and $\tau(y, x_i) < \tau(y, x_j)$ for all $j \neq i$. This is well-defined when there are *unique geodesics*, i.e., \mathbb{P} -a.s. no two paths have the same passage time, as is the case when μ is continuous, but we shall also want to consider measures μ with atoms, so we require a mechanism to break ties. For simplicity we introduce a worst-case model: if two species $i \neq j$ reach an unoccupied site x at the same instant then x is colonized by a species of type -1 , which spreads according to the same rules as the others. Under this convention if a site is occupied by the species $i \neq -1$ then it would be so occupied under any other tie-breaking rule.

Given initial sites x_1, \dots, x_k , consider the set colonized by the i -th species:

$$C_i = \{y \in \mathbb{Z}^2 : y \text{ is eventually occupied by } i\}.$$

One says that μ *admits coexistence of k species* if for some choice of x_1, \dots, x_k ,

$$\mathbb{P}(|C_i| = \infty \text{ for all } i = 1, \dots, k) > 0.$$

Coexistence of infinitely many species is defined similarly.

It is not known, even in simple examples, how many species can coexist. When μ is the exponential distribution, Häggström and Pemantle [5] proved coexistence of 2 species, and for a broad class translation-invariant measures on $(0, \infty)^\mathbb{E}$, including some non-i.i.d. ones, Hoffman [6] demonstrated coexistence of 8 species by establishing a relation with the number of sides of the limit shape in the associated FPP. Using the same relation we obtain the following:

Theorem 1.3. *There exists $\mu \in \mathcal{M}$ (with atoms) which admits coexistence of infinitely many species. For each k there exist continuous $\mu \in \mathcal{M}$ admitting coexistence of k species.*

Finally, the *graph of infection* $K \subseteq \mathbb{E}$ is the union over $x \in \mathbb{Z}^d$ of the edges of geodesics from 0 to x . If μ is continuous this is a.s. a tree. A graph has m ends if, after removing a finite set of vertices, the induced graph contains at least m infinite connected components, and, if there are m ends for every $m \in \mathbb{N}$, we say there are infinitely many ends. Newman [8] has conjectured for a broad class of μ that K has infinitely many ends. Hoffman [6] showed for continuous distributions that in general there are a.s. at least 4 ends.

Theorem 1.4. *There exist $\mu \in \mathcal{M}$ (with atoms) such that \mathbb{P} -a.s., K has infinitely many ends. For each k there exist continuous $\mu \in \mathcal{M}$ such that \mathbb{P} -a.s., K has at least k ends.*

When μ is continuous Theorems 1.3 and 1.4 follow, respectively, from Theorem 1.2 and from Theorems 1.4 and 1.6 of Hoffman [6]. For the non-continuous case we provide the necessary modifications of Hoffman’s arguments in Section 4.

2. BACKGROUND ON THE LIMIT SHAPE

Endow \mathcal{M} with the topology of weak convergence and for convenience fix a compatible metric $d(\cdot, \cdot)$ on \mathcal{M} . Next, fix the ℓ^1 -metric on \mathbb{R}^2 , and write $A^{(\varepsilon)}$ for the ε -neighborhood of $A \subseteq \mathbb{R}$. Let \mathcal{C} denote the space of non-empty, closed, convex subsets of \mathbb{R}^2 endowed with the Hausdorff metric d_H :

$$d_H(A, B) = \inf\{\varepsilon : A \subseteq B^{(\varepsilon)} \text{ and } B \subseteq A^{(\varepsilon)}\}.$$

Theorem 2.1 (Cox-Kesten [2]). *The map $\mu \mapsto B_\mu$ from \mathcal{M} to \mathcal{C} is continuous.*

It is elementary to verify that for $A \in \mathcal{C}$, the map $A \mapsto \text{ext}(A)$ is semi-continuous in the sense that, given $x \in \text{ext}(A)$ and $\varepsilon > 0$, there is a $\delta > 0$ such that if $A' \in \mathcal{C}$ and $d_H(A, A') < \delta$ then there exists $x' \in \text{ext}(A')$ with $\|x - x'\|_1 < \varepsilon$. Combined with the continuity theorem above, we have:

Corollary 2.2. *Let $\mu \in \mathcal{M}$. For every $x_1, \dots, x_k \in \text{ext}(B_\mu)$ and $\varepsilon > 0$ there is a $\delta > 0$ such that, if $\nu \in \mathcal{M}$ and $d(\nu, \mu) < \delta$ then there are $y_1, \dots, y_k \in \text{ext}(B_\nu)$ such that $\|x_i - y_i\| < \varepsilon$ for $i = 1, \dots, k$.*

We next recall some results about limit shapes for a special class of measures. Given $0 < p < 1$, let $\mathcal{M}_p \subseteq \mathcal{M}$ denote the set of measures $\mu \in \mathcal{M}$ with an atom of mass p located at $x = 1$, i.e., $\mu(\{1\}) = p$, and no mass to the left of 1, i.e., $\mu((-\infty, 1)) = 0$. Limit shapes for μ of this form were first studied in Durrett and Liggett [4]. Writing \vec{p}_c for the critical parameter of oriented percolation on \mathbb{Z}^2 (see Durrett [3] for background), it was shown that when $p > \vec{p}_c$ and $\mu \in \mathcal{M}_p$, the limit shape B_μ contains a “flat edge”, or, more precisely, ∂B_μ has sides which lie on the boundary of the ℓ^1 -unit ball. The nature of this edge was fully characterized in [7]. For $p \geq \vec{p}_c$, let α_p be the asymptotic speed of super-critical oriented percolation on \mathbb{Z}^2 with parameter p (see [3]). Define points

$w_p, w'_p \in \mathbb{R}^2$ by

$$\begin{aligned} w_p &= (1/2 + \alpha_p/\sqrt{2}, 1/2 - \alpha_p/\sqrt{2}), \\ w'_p &= (1/2 - \alpha_p/\sqrt{2}, 1/2 + \alpha_p/\sqrt{2}). \end{aligned}$$

Let $[w_p, w'_p] \subseteq \mathbb{R}^2$ denote the line segment with endpoints w_p and w'_p . It will be important to note that α_p is strictly increasing in $p > \vec{p}_c$, so the same is true of $[w_p, w'_p]$.

Theorem 2.3 (Marchand [7]). *Let $\mu \in \mathcal{M}_p$. Then*

- (1) $B_\mu \subseteq \{x \in \mathbb{R}^2 : \|x\|_1 \leq 1\}$.
- (2) If $p < \vec{p}_c$, then $B_\mu \subseteq \{x \in \mathbb{R}^2 : \|x\|_1 < 1\}$.
- (3) If $p > \vec{p}_c$, then $B_\mu \cap [0, \infty)^2 = [w_p, w'_p]$.
- (4) If $p = \vec{p}_c$, then $B_\mu \cap [0, \infty)^2 = \{(1/2, 1/2)\}$.

As noted by Marchand, this implies $\text{sides}(B_\mu) \geq 8$ for $\mu \in \mathcal{M}_p$ and $\vec{p}_c < p < 1$, since w_p, w'_p and their reflections about the axes are extreme points.

3. PROOF OF THEOREM 1.2

Our aim is to construct a $\mu \in \mathcal{M}$ with $\text{sides}(B_\mu) = \infty$. Fix any $p_0 > \vec{p}_c$, $\mu_0 \in \mathcal{M}_{p_0}$ and $\delta_0 > 0$. We will inductively define a sequence $p_1 > p_2 > \dots > \vec{p}_c$, measures $\mu_1 \in \mathcal{M}_{p_1}$, $\mu_2 \in \mathcal{M}_{p_2}, \dots$, and $\delta_1, \delta_2, \dots > 0$ such that for every $n \geq 0$ and all $k \leq n$,

- (1) If $\nu \in \mathcal{M}$ and $d(\nu, \mu_k) < \delta_k$ then $\text{sides}(B_\nu) \geq k$, and
- (2) $d(\mu_k, \mu_n) < \frac{1}{2}\delta_k$.

Note that (1) implies that $\text{sides}(B_{\mu_k}) \geq k$. Assuming p_k, μ_k and δ_k are defined for $k \leq n$, we define them for $n+1$. Fix $p_{n+1} \in (\vec{p}_c, p_n)$ and set $r = p_n - p_{n+1} > 0$. For $y > 1$ construct μ_{n+1}^y from μ_n by moving an amount r of mass from the atom at 1 to y , that is,

$$\mu_{n+1}^y = \mu_n - r\delta_1 + r\delta_y.$$

We claim that for small enough $y > 1$ and any sufficiently small choice of $\delta_{n+1} > 0$ the measure $\mu_{n+1} = \mu_{n+1}^y$ has the desired properties. First, $\mu_{n+1}^y \rightarrow \mu_n$ weakly as $y \downarrow 1$, so, since μ_n satisfies (2), so does μ_{n+1}^y for all sufficiently small y .

Second, we claim that $\text{sides}(B_{\mu_{n+1}^y}) \geq n+1$ for y close enough to 1. Indeed, since $r > 0$ we have $w_{p_{n+1}} \neq w_{p_n}$. Using (1), choose n extreme points $x_1, \dots, x_n \in \text{ext}(B_{\mu_n})$ and let

$$a = \min \{ \|x_i - x_j\|_1, \|x_i - w_{p_{n+1}}\|_1 : i \neq j \}.$$

Note that $a > 0$ by Marchand's theorem. By Corollary 2.2, for y close enough to 1, for each $i = 1, \dots, n$ we can choose an extreme point x'_i of $B_{\mu_{n+1}^y}$ with $\|x'_i - x_i\| < a/2$. By Marchand's theorem, $B_{\mu_{n+1}^y}$ also has an extreme point at $w_{p_{n+1}}$. By definition of a , these extreme points are distinct, giving $\text{sides}(B_{\mu_{n+1}^y}) \geq n+1$.

Finally, by Corollary 2.2, μ_{n+1}^y satisfies (1) for any sufficiently small choice of δ_{n+1} .

Let μ be a weak limit of μ_n . Then $d(\mu, \mu_n) \leq \frac{1}{2}\delta_n$ for all n , so by (2), $\text{sides}(B_\mu) = \infty$.

At each step, rather than creating a new atom at y , one can instead add, e.g., Lebesgue measure on a small interval around y . In this way one can make the atom at 1 be the only atom of μ .

Regarding the degree of denseness of the extreme points in the boundary, note that at each stage if y is small enough and p_{n+1} is close enough to p_n , the new extreme point we introduce can be made arbitrarily close to w_{p_n} (here we use that α_p is continuous in $p > \vec{p}_c$, from [3]), and in the limit we can ensure an extreme point close to it. Thus, if we begin from $\mu_0 = \delta_1$ and choose p_n so that $\lim p_n = \vec{p}_c$, and using Marchand's result that the flat edge in B_{μ_n} then shrinks to a point (and symmetry of the limit shape about the axes), we can ensure ε -density of the extreme points of B_μ .

For the second part of Theorem 1.2, choose a sequence $\nu_n \in \mathcal{M}$ of continuous measures converging weakly to μ . By Corollary 2.2, $\text{sides}(B_{\nu_n}) \rightarrow \infty$, and if $\text{ext}(B_\mu)$ is ε -dense in ∂B_μ then the same holds for B_{ν_n} for sufficiently large n .

Regarding the remark after the theorem, one may verify that if at each stage y is chosen close enough to 1 and $p = \lim p_n$, then w_p is a C^∞ -point of ∂B_μ .

4. PROOF OF THEOREM 1.3.

Let us recall Hoffman's argument relating coexistence to the geometry of the limit shape for continuous μ (Theorem 1.6 of [6]). Extend τ to $\mathbb{R}^2 \times \mathbb{R}^2$ by $\tau(x, y) = \tau(x', y')$ where x' is the unique lattice point in $x + [-1/2, 1/2)^2$. Similarly, a geodesic between x, y is a geodesic between x', y' . For $S \subseteq \mathbb{R}^2$, the Busemann function $B_S : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is defined by

$$B_S(x, y) = \inf_{z \in S} \tau(x, z) - \inf_{w \in S} \tau(y, w).$$

For $v \in \mathbb{R}^2$, write $S + v = \{s + v : s \in S\}$. If $v \in \partial B_\mu$ is a point of differentiability and w is a tangent vector at v , let π_v denote the linear functional $av + bw \mapsto a$. Define the lower density of a set $A \subseteq \mathbb{N}$ by $\underline{d}(A) = \liminf \frac{1}{N} |A \cap \{1, \dots, N\}|$.

Theorem 4.1 (Hoffman [6]). *Let $\mu \in \mathcal{M}$ and let $v \in B_\mu$ be a point of differentiability of ∂B_μ with tangent line $L \subseteq \mathbb{R}^2$. Then for every $\varepsilon > 0$ there exists an $M = M(v, \varepsilon) > 0$ such that, if $x, y \in \mathbb{R}^2$ satisfy $\pi_v(x - y) > M$, then*

$$\mathbb{P}\left(\underline{d}\left(n \in \mathbb{N} : B_{L+nv}(y, x) > (1 - \varepsilon)\pi_v(x - y)\right) > 1 - \varepsilon\right) > 1 - \varepsilon.$$

Hoffman's proof of this result does not use unique passage times.

Theorem 4.1 is related to coexistence as follows. Suppose $\text{sides}(B_\mu) \geq k$. We can then find k points of differentiability $v_1, \dots, v_k \in \partial B_\mu$ with distinct tangent lines L_i , and in particular $\pi_{v_i}(v_i - v_j) > 0$ for all $j \neq i$. Fix $\varepsilon > 0$ and choose $R > 0$ large enough so that the points $x_i = Rv_i$ satisfy $\pi_{v_i}(x_i - x_j) > M(v_i, \varepsilon/k^2)$. Using the elementary relation $\underline{d}(\cap_{i=1}^n A_i) \geq 1 - \sum_{i=1}^n (1 - \underline{d}(A_i))$, for each i we have

$$\mathbb{P}\left(\underline{d}\left(n \in \mathbb{N} : B_{L+nv_i}(x_j, x_i) > 0 \text{ for all } j \neq i\right) > 1 - \frac{\varepsilon}{k}\right) > 1 - \frac{\varepsilon}{k},$$

and hence with positive probability (which can be made arbitrarily close to 1 by decreasing ε), for each i there is a positive density of n such that $B_{L_i+nv_i}(x_j, x_i) > 0$ for all $j \neq i$. For such an n , take $y_{i,n} \in L_i + nv_i$ to be the closest point (in the sense of passage times) to x_i ; assuming unique geodesics, by definition $y_{i,n}$ is reached first by species i . The points $y_{i,n}$ are in C_i , so $|C_i| = \infty$ for $i = 1, \dots, k$, i.e., coexistence occurs.

If geodesics are not unique the argument still applies, using the following observation. If $B_{L_i+nv_i}(x_j, x_i) > 0$, $j \neq i$, then the only way that $y_{i,n}$ could be colonized by a species other than i is if it is colonized by -1 . This can occur only if there is a site z on a geodesic from x_i to $y_{i,n}$ which is reached simultaneously by species i and $j \neq i$. If this happens then by concatenating the geodesic from x_j to z with the geodesic from z to $y_{i,n}$ we get a path from x_j to $L_i + nv_i$ with passage time equal to $\tau(x_i, y_{i,n})$, so $\tau(x_j, L_i + nv_i) \leq \tau(x_i, y_{i,n})$ and hence $B_{L_i+nv_i}(x_j, x_i) \leq 0$, contrary to assumption.

When $\text{sides}(B_\mu) = \infty$, one proves coexistence of infinitely many types similarly. Choose a sequence $\{v_i\}_{i=1}^\infty \subseteq \partial B_\mu$ of points of differentiability of the boundary, ordered clockwise, say. Given $\varepsilon > 0$, define the points x_i inductively by $x_{i+1} = x_i + R_i(v_{i+1} - v_i)$ for a sufficiently large $R_i > 0$ so as to ensure that for $i \neq j$, $\pi_{v_i}(x_i - x_j) > M(v_i, \varepsilon_{i,j})$, where $\sum_{i,j} \varepsilon_{i,j} < \varepsilon$. The rest of the argument is the same as above.

5. PROOF OF THEOREM 1.4.

Recall that K denotes the graph of infection. When there are unique geodesics, Hoffman's results imply that the number of ends of K is at least $\text{sides}(B_\mu)/2$ (Theorem 1.4 of [6]), establishing Theorem 1.4 for non-atomic μ . In this section we deal with the presence of atoms, proving the following result:

Theorem 5.1. *If μ is not purely atomic and has at least $s \in \mathbb{N}$ sides, then the number of ends in K is \mathbb{P} -a.s. at least*

$$(5.1) \quad k = 4 \left\lfloor \frac{s-4}{12} \right\rfloor.$$

See below Theorem 5.3 for an explanation of this bound.

For simplicity, in the following discussion we fix a measure $\mu \in \mathcal{M}_p$ with $p < 1$, and make two further assumptions: (a) the only atom of μ is the atom at 1, and (b) μ is supported on a bounded interval $[1, R]$. The argument can be modified to deal with the general case of the theorem.

Proposition 5.2. *Suppose that \mathbb{P} -a.s. there exist k infinite geodesics $\gamma_1, \dots, \gamma_k$ starting at 0, edges e_1, \dots, e_k , and a finite set $V \subseteq \mathbb{Z}^2$, such that (a) e_i lies on γ_i but not on γ_j for $i \neq j$, (b) the endpoints of e_i are in V , (c) $\tau_{e_i} > 1$, and (d) each pair of geodesics is disjoint outside of V . Then \mathbb{P} -a.s., K has k ends.*

Proof. Under our assumptions on μ , with probability 1 every pair of edges with passage times > 1 has distinct passage times. Consequently, geodesics between $x, y \in \mathbb{Z}^2$ can

differ only in edges e with $\tau_e = 1$, and must share edges with $\tau_e > 1$. We assume we are in this probability-1 event.

We claim that no two of the given geodesics are connected in $K \setminus V$. Suppose for instance that γ_1, γ_2 were connected in $K \setminus V$ by a path σ which we may assume is simple (non self-intersecting) and with endpoints $y_1 \in \gamma_1$ and $y_2 \in \gamma_2$. Denote the sequence of vertices in σ by $y_1 = v_1, v_2, \dots, v_k = y_2$. Write $e = e_1$ and let $J \subseteq \{1, \dots, k\}$ denote the set of j such that there exists a geodesic σ_j from 0 to v_j which contains e . We claim that $k \in J$. This leads to a contradiction because then σ_k and γ_2 are both geodesics connecting 0 and y_2 , but only one of them, σ_k , contains e .

Clearly $1 \in J$. Suppose now that $j \in J$ with corresponding geodesic σ_j . Write f for the edge between v_j and v_{j+1} , and note that $f \neq e$ because the endpoints of e are in V while those of f are not. Also, $\tau(0, v_j) \neq \tau(0, v_{j+1})$, since $f \in K$. If $\tau(0, v_{j+1}) > \tau(0, v_j)$ then we adjoin f to σ_j and obtain a geodesic σ_{j+1} with the desired properties. If $\tau(0, v_{j+1}) < \tau(0, v_j)$ and v_{j+1} lies on σ_j we remove f from σ_j to obtain σ_{j+1} . On the other hand, if v_{j+1} does not lie on σ_j but $\tau(0, v_{j+1}) < \tau(0, v_j)$, then there is a geodesic σ'_{j+1} from 0 to v_j whose last edge is f . Because σ'_{j+1} must reach v_j in the same time as σ_j does, it must pass through each of those edges of σ_j which have passage times > 1 , and in particular through e . We remove f from σ'_{j+1} to obtain σ_{j+1} . \square

Our goal is to establish the hypotheses of the proposition for k as in (5.1). It is enough to show that there exist random variables $m < M$ such that with probability one,

- (1) There exist k geodesics $\gamma_1, \dots, \gamma_k$ which are disjoint outside of mB_μ .
- (2) There are edges e_i in γ_i , with endpoints in $MB_\mu \setminus mB_\mu$, such that $\tau_{e_i} > 1$.

This suffices because we can then set $V = MB_\mu$ in the proposition. To show that such m, M exist, it is enough to show that for every $\varepsilon > 0$ there exists deterministic $m < M$ such that each of the conditions above holds on an event of probability $> 1 - \varepsilon$.

Given $u, v, w \in \partial B_\mu$ which are points of differentiability of ∂B_μ , let $C(u, v, w)$ denote the open arc in ∂B_μ from u to w containing v . We rely on the following result, whose proof does not require unique geodesics:

Theorem 5.3 (Hoffman [6]). *Let $u, v, w \in \partial B_\mu$ be points of differentiability of ∂B_μ , let L be the tangent line at v , and write $C = C(u, v, w)$. Then for every $\varepsilon > 0$ there is an $M_0 = M_0(\varepsilon)$ such that for every $M > M_0$, with probability $> 1 - \varepsilon$ the set*

$$I = \{n \in \mathbb{N} : \gamma \cap M\partial B_\mu \subseteq MC \text{ for all geodesics } \gamma \text{ from } 0 \text{ to } L + nv\}$$

satisfies $\underline{d}(I) > 1 - \varepsilon$.

Henceforth fix k as in (5.1) and $\varepsilon > 0$ and for $i = 1, \dots, k$ choose points $u_i, v_i, w_i \in \partial B_\mu$ and lines L_i as in the theorem, and such that the closed sets $C_i = \overline{C(u_i, v_i, w_i)}$ are pairwise disjoint and do not intersect the boundary of the ℓ^1 unit ball; write $C = \bigcup_{i=1}^k C_i$. Note that k was picked so that such a choice is possible: note that $\frac{1}{4}(\text{sides}(B_\mu) - 4)$ is the number of distinct sides on each of the four curves in ∂B_μ which constitute the

complement of the ℓ^1 unit ball; dividing this number by 3 gives an upper bound on the number of triples we can choose in each of these curves. Taking integer part and multiplying by 4 gives k .

Claim 5.4. *There exists M_0 and $\rho > 0$ such that with probability at least $1 - \varepsilon$, for all $M > M_0$, every $x \in MC$ and every geodesic γ from 0 to x , at least a ρ -fraction of the edges of γ have passage times > 1 .*

Proof. Pick $\delta > 0$ and define edge weights $\{\tau'_e : e \in \mathbb{E}\}$ by the rule that if $\tau_e > 1$ then $\tau'_e = \tau_e + \delta$ and $\tau'_e = \tau_e$ otherwise. Let μ' denote the marginal distribution of τ'_e .

Choose $\eta > 0$ so that $(1 - \eta)C \cap B_{\mu'} = \emptyset$ (we can do so by a theorem of Marchand [7, Theorem 1.5] and the fact that C is disjoint from the ℓ^1 unit ball). For a path σ let $f(\sigma)$ denote the fraction of edges of σ with passage time > 1 . By Theorem 1.1, there is an event A with $\mathbb{P}(A) > 1 - \varepsilon$ and an M_0 such that for all $M > M_0$ and $y \in \partial B_\mu$, the τ -geodesic γ from 0 to y satisfies $(1 - \eta^2)M < \tau(\gamma) < (1 + \eta^2)M$, and similarly for $y' \in M\partial B'_\mu$ and the τ' -length of τ' -geodesics from 0 to y' . We claim that A is the desired event. Indeed, let $M > M_0$ and let γ be a τ -geodesic from 0 to some $x \in MC$. Since $\tau_e \geq 1$ for all $e \in \mathbb{E}$, the number of edges in γ is at most $\tau(\gamma)$. Hence

$$\begin{aligned} \tau'(\gamma) &= \tau(\gamma) + \delta f(\gamma) \#\{\text{edges of } \gamma\} \\ &\leq (1 + \delta f(\gamma))\tau(\gamma) \\ &\leq (1 + \delta f(\gamma))(1 + \eta^2)M. \end{aligned}$$

On the other hand $x = \frac{M}{s}y$ for some $y \in \partial B_{\mu'}$ and $s < 1 - \eta$, so

$$\tau'(\gamma) \geq (1 - \eta^2) \frac{M}{1 - \eta}.$$

Combining these we find that $f(\gamma) \geq \frac{\eta}{\delta} \cdot \frac{1 - \eta}{1 + \eta^2}$. This lower bound on f can serve as ρ . \square

Let $\alpha > 0$ be the quantity

$$(5.2) \quad \alpha = \frac{1}{2} \min\{\pi_{v_i}(x_i - x_j) : x_i \in C_i, x_j \in C_j, i \neq j\}.$$

and fix finite, $\frac{\alpha}{100R}$ -dense sets $D_i \subseteq C_i$. Let ρ be as in the claim and $L \gg m/\rho$. By Theorems 5.3 and 1.1, we can choose an integer m and $M = Lm$ such that, with probability $> 1 - \varepsilon$, there is a set $I \subseteq \mathbb{N}$ of density $> 1 - \varepsilon$ such that, for $n \in I$,

- (A) Every geodesic $\gamma_{i,n}$ from 0 to $L_i + nv_i$ intersects $m\partial B_\mu$ in mC_i and intersects $M\partial B_\mu$ in MC_i .
- (B) If $i \neq j$ then $B_{L_i + nv_i}(x_j, x_i) > m\alpha$ for all $x_i \in mD_i$ and $x_j \in mD_j$.
- (C) $|\tau(0, x) - m| < \frac{m\alpha}{10}$ for all $x \in mD_i$.
- (D) At least a $\rho/2$ -fraction of edges on $\gamma_{i,n} \cap (MB_\mu \setminus mB_\mu)$ have passage time > 1 .

Fix $\gamma_{i,n}$ as in (A). We may choose an infinite $J \subseteq I$ such that $\lim_{n \in J} \gamma_{i,n} \rightarrow \gamma_i$ for some infinite geodesics γ_i originating at 0, i.e., for every $r > 0$ we have $\gamma_i \cap [-r, r]^2 = \gamma_{i,n} \cap [-r, r]^2$ for all large enough $n \in J$. Henceforth we only consider such n .

Let $y_{i,n}$ be the first intersection point of $\gamma_{i,n}$ with mC_i , and choose $x_{i,n} \in D_i$ such that $|x_{i,n} - y_{i,n}| \leq \frac{m\alpha}{R10}$. Since μ is supported on $[0, R]$, we conclude that $\tau(x_{i,n}, y_{i,n}) \leq \frac{m\alpha}{10}$, so

$$(5.3) \quad |B_{L_i+nv_i}(y_{j,n}, y_{i,n}) - B_{L_i+nv_i}(x_{j,n}, x_{i,n})| < \frac{2m\alpha}{10} \text{ for } i \neq j.$$

Claim 5.5. *The γ_i 's are disjoint outside of mB_μ .*

Proof. Suppose for example that γ_1, γ_2 intersect at some point z outside of mB_μ . Then for large enough $n \in J$ the same is true of $\gamma_{1,n}$ and $\gamma_{2,n}$. Then

$$\tau(0, y_{1,n}) + \tau(y_{1,n}, z) = \tau(0, y_{2,n}) + \tau(y_{2,n}, z).$$

By (C) we have $|\tau(0, y_{1,n}) - \tau(0, y_{2,n})| < \frac{2m\alpha}{10}$, so

$$|\tau(y_{1,n}, z) - \tau(y_{2,n}, z)| < \frac{2m\alpha}{10}.$$

Write σ_1 for the part of $\gamma_{1,n}$ from $y_{1,n}$ to $L_1 + nv_1$. Let σ_2 be path which starts at $y_{2,n}$, follows $\gamma_{2,n}$ until z , and then follows $\gamma_{1,n}$ until $L_1 + nv_1$. We find that $|\tau(\sigma_1) - \tau(\sigma_2)| < \frac{2m\alpha}{10}$. But $\gamma_{1,n}$ is a shortest path from 0 to $L_1 + nv_1$, so σ_1 is a shortest path from $y_{1,n}$ to $L_1 + nv_1$. Hence $B_{L_1+nv_1}(y_{2,n}, y_{1,n}) \leq \frac{2m\alpha}{10}$. By (5.3), this contradicts (B). \square

Finally, combining the last claim with (D) establishes the two claims stated after Proposition 5.2. This completes the proof of Theorem 1.4.

REFERENCES

- [1] Cox, J. T. and Durrett, R. (1981). Some limit theorems for percolation with necessary and sufficient conditions. *Ann. Probab.* **9** 583–603.
- [2] Cox, J. T. and Kesten, H. (1981). On the continuity of the time constant of first-passage percolation. *J. Appl. Probab.* **18** 809–819.
- [3] Durrett, R. (1984). Oriented percolation in two dimensions. *Ann. Probab.* **12** 999–1040.
- [4] Durrett, R. and Liggett, T. (1981). The shape of the limit set in Richardson's growth model. *Ann. Probab.* **9** 186–193.
- [5] Häggström, O. and Pemantle, R. (1998). First passage percolation and a model for competing spatial growth. *J. Appl. Probab.* **35** 683–692.
- [6] Hoffman, C. (2008). Geodesics in first passage percolation. *Ann. Appl. Probab.* **18** 1944–1969.
- [7] Marchand, R. (2002). Strict inequalities for the time constant in first passage percolation. *Ann. Appl. Probab.* **12** 1001–1038.
- [8] Newman, C. A surface view of first-passage percolation. Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 1017–1023, Birkhauser, Basel, 1995.
- [9] Häggström, O. and Meester, R. (1995). Asymptotic shapes for stationary first passage percolation. *Ann. Probab.* **23** 1511–1522.
- [10] Zhang, Yu, (2007). Shape curvatures and transversal fluctuations in the first passage percolation model. *Preprint*, <http://arxiv.org/abs/math.PR/0701689>

MATHEMATICS DEPARTMENT, PRINCETON UNIVERSITY, FINE HALL, WASHINGTON RD., PRINCETON, NJ 08544.

E-mail address: Michael Damron: mdamron@math.princeton.edu,

Michael Hochman: hochman@math.princeton.edu