

Approachable Free Subsets and Fine Structure Derived Scales*

Dominik Adolf and Omer Ben-Neria

February 16, 2023

Abstract

Shelah showed that the existence of free subsets over internally approachable subalgebras follows from the failure of the PCF conjecture on intervals of regular cardinals. We show that a stronger property called the Approachable Bounded Subset Property can be forced from the assumption of a cardinal λ for which the set of Mitchell orders $\{o(\mu) \mid \mu < \lambda\}$ is unbounded in λ . Furthermore, we study the related notion of continuous tree-like scales, and show that such scales must exist on all products in canonical inner models. We use this result, together with a covering-type argument, to show that the large cardinal hypothesis from the forcing part is optimal.

1 Introduction

The study of set theoretic algebras has been central in many areas, with many applications to compactness principles, cardinal arithmetic, and combinatorial set theory.

An algebra on a set X is a tuple $\mathfrak{A} = \langle X, f_n \rangle_{n < \omega}$ where $f_n : X^{k_n} \rightarrow X$ is a function. A sub-algebra is a subset $M \subseteq X$ such that $f_n(x_0, \dots, x_{k_n-1}) \in M$ for all $(x_0, \dots, x_{k_n-1}) \in M^{k_n}$ and $n < \omega$. The set of sub-algebras of \mathfrak{A} is known as a club (in $\mathcal{P}(X)$). The characteristic function χ_M of M is defined on the ordinals of M by $\chi_M(\tau) = \sup(M \cap \tau)$.

Shelah's celebrated bound in cardinal arithmetic ([25]) states that if \aleph_ω is a strong limit cardinal then

$$2^{\aleph_\omega} < \min\{\aleph_{\omega_4}, \aleph_{(2^{\aleph_0})^+}\}.$$

Starting from a supercompact cardinal, Shelah proved that for every $\alpha < \omega_1$, there exists a generic extension in which $2^{\aleph_\omega} = \aleph_{\alpha+1}$ (see [14]). It is a central open problem in cardinal arithmetic if $2^{\aleph_\omega} \geq \aleph_{\omega_1}$ is consistent. A major work towards a possible solution is the work of Gitik ([13]) on the failure of the

*2010 *Mathematics Subject Classification*. Primary 03E04, 03E45, 03E55

PCF-conjecture. Shelah’s PCF conjecture states that $|\text{pcf}(A)| \leq |A|$ for every progressive¹ set A of regular cardinals. In [26], Shelah has extracted remarkable freeness properties of sets over subalgebras, from the assumption of $2^{\aleph_\omega} \geq \aleph_{\omega_1}$, or more generally, from the assumption $|\text{pcf}(A)| > |A|$ for a progressive interval of regular cardinals $|A|$.

Definition 1. Let $\mathfrak{A} = \langle X, f_n \rangle_n$ be an algebra and $x \subseteq X$. We say that x is **free** with respect to \mathfrak{A} if for every $\delta \in x$ and $n < \omega$, $\delta \notin f_n \text{``}(x \setminus \{\delta\})^{<\omega}$. More generally, x is free over a subalgebra $N \subseteq \mathfrak{A}$ if for every $\delta \in x$ and $n < \omega$, $\delta \notin f_n \text{``}(N \cup (x \setminus \{\delta\}))^{<\omega}$.

A cardinal λ has the Free Subset Property if every algebra \mathfrak{A} on λ or a bigger H_θ , has a free subset $x \subseteq \lambda$ which is cofinal in λ . A regular cardinal λ with the Free Subset Property is Jonsson. Koepke [18] has shown that the free subset property at \aleph_ω is equiconsistent with the existence of a measurable cardinal. For a singular limit λ of a progressive interval $|A|$, it is shown in [26] that if $|\text{pcf}(A)| > |A|$ then λ satisfies the Free Subset Property. In his PhD thesis ([22]), Pereira has isolated the notion of the Approachable Free Subset Property (AFSP) to play a critical role in the result from [26]. The Approachable Free Subset Property for a singular cardinal λ asserts that there exists some sufficiently large H_θ , $\theta > \lambda$ and an algebra \mathfrak{A} on H_θ such that for every internally approachable substructure² $N \prec \mathfrak{A}$ with $|N| < \lambda$, there exists an infinite sequence of regular cardinal $\langle \tau_i \mid i < \text{cof}(\lambda) \rangle \in N$ such that the set $x = \{\chi_N(\tau_i) \mid i < \text{cof}(\lambda)\}$ is free over N .

Pereira showed that Shelah’s proof yields that if λ is a limit of a progressive interval A of regular cardinals and $|\text{pcf}(A)| > |A|$ then the Approachable Free Subset Property holds at λ .

Working with fixed sequences $\langle \tau_n \mid n < \omega \rangle$ of regular cardinal, we consider here the following version of this property.

Definition 2. The **Approachable Free Subset Property (AFSP)** with respect to $\langle \tau_n \rangle_n$ asserts that for every sufficiently large regular $\theta > \lambda = (\cup_n \tau_n)$ and for every internally approachable subalgebra $N \prec \mathfrak{A}$, of an algebra \mathfrak{A} extending $(H_\theta, \in, \langle \tau_n \rangle_n)$, satisfying $|N| < \lambda$ there exists a cofinite set $x \subseteq \{\chi_N(\tau_n) \mid n < \omega\}$ which is free over N .

By moving from one cardinal θ to $\theta' > \theta$ if needed, it is routine to verify the definition of AFSP with respect to a sequence $\langle \tau_n \rangle_n$ can be replaced with a similar assertion in which the requirement of “every internally approachable N ” is replaced with “for every internally approachable in some closed unbounded subset of $\mathcal{P}_\lambda(\mathfrak{A})$ ”. Clearly, if AFSP holds with respect to a sequence $\langle \tau_n \rangle_n$ then AFSP holds with respect to the singular limit $\lambda = \cup_n \tau_n$, as in the original definition of [22].

¹I.e., $\min(A) > |A|$.

²See Definition 10

The above mentioned results, suggest that AFSP can provide a path to possibly improving Shelah's bound, to $2^{\aleph_\omega} < \aleph_{\omega_1}$. I.e., proving (in ZFC) that AFSP must fail at \aleph_ω (or AFSP fails w.r.t every subsequence $\langle \tau_n \rangle_n$ of $\{\aleph_k \mid k < \omega\}$) would imply that $2^{\aleph_\omega} < \aleph_{\omega_1}$. To this end, Pereira ([22]) has isolated the notion of **tree-like scales**, as a potential tool of proving AFSP must fail.

Definition 3. Let $\langle \tau_n \rangle_{n < \omega}$ be an increasing sequence of regular cardinals. A scale³ $\vec{f} = \langle f_\alpha \mid \alpha < \eta \rangle$ is a tree-like scale on $\prod_{n < \omega} \tau_n$ if for every $\alpha \neq \beta < \eta$ and $n < \omega$, $f_\alpha(n+1) = f_\beta(n+1)$ implies $f_\alpha(n) = f_\beta(n)$.

Pereira shows in [23] that the existence of a continuous tree-like scale on a product $\prod_{n < \omega} \tau_n$ guarantees the failure of AFSP with respect to $\langle \tau_n \rangle_n$ (see also Lemma 15), and further proves that continuous tree-like scales, unlike other well-known types of scales, such as good scales, can exist in models with some of the strongest large cardinal notions, e.g. I_0 -cardinals. Moreover, Cummings [5] proved that tree-like scales can exist above supercompact cardinals. These results show that as opposed to other well-known properties of scales such as good and very-good scales, which exhibit desirable "local" behaviour but cannot exist in the presence of certain large cardinals ([6]), the notion of continuous tree-like scales may coexist with the some of the strongest large cardinals hypothesis.

The consistency of the inexistence of a continuous tree-like scale on a product $\prod_n \tau_n$ of regular cardinal has been established by Gitik in [11], from the consistency assumption of a cardinal κ satisfying $o(\kappa) = \kappa^{++} + 1$. The argument makes a sophisticated use of the key features of Gitik's extender based Prikry forcing by a (κ, κ^{++}) -extender.⁴ Concerning the possible consistency of the Approachable Free Subset Property, Welch ([30]) has shown that its strengthening ABSP (Definition 7 below) with respect to a sequence $\langle \tau_n \rangle_n$ implies that the large cardinal assumption of Theorem 4 holds in an inner model.

It remained open whether AFSP with respect to some sequence $\langle \tau_n \mid n < \omega \rangle$ is consistent at all, and if so, whether its consistency strength is strictly stronger than the (seemingly) weaker property, of no continuous tree-like scale on $\prod_n \tau_n$. The current work answers both questions:

Theorem 4. *It is consistent relative to the existence of a cardinal λ such that the set of Mitchell orders $\{o(\mu) \mid \mu < \lambda\}$ is unbounded in λ , that the Approachable Free Subset Property holds with respect to some sequence of regular cardinals $\vec{\tau} = \langle \tau_n \rangle_n$.*

Moreover, the sequence τ_n can be made to be a subsequence of the first uncountable cardinals, in a model where $\lambda = \aleph_\omega$.

Theorem 5. *Let λ be a singular cardinal of countable cofinality such that there is no inner model M with $\lambda = \sup\{o^M(\mu) \mid \mu < \lambda\}$. Let $\langle \tau_n \mid n < \omega \rangle$ be a*

³see Definition 9 for the definition of a continuous scale

⁴E.g., on the fact that there are unboundedly many pairs $(\alpha, \alpha^*) \in [\kappa]^2$, sharing the same Rudin-Keisler projection map $\pi_{\alpha^*, \alpha}$.

sequence of regular cardinals cofinal in λ . Then $\prod_{n < \omega} \tau_n$ carries a continuous tree-like scale.

To achieve the proof of Theorem 5, we establish a result of an independent interest, that the continuous tree-like scales naturally appear in fine-structural canonical inner models. Thus obtaining complementary result to aforementioned theorems by Pereira and Cummings, i.e. we know that no large cardinal property that can consistently appear in canonical inner models disproves the existence of products with continuous tree-like scales (e.g., Woodin cardinals).

Theorem 6. *Let \mathcal{M} be a premouse such that each countable hull has an ω -maximal $(\omega_1 + 1)$ -iteration strategy. Let $\lambda \in \mathcal{M}$ be a singular cardinal of countable cofinality. Let $\langle \kappa_i : i < \omega \rangle$ be a sequence of regular cardinals cofinal in λ . Then $\prod_{i < \omega} \kappa_i / J_{bd}$ carries a continuous tree-like scale.*

Continuous tree-like scales on products of successor cardinals in L where implicitly constructed by Donder, Jensen, and Stanley in [7]. In the course of proving Theorem 4, we establish the consistency of a principle stronger than AFSP, which we call the Approachable Bounded Subset Property.

Let N be a subalgebra of $\mathfrak{A} = \langle H_\theta, f_n \rangle_n$ and $\vec{\tau} = \langle \tau_n \rangle_n$ be an increasing sequence of cardinals. Given a set $x \subseteq H_\theta$, we define $N[x]$ to be the \mathfrak{A} -closure of the set $(x \cup N)$. We say that N satisfies the **Bounded Appending Property** with respect to $\vec{\tau}$ if for every $n_0 < \omega$, setting $x = \{\chi_N(\tau_n) \mid n \neq n_0\}$ then the addition of x to N does not increase the supremum below τ_{n_0} , namely $\chi_{N[x]}(\tau_{n_0}) = \chi_N(\tau_{n_0})$.

Definition 7. The **Approachable Bounded Subset Property** holds with respect to $\langle \tau_n \rangle_n$ if for every sufficiently large regular $\theta > \lambda = (\cup_n \tau_n)$ and internally approachable subalgebra $N \prec \mathfrak{A}$ of an algebra \mathfrak{A} extending $(H_\theta, \in, \langle \tau_n \rangle_n)$ of size $|N| < \lambda$, N satisfies the bounded appending property with respect to a tail of $\langle \tau_n \rangle_n$.

We show in Lemma 15 ABSP with respect to a sequence $\langle \tau_n \rangle_n$ implies AFSP with respect to the same sequence, as well as the inexistence of a continuous **essentially tree-like** scale; a weakening of tree-like scale introduced by Pereira (see Definition 9). The proof of the forcing Theorem 4, stated above, goes through proving that ABSP is consistent with respect to a sequence of regulars $\langle \tau_n \rangle_n$.

The following summarizes the main results of this paper:

Corollary 8. *The following principles are equiconsistent:*

1. *There exists a sequence of regular cardinals $\langle \tau_n \mid n < \omega \rangle$ for which the Approachable Bounded Subset Property holds.*
2. *There exists a sequence of regular cardinals $\langle \tau_n \mid n < \omega \rangle$ for which the Approachable Free Subset Property holds.*

3. *There exists a sequence of regular cardinals $\langle \tau_n \mid n < \omega \rangle$ for which the product $\prod_n \tau_n$ does not carry a continuous Tree-Like scale.*
4. *There exists a cardinal λ such that the set of Mitchell orders $\{o(\mu) \mid \mu < \lambda\}$ is unbounded in λ .*

The paper is organized as follows: The remainder of **this section** will be dedicated to discussing preliminary material in PCF theory and the theory of inner models. **Section 2** will be dedicated to the forcing argument establishing the proof Theorem 4. In **Section 3** we discuss how to construct tree-like scales from the fine structure of canonical inner models. In **Section 4** we will use these fine structural scales to derive scales on products in V using a covering-like argument. Finally, in **Section 5** we finish with a list of open problems.

Acknowledgments:

The work on this project was initiated following a suggestion by Assaf Rinot to study the consistency of the Approachable Free Subset Property. The authors are grateful for this suggestion and for supporting the first author during the academic year of 2018-2019 at Bar-Ilan University under a grant from the European Research Council (grant agreement ERC-2018-StG 802756). The initial idea for the inner model construction of continuous tree-like scale was conceived during the Berkeley conference on Inner Model theory in July 2019. The authors would like to thank Ralf Schindler and John Steel for organizing the meeting and creating the opportunity for this collaboration. The first author would like to thank Grigor Sargsyan for his generous support and warm hospitality during the Spring of 2020 (NSF career award DMS-1352034). During that time the first author had the opportunity to travel to Pittsburgh. It was there that some significant improvements were made to the lower bound argument, and would like to thank James Cummings for the opportunity to present this research and insightful conversations. The second author was partially supported by the Israel Science Foundation (Grant 1832/19). He would like thank Luis Pereira for insightful discussions on the subject and many valuable remarks on this paper, and to Spencer Unger and Philip Welch for valuable comments and suggestions. Finally, the authors would like to thank the referee for a thorough report and valuable corrections and comments.

1.1 Preliminaries

For a set X and a cardinal λ , $\mathcal{P}_\lambda(X)$ denotes the collections of all subsets $a \subseteq X$ of size $|a| < \lambda$. J_{bd} denotes the ideal of bounded subsets of ω . Let I be an ideal on ω and f, g two functions from ω to ordinals. We write $f <_I g$ if $\{n < \omega \mid f(n) \geq g(n)\} \in I$. We write $f <^* g$ for $f <_{J_{bd}} g$.

1.1.1 Continuous and Tree-Like Scales

Let $\langle \tau_n \mid n < \omega \rangle$ be a sequence of ordinals of strictly increasing cofinalities. A sequence of functions $\vec{f} = \langle f_\alpha \mid \alpha < \eta \rangle \subseteq \prod_n \tau_n$ of a regular length η , is a **pre-**

scale in $(\prod_n \tau_n, <_I)$ if \vec{f} is strictly increasing in the ordering $<_I$. A prescale is a **scale** if it is cofinal in $\prod_n \tau_n$. As we focus on J_{bd} from this point forward, we will frequently say that \vec{f} is a (pre-)scale in $\prod_n \tau_n$, without mentioning the ideal J_{bd} .

Definition 9. Suppose that $\vec{f} = \langle f_\alpha \mid \alpha < \eta \rangle$ is a (pre-)scale in $\prod_n \tau_n$.

1. \vec{f} is said to be **continuous** if for every limit ordinal $\delta < \eta$ of uncountable cofinality, whenever there is a function $g \in \prod_n \tau_n$ so that the sequence $\vec{f} \upharpoonright \delta$ is $<^*$ -cofinal in $\prod_n g(n)$ then $f_\delta(n) = g(n)$ for all but finitely many $n < \omega$.
2. \vec{f} is **Tree-like** if for every $\alpha \neq \beta < \eta$ and $n < \omega$, if $f_\alpha(n+1) = f_\beta(n+1)$ then $f_\alpha(n) = f_\beta(n)$.
3. \vec{f} is **Essentially Tree-like** if for every $n < \omega$ and $\mu \in [\tau_n, \tau_{n+1})$ the set

$$\{\mu' < \tau_n \mid \exists \beta < \eta, f_\beta(n+1) = \mu \text{ and } f_\beta(n) = \mu'\}$$

is nonstationary in τ_n .

If a product $\prod_n \tau_n$ carries a scale, it is not difficult to find another scale on it with the tree-like property (see Pereira [22]), but such a scale need not be continuous. We note that the notion of a continuous scale is weaker than the one of a good scale $\vec{f} = \langle f_\alpha \mid \alpha < \eta \rangle$, in which for every uncountable cofinality ordinal $\delta < \eta$, f_δ is required to be an exact upper bound of $\vec{f} \upharpoonright \delta$ and satisfy that $\text{cof}(f_\delta(n)) = \text{cof}(\delta)$ for almost all $n < \omega$. Cummings ([5]) has shown that the existence of a continuous tree-like scale which is not good is consistent.

1.1.2 Internally Approachable Structures and related Principles

Considering notions such as the Approachable Free Subset Property or the Approachable Bounded Subset Property with respect to subalgebras of Algebras $\mathfrak{A} = (\theta, f_n)_n$, there is no harm in replacing the domain θ with another set of the same size, such as H_θ in cases relevant to us, and adding more structure to the algebra. Therefore, from this point on, we will only restrict ourselves to set theoretic algebras \mathfrak{A} of the form $\mathfrak{A} = (H_\theta, \in, f_n)_n$, which extend the model (H_θ, \in) in the language of set theory, and include Skolem functions. In particular, a subalgebra $N \prec \mathfrak{A}$ will always be an elementary substructure.

This allows us to reformulate our notion of freeness. Assuming⁵ the algebra \mathfrak{A} is rich enough to satisfy a fraction of ZFC⁶, and $N \subseteq \mathfrak{A}$ is sufficiently closed so it is an elementary substructure $N \prec \mathfrak{A}$, then the fact that a set x is free over N is equivalent to having that for every $\delta \in x$ and a function $f \in N$, $\delta \notin f^{<\omega}(x \setminus \{\delta\})^{<\omega}$.

⁵we will always be able to assume so

⁶specifically, the Replacement property

The notion of internally approachable structures was formally introduced in [9]. We refer the reader to [8] for further exposition. The definition below is similar to the standard ones, with the addition that here, we will focus on internally approachable unions of uncountable cofinality.

Definition 10. An elementary subalgebra (substructure) $N \prec \mathfrak{A}$ of an algebra $\mathfrak{A} = (H_\theta; \in, f_n)_n$ is said to be internally approachable of length ρ if $N = \bigcup_{i < \rho} N_i$ is a union of a sequence $\vec{N} = \langle N_i \mid i < \rho \rangle$ of elementary subalgebras $N_i \prec N$, and for every $j < \rho$, $\vec{N} \upharpoonright j = \langle N_i \mid i < j \rangle$ belongs to N .

We say that $N \prec \mathfrak{A}$ is **internally approachable** if it is internally approachable of length ρ for some ρ of uncountable cofinality $\text{cof}(\rho) > \aleph_0$.

Notation 11. Let $N \prec (H_\theta; \in)$ for a regular cardinal θ .

- For every regular cardinal $\tau \in N$, define $\chi_N(\tau) = \sup(N \cap \tau)$.
- Given a sequence $\vec{\tau} = \langle \tau_n \mid n < \omega \rangle \subseteq N$ define the function $\chi_N^{\vec{\tau}} \in \prod_n \tau_n$ by $\chi_N^{\vec{\tau}}(n) = \chi_N(\tau_n)$ if the last ordinal is strictly smaller than τ_n , and 0 otherwise.

The following folklore result connects continuous scales with characteristic functions of internally approachable structures. We include a proof for completeness.

Lemma 12. *Suppose that $\vec{\tau} = \langle \tau_n \mid n < \omega \rangle \in N$ is a strictly increasing sequence of regular cardinals for which $\prod_n \tau_n$ carries a continuous scale $\vec{f} = \langle f_\alpha \mid \alpha < \eta \rangle$. For every $N \prec H_\theta$ which is internally approachable of size $|N| < \bigcup_n \tau_n$, with $\vec{f} \in N$, if $\delta = \chi_N(\eta)$ then $\chi_N^{\vec{\tau}}(n) = f_\delta(n)$ for all but finitely many $n < \omega$.*

Proof. Let $\vec{N} = \langle N_i \mid i < \rho \rangle$ be a sequence witnessing $N = \bigcup_i N_i$ is internally approachable of length ρ which has uncountable cofinality. Since \vec{f} is continuous, it suffices to show that $\vec{f} \upharpoonright \delta$ is $<^*$ -cofinally interleaved with the functions in $\prod_n \chi_N^{\vec{\tau}}(n)$ to prove that $\chi_N^{\vec{\tau}}(n) = f_\delta(n)$ for almost all $n < \omega$. First, for every $f_\alpha \in \vec{f} \upharpoonright \delta$ there exists some $\beta \in N \cap \delta$ so that $\alpha < \beta$, and thus $f_\alpha <^* f_\beta$. But $f_\beta \in N$ since $\beta \in N$, which means that $f_\beta \in \prod_n \chi_N^{\vec{\tau}}(n)$. Next, fix $g \in \prod_n \chi_N(\tau_n)$. We show that $g <^* f_\alpha$ for some $\alpha < \delta$. To this end, $N = \bigcup_{i < \rho} N_i$ guarantees that for each $n < \omega$ there is $i < \rho$ such that $g(n) < \chi_{N_i}(\tau_n)$. Since $\text{cof}(\rho) > \aleph_0$ there is $i < \rho$ such that $g(n) < \chi_{N_i}(\tau_n)$ for all n , and in particular, $g <^* \chi_{N_i}^{\vec{\tau}}$. Since $\vec{f} \in N$ is $<^*$ -cofinal in $\prod_n \tau_n$ and $\chi_{N_i}^{\vec{\tau}} \in N$, there is some $\alpha \in N \cap \eta \subseteq \delta$ so that $\chi_{N_i}^{\vec{\tau}} <^* f_\alpha$, and thus $g <^* f_\alpha$. \square

Lemma 13. *Let $\lambda \leq \eta < \theta$ be cardinals with η, θ regular and $\rho^{\aleph_0} < \eta$ for every $\rho < \lambda$. Suppose that \triangleleft is a well-ordering of H_θ , $\mathcal{S} \subseteq \mathcal{P}_\lambda(H_\theta)$ is a stationary set of internally approachable structures $N \prec (H_\theta; \in, \{\eta\}, \triangleleft)$, and $X \in \bigcap_{N \in \mathcal{S}} N$ is a set satisfying $|X^\omega| \leq \eta$. For every assignment that maps $N \in \mathcal{S}$ to $\langle x_n^N \mid n < \omega \rangle \in (X \cap N)^\omega$ there exists a stationary subset $S^* \subseteq \eta$ and a constant sequence $\langle x_n \mid n < \omega \rangle$ such that for every $\delta \in S^*$ there is $N \in \mathcal{S}$ with $\chi_N(\eta) = \delta$ and $\langle x_n^N \rangle_n = \langle x_n \rangle_n$.*

Proof. Let $\langle \bar{x}^\alpha \mid \alpha < \eta \rangle$ be the \triangleleft -least enumeration of X^ω in H_θ , where each \bar{x}^α is of the form $\langle x_n^\alpha \mid n < \omega \rangle$. For each $N \in \mathcal{S}$ let $\alpha_N < \eta$ be such that $\langle x_n^{\alpha_N} \mid n < \omega \rangle = \bar{x}^{\alpha_N}$. Note that α_N may not be a member of N as $\langle x_n^{\alpha_N} \mid n < \omega \rangle$ need not be in N . Since each $N \in \mathcal{S}$ is the union of a sequence $\langle N_i \mid i < \rho \rangle$ with $\text{cof}(\rho) > \aleph_0$, and $\langle x_n^{\alpha_N} \mid n < \omega \rangle \subseteq N$, there is some $i < \rho$ so that $\langle x_n^{\alpha_N} \mid n < \omega \rangle \subset N_i$, and thus $\langle x_n^{\alpha_N} \mid n < \omega \rangle \in (X \cap N_i)^\omega \in N$. Moreover, as $|X \cap N_i| < \lambda$, we have that $|(X \cap N_i)^\omega| < \eta$, and therefore there exists some $\beta_N \in \eta \cap N$ so that $(X \cap N_i)^\omega \subset \langle \bar{x}^\alpha \mid \alpha < \beta_N \rangle$. We conclude that, $\alpha_N < \beta_N < \chi_N(\eta)$. Next, define $S = \{\chi_N(\eta) \mid N \in \mathcal{S}\}$. $S \subseteq \eta$ is stationary, and by choosing for each $\delta \in S$ a specific structure $N_\delta \in \mathcal{S}$ with $\delta = \chi_{N_\delta}(\eta)$, we can form a pressing down assignment taking each $\delta \in S$ to $\alpha_{N_\delta} < \delta$. Let $\alpha^* < \eta$ and $S^* \subseteq S$ be so that $\alpha_{N_\delta} = \alpha^*$ for all $\delta \in S^*$. The claim follows for S^* and $\langle x_n \mid n < \omega \rangle = \bar{x}^{\alpha^*}$. \square

Let $\vec{\tau} = \langle \tau_n \mid n < \omega \rangle$ be an increasing sequence of regular cardinals, $\lambda = \cup_n \tau_n$, and $\theta > \lambda^+$ regular. A set $C \subseteq \mathcal{P}_\lambda(H_\theta)$ is a closed unbounded set if it contains all elementary substructures $M \prec \mathfrak{A}$ of size $|M| < \lambda$ of some algebra $\mathfrak{A} = (H_\theta, \in, f_n)_n$ on H_θ . We reformulate the definitions of Approachable Free Subset Property and Approachable Bounded Subset Property from the introduction.

- Definition 14.** 1. Let $F : [\lambda]^{<\omega} \rightarrow \lambda$ be a function. We say that a subset $X \subseteq \lambda$ is **free with respect to F** if for every $\gamma \in X$, $\gamma \notin F[X \setminus \{\gamma\}]^{<\omega}$.
2. The **Approachable Free Subset Property (AFSP)** with respect to $\vec{\tau}$ asserts that there exists a closed unbounded set $C \subseteq \mathcal{P}_\lambda(H_\theta)$ of structures $N \prec (H_\theta; \in)$ so that for every internally approachable structure $N \in C$ there exists some $m < \omega$ such that the set $\{\chi_N(\tau_n) \mid m \leq n < \omega\}$ is free with respect to every function $F \in N$
3. The **Approachable Bounded Subset Property (ABSP)** with respect to $\vec{\tau}$ asserts that there exists a closed unbounded set $C \subseteq \mathcal{P}_\lambda(H_\theta)$ of structures $N \prec (H_\theta; \in)$ so that for every internally approachable structure $N \in C$ there exists some $m < \omega$ such that for every $F \in N$, $F : [\lambda]^k \rightarrow \lambda$ of finite arity $k < \omega$, and distinct numbers $d, d_1, d_2, \dots, d_k \in \omega \setminus m$, if

$$F(\chi_N(\tau_{d_1}), \dots, \chi_N(\tau_{d_k})) < \tau_d$$

then

$$F(\chi_N(\tau_{d_1}), \dots, \chi_N(\tau_{c_d})) < \chi_N(\tau_d).$$

To see that the formulations in Definition 14 are equivalent to the ones given in the introduction, note that if $\theta > \lambda = \cup_n \tau_n$ is the first for which there exists a club $C \subseteq \mathcal{P}_\lambda(H_\theta)$ which is definable in $\vec{\tau}$ and consists of subalgebras $M \subseteq \mathfrak{A} = (H_\theta, \in, f_n)_n$, then for every $\theta' > \theta$ and $M' \prec H_{\theta'}$, if $\vec{\tau} \in M'$ then $\theta, C \in M'$ and $M' \cap \mathcal{P}_\lambda(H_\theta) \in C$.

Lemma 15. *Suppose that $\vec{\tau} = \langle \tau_n \mid n < \omega \rangle$ is an increasing sequence of regular cardinals.*

1. If there is no continuous essentially tree-like scale on $\prod_n \tau_n$ then there is no continuous tree-like scale on $\prod_n \tau_n$.
2. AFSP w.r.t $\vec{\tau}$ implies that there is no continuous tree-like scale on $\prod_n \tau_n$.
3. ABSP w.r.t $\vec{\tau}$ implies both
 - (i) AFSP w.r.t $\vec{\tau}$, and
 - (ii) there is no continuous essentially tree-like scale on $\prod_n \tau_n$.

Proof. 1. This is an immediate consequence of the definitions of an essentially tree-like scale and a tree-like scale on $\prod_n \tau_n$.

2. We prove the contrapositive statement, that if there exists a continuous tree-like scale on $\prod_n \tau_n$ then AFSP fails with respect to $\vec{\tau}$. Suppose that \vec{f} is a continuous tree-like scale on $\prod_n \tau_n$. Since \vec{f} is tree-like, we can assign to it a function $F : \lambda \rightarrow \lambda$, $\lambda = \cup_n \tau_n$, defined as follows: For every $n < \omega$ and μ , $\tau_n \leq \mu < \tau_{n+1}$, define

$$F(\mu) = \begin{cases} f_\alpha(n) & \text{if } \mu = f_\alpha(n+1) \text{ for some } \alpha < \eta \\ 0 & \text{otherwise.} \end{cases}$$

$F(\mu)$ is well defined, i.e., does not depend on the choice of α such that $\mu = f_\alpha(n+1)$, since \vec{f} is tree-like. It is clear from the definition of F that for every $\delta < \eta$ and $n < \omega$, $F(f_\delta(n+1)) = f_\delta(n)$. Now, if $C \subseteq \mathcal{P}_\lambda(H_\theta)$ is a closed unbounded subset, $N \in C$ is an internally approachable structure with $F \in N$, and $\delta = \chi_N(\eta)$, then $\chi_N^\vec{\tau}(n) = f_\delta(n)$ for all but finitely many $n < \omega$. Hence, for all but finitely many $n < \omega$, $F(\chi_N(\tau_{n+1})) = \chi_N(\tau_n)$, which means that $\{\chi_N(\tau_{n+1}), \chi_N(\tau_n)\}$ is not free with respect to $F \in N$. Since C was an arbitrary closed and unbounded subset, AFSP with respect to $\vec{\tau}$ fails.

3. The fact that ABSP implies AFSP is immediate from the definition of the two properties. To show that ABSP w.r.t $\vec{\tau}$ implies that there is no continuous scale on $\prod_n \tau_n$ which is essentially tree-like, we prove the contrapositive statement. Suppose that $\langle f_\alpha \mid \alpha < \eta \rangle$ is a continuous essentially tree-like scale on a product $\prod_n \tau_n$. Then by Definition 9 for every $n < \omega$, there is a function $C_n : \tau_{n+1} \rightarrow \mathcal{P}(\tau_n)$ so that for every $\mu < \tau_{n+1}$, $C_n(\mu)$ is a closed and unbounded subset of τ_n which is disjoint from $\{\mu_n < \tau_n \mid \exists \beta < \eta, f_\beta(n+1) = \mu \text{ and } f_\beta(n) = \mu_n\}$. Let C be any club of elementary substructures of $(H_\theta; \in)$. Take an internally approachable substructure $N \in C$ and of size $|N| < \lambda = \cup_n \tau_n$, so that both $\langle \tau_n \mid n < \omega \rangle$ and $\langle C_n \mid n < \omega \rangle$ belong to N . Define $\delta = \chi_N(\eta)$ and let $m < \omega$ so that $f_\delta(n) = \chi_N(\tau_n)$ for all $n \geq m$. Fixing $n \geq m$ and examining the elementary extension $N' = N[\{f_\delta(n+1)\}] = \{F(f_\delta(n+1)) \mid F \in N\} \prec (H_\theta; \in)$ of N , we have that $C_n(f_\delta(n+1)) \in N'$ since $C_n \in N$. Now, as $C_n(f_\delta(n+1)) \subseteq \tau_n$ is closed unbounded, we must have that $\chi_{N'}(\tau_n) \in C_n(f_\delta(n+1))$. However $\chi_N(\tau_n) = f_\delta(n) \notin C_n(f_\delta(n+1))$

by the definition of C_n . This implies that $\chi_{N'}(\tau_n) > \chi_N(\tau_n) = f_\delta(n)$, which in turn, implies that $F(f_\delta(n+1)) > f_\delta(n)$ for some $F \in N$. Since $N \in C$ where C is an arbitrary closed unbounded collection, and n is an arbitrarily large finite ordinal, we conclude that ABSF fails with respect to $\langle \tau_n \mid n < \omega \rangle$. □

1.2 Fine structure primer

1.2.1 Ultrafilters

We shall take our fine structure from [29]. Our result almost certainly also applies to different forms of fine structure such as the fine structure theory of [16], in fact, the proof of Theorem 6 in particular would be greatly simplified, but at the cost of significantly complicating the arguments in the core model part of this paper. As there is currently no account of the covering lemma for λ -indexing, we think it prudent to choose Mitchell-Steel mice at this time. We don't use [31], as $\neg O^\sharp$ is much too strong a limitation for this section. (While technically Mitchell and Steel operate under the assumption of $\neg M_1^\#$ in [29], it is well understood by now that their fine structure theory functions well past this point.)

For our purposes an extender F is a directed system of ultrafilters $\{(a, X) \mid a \in [\text{lh}(F)]^{<\omega}, X \subset [\text{crit}(F)]^{|a|}\}$ as described in [15, p. 384]. The individual ultrafilters will be denoted as $F_a := \{X \subset \text{crit}(F)^{|a|} \mid (a, X) \in F\}$. For $a \subset b$ and f a function with domain $[\text{crit}(F)]^{|a|}$, we let $f^{a,b}$ be the function with domain $[\text{crit}(F)]^{|b|}$ determined by $f^{a,b}(\bar{b}) = f(\bar{a})$ where \bar{a} is the unique subset of \bar{b} determined by the type of a and b . This gives rise to an embeddings from $\text{Ult}(\mathcal{M}, F_a)$ into $\text{Ult}(\mathcal{M}, F_b)$. The direct limit along those embeddings is the extender ultrapower $\text{Ult}(\mathcal{M}, F)$, elements of which we will present as pairs $[f, a]_F^\mathcal{M}$ where $f \in \mathcal{M}$ is a function with domain $[\text{crit}(F)]^{|a|}$ and $a \in [\text{lh}(F)]^{<\omega}$. The direct limit map shall be denoted $\iota_F^\mathcal{M} : \mathcal{M} \rightarrow \text{Ult}(\mathcal{M}, F)$. We will generally omit the superscript in this notation. This should not lead to confusion. Note that we will later form ultrapowers where some functions involved in the construction are not elements of the structure but merely definable over it.

$\beta < \text{lh}(F)$ is a generator of F if it cannot be represented as $[f, a]_F$ for any $f \in \text{crit}(F)^{\text{crit}(F)} \cap \mathcal{M}$ and $a \in [\beta]^{<\omega}$, i.e. $\{b \cup \{\xi\} \mid f(b) = \xi\} \notin F_{a \cup \{\beta\}}$. Let $\text{gen}(F)$ denote the strict supremum of the generators of F . Also let $\nu(F) = \max\{\text{gen}(F), (\text{crit}(F)^+)^{\mathcal{M}}\}$.

For a subset A of α we will write $F \upharpoonright A := \{(a, X) \in F \mid a \subset A\}$. We will consider this an extender, forming ultrapowers etc, even if A is not an ordinal. Let $\eta < \alpha$ be such that $\eta = \text{gen}(F \upharpoonright \eta)$, then the trivial completion is the $(\text{crit}(F), (\eta^+)^{\text{Ult}(\mathcal{M}; F \upharpoonright \eta)})$ -extender derived from $\iota_{F \upharpoonright \eta}$.

1.2.2 Premice

A potential premouse is a structure of the form $\mathcal{M} = \langle J_\alpha^{\vec{E}}; \in, \vec{E}, F \rangle$ where $J_\alpha^{\vec{E}}$ is a model constructed from a sequence of extenders \vec{E} using the Jensen hierarchy. For $\beta \leq \alpha$ we define $\mathcal{M}|\beta := (J_\beta^{\vec{E} \upharpoonright \beta}; \in, \vec{E} \upharpoonright \beta, \vec{E}_\beta)$ and $\mathcal{M}||\beta := (J_\beta^{\vec{E} \upharpoonright \beta}; \in, \vec{E} \upharpoonright \beta)$. (The difference between the two notations lies in including a top predicate.) If \mathcal{N} is of one of the above forms then we write $\mathcal{N} \trianglelefteq \mathcal{M}$ and say \mathcal{N} is an initial segment of \mathcal{M} .

\vec{E} must be good, i.e. it has the following properties:

- (Idx) for all $\beta < \alpha$ if $\vec{E}_\beta \neq \emptyset$, then $\beta = (\nu(\vec{E}_\beta)^+)^{\text{Ult}(\mathcal{M}|\beta; \vec{E}_\beta)}$;
- (Coh) for all $\beta < \alpha$ if $\vec{E}_\beta \neq \emptyset$, then $\mathcal{M}||\beta = \text{Ult}(\mathcal{M}|\beta; \vec{E}_\beta)|\beta$;
- (ISC) for all $\beta < \alpha$ if $\vec{E}_\beta \neq \emptyset$, then for all $\eta < \alpha$ such that $\eta = \text{gen}(\vec{E}_\beta \upharpoonright \eta)$ the trivial completion of $\vec{E}_\beta \upharpoonright \eta$ is on \vec{E} or $\vec{E}_\eta \neq \emptyset$ and it is on $\iota_{\vec{E}_\eta}(\vec{E})$.

Note that \vec{E}_β measures exactly those subsets of its critical point that are in $\mathcal{M}||\beta$ for any $\beta < \alpha$ such that $\vec{E}_\beta \neq \emptyset$. F the top extender must be such that $\vec{E} \hat{\ } F$ remains good. F can be empty in which case \mathcal{M} is called passive, otherwise \mathcal{M} is active.

To an active potential premouse we associate three constants: $\mu^{\mathcal{M}}$ the critical point of the top extender; $\nu^{\mathcal{M}}$ the strict supremum of the generators of \mathcal{M} 's top extender or $(\mu^{\mathcal{M}})^+$ whichever is larger; $\gamma^{\mathcal{M}}$ the index of the longest initial segment of \mathcal{M} 's top extender (if it exists).

We distinguish three different types of active potential premouse: \mathcal{M} is active type I if $\nu^{\mathcal{M}} = (\mu^{\mathcal{M},+})^{\mathcal{M}}$; \mathcal{M} is active type II if $\nu^{\mathcal{M}}$ is a successor ordinal; \mathcal{M} is an active type III if it is neither type I or type II, i.e. the set of the generators of \mathcal{M} 's top extender has limit type.

1.2.3 Fine structure

A disadvantage of Mitchell-Steel indexing is that we cannot deal directly with definability over \mathcal{M} , but instead need to work with an amenable code of our original structure. The exact nature of this coding is dependant on the type of \mathcal{M} . We will take inspiration from [28] and use a uniform notation $\mathcal{C}_0(\mathcal{M})$ for this code.

If $\mathcal{M} := \langle |\mathcal{M}|; \in, \vec{E}, F \rangle$ is an active potential premouse of type I or II, we will define an alternative predicate F^c coding the top extender F : F^c consists of tuples (γ, ξ, a, X) such that $\xi \in (\mu^{\mathcal{M}}, (\mu^{\mathcal{M},+})^{\mathcal{M}})$ and $\gamma \in (\nu(F), \text{On} \cap |\mathcal{M}|)$ is such that $(F \cap ([\nu(F)]^{<\omega} \times \mathcal{M}||\xi)) \in \mathcal{M}||\gamma$, and $(a, X) \in (F \cap ([\gamma]^{<\omega} \times \mathcal{M}||\xi))$. The point is that F^c is amenable. We let $\mathcal{C}_0(\mathcal{M}) := \langle |\mathcal{M}|; \in, \vec{E}, F^c \rangle$.

If \mathcal{M} on the other hand is active type III we have to make bigger changes. In the language of [29] we have to “squash”, that is remove ordinals from the structure. (This is to ensure that the initial segment condition is preserved by iterations.) We let $\mathcal{C}_0(\mathcal{M}) := \langle J_{\nu(F)}^{\vec{E}}; \in, \vec{E} \upharpoonright \nu(F), F \upharpoonright \nu(F) \rangle$.

We then define $r\Sigma_1$ -formulae to be Σ_1 over $\mathcal{C}_0(\mathcal{M})$, and $r\Sigma_{n+1}$ -formulae to be Σ_1 in a predicate coding an appropriate segment of the $r\Sigma_n$ -theory of $\mathcal{C}_0(\mathcal{M})$. We will let $\text{Th}_n^{\mathcal{M}}(\alpha, q) := \{([\phi], b) \mid \phi \text{ is } r\Sigma_n, b \in [\alpha]^{<\omega}, \mathcal{C}_0(\mathcal{M}) \models \phi(b, q)\}$.

Projecta can then be defined relative to these formulas, i.e. $\rho_{n+1}(\mathcal{M})$ is the least ordinal such that some $r\Sigma_{n+1}$ -definable (in parameters) subset of it is not in $\mathcal{C}_0(\mathcal{M})$. $\rho_0(\mathcal{M}) = \text{On} \cap \mathcal{C}_0(\mathcal{M})$ (which might be smaller than $\text{On} \cap \mathcal{M}$).

As usual we define $p_{n+1}(\mathcal{M})$, the $(n+1)$ -th standard parameter, to be the lexicographically least $p \in [\text{On} \cap \mathcal{C}_0(\mathcal{M}) \setminus \rho_{n+1}(\mathcal{M})]^{<\omega}$ that defines a missing subset of $\rho_{n+1}(\mathcal{M})$.

We can also define canonical $r\Sigma_{n+1}$ -Skolem function allowing us to form $\text{Hull}_{n+1}^{\mathcal{M}}(A)$ given a subset A of $\mathcal{C}_0(\mathcal{M})$. Note that while our notation makes it look like a hull of \mathcal{M} it is a substructure of $\mathcal{C}_0(\mathcal{M})$ not \mathcal{M} .

We say \mathcal{M} is n -sound above β relative to p iff $\mathcal{C}_0(\mathcal{M}) = \text{Hull}_n^{\mathcal{M}}(\beta \cup \{p\})$. We will not mention the parameter if \mathcal{M} is n -sound above β relative to $p_n(\mathcal{M})$. If \mathcal{M} is n -sound above $\rho_n(\mathcal{M})$, we simply say that \mathcal{M} is n -sound.

A potential premouse is then a premouse if all its initial segments are n -sound for all n . We can now also define fine structural ultrapowers. Let \mathcal{M} be a premouse and let F be an extender that measure all subsets of its critical point in \mathcal{M} . Let n be such that $\text{crit}(F) < \rho_n(\mathcal{M})$ and \mathcal{M} is n -sound. Then $\text{Ult}_n(\mathcal{M}, F)$ is the ultrapower formed using all equivalence classes $[f, a]_F$ where $a \in [\text{lh}(F)]^{<\omega}$ and f is a function with domain $[\text{crit}(F)]^{|a|}$ that is $r\Sigma_n$ -definable over \mathcal{M} (in parameters).

Lemma 16. *Let \mathcal{M} be a premouse, and let $\kappa \in \mathcal{C}_0(\mathcal{M})$ be a regular cardinal there. Assume $\rho_{n+1}(\mathcal{M}) \leq \beta < \kappa \leq \rho_n(\mathcal{M})$ for some n such that \mathcal{M} is $(n+1)$ -sound above β . Then $\text{cof}(\kappa) = \text{cof}(\rho_n(\mathcal{M}))$.*

Proof. For $\xi < \rho_n(\mathcal{M})$ we let \mathcal{N}_ξ be the structure $\mathcal{M} \upharpoonright \xi$ with $\text{Th}_n^{\mathcal{M}}(\xi, p_n(\mathcal{M}))$ as an additional predicate. Let then κ_ξ be the supremum of ordinals less than κ which are Σ_1 -definable over \mathcal{N}_ξ from $p_{n+1}(\mathcal{M})$ and ordinals less than β . As all objects involved are elements of \mathcal{M} , we must have $\kappa_\xi < \kappa$. On the other hand $\sup_{\xi < \rho_n(\mathcal{M})} \kappa_\xi = \kappa$ as \mathcal{M} was $(n+1)$ -sound above β . \square

An additional fact that we will need is that if \mathcal{M} is an active (potential) premouse, then $\text{cof}(\text{On} \cap \mathcal{M}) = \text{cof}((\mu^{\mathcal{M},+})^{\mathcal{M}})$. See the last remark of Chapter 1 in [29].

1.2.4 Iterability

A (normal, ω -maximal) iteration tree on a premouse \mathcal{M} is a tuple $\mathcal{T} := \langle \langle \mathcal{M}_\alpha^{\mathcal{T}} : \alpha \leq \text{lh}(\mathcal{T}) \rangle, \langle E_\alpha^{\mathcal{T}} : \alpha < \text{lh}(\mathcal{T}) \rangle, D^{\mathcal{T}}, \langle \iota_{\alpha,\beta}^{\mathcal{T}} : \alpha \leq_{\mathcal{T}} \beta \leq \text{lh}(\mathcal{T}) \rangle \rangle$ where $\mathcal{M}_\alpha^{\mathcal{T}}$ is a premouse for all $\alpha \leq \text{lh}(\mathcal{T})$ ($\mathcal{M}_0^{\mathcal{T}} = \mathcal{M}$); $E_\alpha^{\mathcal{T}}$ is an extender from the $\mathcal{M}_\alpha^{\mathcal{T}}$ -sequence for all $\alpha < \text{lh}(\mathcal{T})$, $\alpha < \beta$ implies $\text{lh}(E_\alpha^{\mathcal{T}}) < \text{lh}(E_\beta^{\mathcal{T}})$; $\iota_{\alpha,\beta}^{\mathcal{T}} : \mathcal{C}_0(\mathcal{M}_\alpha^{\mathcal{T}}) \rightarrow \mathcal{C}_0(\mathcal{M}_\beta^{\mathcal{T}})$ is the (possibly) partial iteration map for all $\alpha \leq_{\mathcal{T}} \beta \leq \text{lh}(\mathcal{T})$, it is total iff $D^{\mathcal{T}} \cap (\alpha, \beta]_{\leq_{\mathcal{T}}} \neq \emptyset$; $\leq_{\mathcal{T}}$ is the tree order on

$\text{lh}(\mathcal{T})$ with root 0, if $\gamma + 1 \leq \text{lh}(\mathcal{T})$, then the \mathcal{T} -predecessor is the least β such that $\text{crit}(E_\gamma^\mathcal{T}) < \text{gen}(E_\beta^\mathcal{T})$, in that case $(\mathcal{M}_{\gamma+1}^\mathcal{T})^*$ is the segment of $\mathcal{M}_\beta^\mathcal{T}$ to which $E_\gamma^\mathcal{T}$ is applied, if $\lambda \leq \text{lh}(\mathcal{T})$ is a limit, then $b_\lambda^\mathcal{T} := [0, \lambda]_{\leq \mathcal{T}}$ is a cofinal branch whose intersection with $D^\mathcal{T}$ is finite, $\mathcal{M}_\lambda^\mathcal{T}$ must be the direct limit of $\langle \mathcal{M}_\alpha^\mathcal{T}, \iota_{\alpha, \beta}^\mathcal{T} : \alpha \leq \mathcal{T} \beta \in b_\lambda^\mathcal{T} \rangle$; finally, $\gamma + 1 \in D^\mathcal{T}$ if and only if $(\mathcal{M}_{\gamma+1}^\mathcal{T})^* \neq \mathcal{M}_\beta^\mathcal{T}$. For an ordinal γ , a γ iteration strategy Σ for a premouse \mathcal{M} is a function such that $\Sigma(\mathcal{T})$ is a cofinal branch b in \mathcal{T} for which the direct limit $\mathcal{M}_b^\mathcal{T}$ along the branch is wellfounded branch for every iteration tree on \mathcal{T} of limit length $< \gamma$ and with the property that $\Sigma(\mathcal{T} \upharpoonright \alpha) = [0, \alpha]_{\leq \mathcal{T}}$ for all limit $\alpha < \text{lh}(\mathcal{T})$. \mathcal{M} is γ -iterable if there exists a γ -iteration strategy for \mathcal{M} . We will just say \mathcal{M} is iterable if it is γ -iterable for all ordinals γ .

Let \mathcal{M} be a premouse, $n < \omega$ and let $p_{n+1}(\mathcal{M}) = \langle \xi_0, \dots, \xi_{k-1} \rangle$ listed in a decreasing order. The $(n+1)$ -th solidity witness $w_{n+1}(\mathcal{M})$ is a tuple $\langle t_0, \dots, t_{k-1} \rangle$ where

$$t_i = \text{Th}_{n+1}^{\mathcal{M}}(\xi_i, \langle \xi_0, \dots, \xi_{i-1} \rangle).$$

We say \mathcal{M} is $(n+1)$ -solid if $w_{n+1}(\mathcal{M}) \in \mathcal{C}_0(\mathcal{M})$.

A core result of [29] is that any reasonably iterable n -sound premouse is $(n+1)$ -solid. Mitchell-Steel also showed the following with similar methods, see the remark after Theorem 8.2. Note that the requirement for unique branches can be replaced by the weak Dodd-Jensen property from [28].

Lemma 17 (Condensation Lemma). *Let $\mathcal{M} := (|\mathcal{M}|; \in, \vec{E}, F)$ be a $(n+1)$ -sound premouse such that every countable hull of \mathcal{M} has a $(\omega_1 + 1)$ -iteration strategy. Let \mathcal{N} be a premouse such that there exist an $r\Sigma_{n+1}$ -elementary embedding $\pi : \mathcal{C}_0(\mathcal{N}) \rightarrow \mathcal{C}_0(\mathcal{M})$ with $\text{crit}(\pi) \geq \rho_{n+1}(\mathcal{N})$. Then \mathcal{N} is an initial segment of \mathcal{M} or of $\text{Ult}(\mathcal{M}, \vec{E}_{\text{crit}(\pi)})$.*

Both these results use the notion of a phalanx (although this notion was not yet fully developed by the time of [29]) of which we too will have need. A phalanx is a tuple $\langle \langle \mathcal{M}_i : i \leq \alpha \rangle, \langle \kappa_i : i < \alpha \rangle \rangle$ where \mathcal{M}_i agrees with \mathcal{M}_j up to $(\kappa_i^+)^{\mathcal{M}_j}$ for all $i < j \leq \alpha$.

Phalanxes are a natural byproduct of iteration trees, i.e. if \mathcal{T} is a normal iteration tree on some premouse, then $\langle \langle \mathcal{M}_i^\mathcal{T} : i \leq \text{lh}(\mathcal{T}) \rangle, \langle \nu(E_i^\mathcal{T}) : i < \text{lh}(\mathcal{T}) \rangle \rangle$ is a phalanx.

We can then also define iterability on phalanxes as a natural extension of the structure of iteration trees. Given a phalanx $\langle \langle \mathcal{M}_i : i \leq \alpha \rangle, \langle \kappa_i : i < \alpha \rangle \rangle$ and an extender E we can extend the phalanx by applying E to \mathcal{M}_i where i is minimal with $\text{crit}(E) < \kappa_i$. (Note we have to require that the length of E is above $\sup_{i < \alpha} \kappa_i$ to maintain “normality”.)

A notion of iteration then follows naturally. The most critical difference here is that we have to keep track above which element of the phalanx any given model of the iteration tree lies. The art of phalanx iteration lies in arranging things such that the last model of a co-iteration lies above the “right” model.

2 Forcing the Approachable Bounded Subset Property

Our forcing notations are mostly standard. We use the Jerusalem forcing convention by which “a condition p extends (is more informative than) q ” is denoted by $p \geq q$. In general, names for a set x in a generic extension will be denoted by \dot{x} . If x is in the ground model then its canonical name is denoted by \check{x} .

We denote our initial ground model by V' , which we assume to satisfy the following assumptions: there are two increasing sequences $\langle \kappa_n \mid n < \omega \rangle$, $\langle \lambda_n \mid n < \omega \rangle$ of regular cardinals, with $\lambda_n < \kappa_{n+1} < \lambda_{n+1}$ for all n , and that each λ_n is measurable of Mitchell order $o(\lambda_n) = \kappa_n$.

For each $n < \omega$, let $\langle U_{\lambda_n, \alpha} \mid \alpha < \kappa_n \rangle$ be a \triangleleft -increasing of normal measures on λ_n . I.e., $U_{\lambda_n, \alpha}$ belongs to the ultrapower by $U_{\lambda_n, \beta}$, whenever $\alpha < \beta$. Denote $\lambda = \cup_n \lambda_n$.

In order to apply our main extender-based forcing notion, we first force with a preparatory forcing \mathbb{P}' over V' to transform the Mitchell-order increasing sequences $\langle U_{\lambda_n, \alpha} \mid \alpha < \kappa_n \rangle$ of normal measures, to Rudin-Keisler increasing sequences. For this, we force with a Gitik-iteration \mathbb{P}' ([10]) for changing the cofinality of measurable cardinals between the cardinals (κ_n, λ_n) for all $n < \omega$. Let $G' \subseteq \mathbb{P}'$ be a generic filter over V' , and set $V = V[G']$. We list a number of facts concerning the extensions in V of the measures $\langle U_{\lambda_n, \alpha} \mid \alpha < \kappa_n \rangle$ from V' . The analysis leading to these facts can be found in [10], or [3] for a similar type of poset. The Mitchell-order increasing sequence $\langle U_{\lambda_n, \alpha} \mid \alpha < \kappa_n \rangle$ extends to a Rudin-Keisler increasing sequence of λ_n -complete measures $\langle U_{\lambda_n, \alpha}^* \mid \alpha < \kappa_n \rangle$, with Rudin-Keisler projections $\pi_{\beta, \alpha}^n : \lambda_n \rightarrow \lambda_n$ for each $\alpha < \beta < \kappa_n$. We note that the least measure $U_{\lambda_n, 0}^*$ remains normal. We denote for each $n < \omega$ the linear directed system of measures $\{U_{\lambda_n, \alpha}^*, \pi_{\beta, \alpha}^n \mid \alpha \leq \beta < \kappa_n\}$ by E_n , and further denote each $U_{\lambda_n, \alpha}^*$ by $E_n(\alpha)$. Let

$$j_{E_n} : V \rightarrow M_{E_n} = \text{Ult}(V, E_n) = \text{dirlim}_{\alpha < \kappa_n} \text{Ult}(V, E_n(\alpha))$$

Each measure $E_n(\alpha)$ can be derived from j_{E_n} using a generator $\gamma_\alpha^{E_n} < j_{E_n}(\lambda_n)$. The following list summarizes the key properties of the extenders E_n (see [11]):

Fact 18.

1. $\text{cp}(j_{E_n}) = \lambda_n$ and $M_{E_n}^{< \kappa_n} \subseteq M_{E_n}$
2. $\gamma_0^{E_n} = \lambda_n$ and $\langle \gamma_\alpha^{E_n} \mid \alpha < \kappa_n \rangle$ is a strictly increasing and continuous sequence
3. $\gamma^{E_n} = \sup_{\alpha < \kappa_n} \gamma_\alpha^{E_n}$ is strongly inaccessible in M_{E_n} , and we may assume that there exists a function $g_n : \lambda_n \rightarrow \lambda_n$ such that $\gamma^{E_n} = j_{E_n}(g_n)(\lambda_n)$
4. for each $\alpha < \beta < \kappa_n$, $E_n(\alpha)$ is strictly weaker than $E_n(\beta)$ in the Rudin-Keisler order, i.e., for every $A \in E_n(\alpha)$ there is $\nu \in A$ such that $\pi_{\beta, \alpha}^{-1}(\{\nu\})$ is unbounded in λ_n .

5. for every $\alpha < \kappa_n$ and $h : \lambda_n \rightarrow \lambda_n$ such that $j_{E_n}(h)(\gamma_\alpha^{E_n}) < \gamma^{E_n}$.
 $j_{E_n}(h)(\gamma_\alpha^{E_n}) < \gamma_\beta^{E_n}$ for all $\beta > \alpha$.

Next, we force over V with a short extender-based-type forcing \mathbb{P} , associated with the extenders E_n , $n < \omega$. \mathbb{P} is a variant of the forcing in [2]. Extending the arguments of [2], we focus here on the generic scale associated with the extender-based-forcing, and use it to analyze the possible internally approachable structures in the generic extensions. This approach follows the one taken in [1], where an extender-based forcing has been used to obtain results concerning internally-approachable structures witnessing that ground model sequences $\langle S_n \mid n < \omega \rangle$ being tightly-stationary.

Definition 19. Conditions $p \in \mathbb{P}$ are sequences $p = \langle p_n \mid n < \omega \rangle$ such that there is some $\ell < \omega$ for which the following requirements hold:

1. for $n < \ell$, $p_n = \langle f_n \rangle$, where $f_n : \lambda^+ \rightarrow \lambda_n$ is a partial function of size $|f_n| \leq \lambda$, with $0 \in \text{dom}(f_n)$ and both $f_n(0), g_n(f_n(0)) < \lambda_n$ are strongly inaccessible cardinals
2. For $n \geq \ell$, $p_n = \langle f_n, a_n, A_n \rangle$, where f_n is as above, $a_n : \lambda^+ \rightarrow \kappa_n$ is a partial continuous and order-preserving function, whose domain d_n is a closed and bounded set of λ^+ of has size $|a_n| < \kappa_n$.
We define $\text{mc}(a_n)$ to be $a_n(\max(d_n)) = \max(\text{rng}(a_n))$, and require that the set A_n to be contained in $\lambda_n \setminus \lambda_{n-1}$ and belong to $E_n(\text{mc}(a_n))$.
3. $\text{dom}(a_n) \cap \text{dom}(f_n) = \emptyset$ and $\text{dom}(a_n) \subseteq \text{dom}(a_{n+1})$ for every $n \geq \ell$, $a_n(0) = 0$, and for every $\delta \in \cup_n \text{dom}(f_n)$ there exists some $m < \omega$ such that $\delta \in \text{dom}(a_m)$.

For a condition $p \in \mathbb{P}$ as above, we denote ℓ, f_n, a_n, A_n by $\ell^p, f_n^p, a_n^p, A_n^p$ respectively. Direct extensions and end-extensions of conditions are defined as follows. A condition p^* is a direct extension of p , if $\ell^{p^*} = \ell^p$, $f_n^p \subseteq f_n^{p^*}$ for all $n < \omega$, and $a_n^p \subseteq a_n^{p^*}$, $A_n^{p^*} \subseteq (\pi_{\text{mc}(a_n^{p^*}), \text{mc}(a_n^p)}^n)^{-1} A_n^p$ for all $n \geq \ell^p$.

For every $\nu \in A_n^p$, define $p_n \frown \langle \nu \rangle = \langle f'_n \rangle$, where

$$f'_n = f_n^{p_n} \cup \{ \langle \alpha, \pi_{\text{mc}(a_n^p), a_n(\alpha)}^n(\nu) \rangle \mid \alpha \in \text{dom}(a_n^p) \}.$$

If $\vec{\nu} = \langle \nu_{\ell^p}, \dots, \nu_{n-1} \rangle$ belong to $\prod_{i=\ell^p}^{n-1} A_i^p$, we define the end extension of p by $\vec{\nu}$, denoted $p \frown \vec{\nu}$, to be the condition $p' = \langle p'_n \mid n < \omega \rangle$, defined by $p'_k = p_k$ for every $k \notin \{ \ell^p, \dots, n-1 \}$, and $p'_k = p_k \frown \langle \nu_k \rangle$ otherwise. A condition $q \in \mathbb{P}$ extends p if q is obtained from p by a finite sequence of end-extensions and direct extensions. Equivalently, q is a direct extension of an end-extension $p \frown \vec{\nu}$ of p . Following the Jerusalem forcing convention, we write $p \geq q$ if p extends q , and $p \geq^* q$ if p is a direct extension of q .

Notation 20. We introduce the following notational convention for the Rudin-Keisler projections $\pi_{\alpha, \beta}^n$ to be applied in the context of the forcing \mathbb{P} . Let p be a condition and $\nu \in A_n^p$ for some $n \geq \ell^p$ and $\alpha \in \text{dom}(a_n^p)$. We write $\pi_{\text{mc}(p), \alpha}^p(\nu)$ for $\pi_{\text{mc}(a_n^p), a_n(\alpha)}^n(\nu)$.⁷ Similarly, for a sequence $\vec{\nu} = \langle \nu_i \mid \ell^p \leq i < n \rangle \in \prod_{\ell^p \leq i < n} A_i^p$,

⁷Note that the index n is determined from the fact that $\nu \in A_n^p \subseteq \lambda_n \setminus \lambda_{n-1}$.

we write $\pi_{\text{mc}(p),\alpha}^p(\vec{\nu})$ for the projected sequence $\langle \pi_{\text{mc}(p),\alpha}^p(\nu_i) \mid \ell^p \leq i < n \rangle$.

We proceed to list several standard basic properties of the poset \mathbb{P} , referring the reader to [2] for details.

Lemma 21.

1. $(\mathbb{P}, \leq, \leq^*)$ is a Prikry-type forcing
2. for each $p \in \mathbb{P}$, the direct extension order \leq^* of \mathbb{P}/p is κ_{ℓ^p} -closed
3. \mathbb{P} satisfies the λ^{++} -c.c
4. (Strong Prikry Property) Let $D \subseteq \mathbb{P}$ be a dense open set. For every $p \in \mathbb{P}$ there are $p^* \geq^* p$ and $n < \omega$, such that for every $\vec{\nu} \in \prod_{\ell^p \leq i < n} A_i^{p^*}$, $p^* \frown \vec{\nu} \in D$.

It is routine to verify that the above properties imply that \mathbb{P} does not add new bounded subsets to λ , and does not collapse λ^{++} . We extend our analysis of \mathbb{P} below to show that it preserves λ^+ . This result can be also derived using a standard application of the Weak Covering Theorem. To extend our study of the poset \mathbb{P} , we introduce a notation of orderings \leq^m , $m < \omega$, which refine the direct extension ordering \leq^* .

Notation 22. Let p, q be two conditions in \mathbb{P} . For $m < \omega$ we write $p \leq^m q$ if $p \leq^* q$ and $a_n^p = a_n^q$, $A_n^p = A_n^q$ for all $n < m$.

Therefore, for each $m < \omega$, \leq^m is κ_m -closed and $\leq^{m+1} \subseteq \leq^m$.

Lemma 23. Let $\theta > \lambda^+$ regular, \triangleleft be a well-ordering of H_θ , and $M \triangleleft (H_\theta; \in, \triangleleft)$ satisfying $\mathbb{P} \in M$ and $|M| = \lambda, V_\lambda \subseteq M$. Suppose that there exists an enumeration $\vec{D} = \langle D_\mu \mid \mu < \lambda \rangle$ of all dense open subsets of \mathbb{P} in M , so that $\vec{D} \upharpoonright \nu \in M$ for every $\nu < \lambda$. Then for every condition $p \in \mathbb{P} \cap M$ and $\ell^*, \ell^p \leq \ell^* < \omega$, there exists $p^* \geq^{\ell^*} p$ so that for each dense open set $D \in M$ there are -

- $q \in M$ with $p \leq^{\ell^*} q \leq^{\ell^*} p^*$,
- a finite ordinal $n^D < \omega$, and
- a function $N^D : \prod_{\ell^p \leq i < n^D} A_i^q \rightarrow \omega$,

such that for every pair of sequences $\vec{\nu}^1, \vec{\nu}^2$, satisfying

$$\vec{\nu}^1 \in \prod_{i=\ell^p}^{n^D-1} A_i^q, \text{ and } \vec{\nu}^2 \in \prod_{i=n^D}^{N^D(\vec{\nu}^1)} A_i^q,$$

the condition $q \frown \vec{\nu}^1 \frown \vec{\nu}^2$ belongs to D .

Remark 24. We note that the condition $q \frown \vec{v}^1 \frown \vec{v}^2$ in the statement of the Lemma belongs to M as $V_\lambda \subseteq M$. Therefore, Lemma 23 implies that p^* is a generic condition for (M, \mathbb{P}) , namely, it forces the statement $\dot{G} \cap \dot{M} \cap \dot{D} \neq \emptyset$ for every dense open set $D \in M$.

Proof. We assume for notational simplicity that $\ell^p = 0$. The proof for the general case is similar. We fix for each $n < \omega$ a bijection $\psi_n : \lambda_n \rightarrow [\lambda_n]^{n+1} \times \lambda_n$ in M . Our final condition p^* will be obtained as a limit of a carefully constructed sequence $\langle p^n \mid \ell^* \leq n < \omega \rangle$, starting from $p^{\ell^*} = p$, and consisting of conditions in M . Moreover, it will satisfy $p^n \leq^{n+1} p^{n+1}$ for all $n \geq \ell^*$. Suppose that p^n has been defined for some $n \geq \ell^*$. Our goal is to construct an extension $p^{n+1} \geq^{n+1} p^n$, so that for every ordinal μ , $\lambda_{n-1} \leq \mu < \lambda_n$ there exists a function $N^{D_\mu} : \prod_{i \leq n} A_i^{p^{n+1}} \rightarrow \omega$ so that for every

$$\vec{v}^1 \in \prod_{i \leq n} A_i^{p^{n+1}} \text{ and } \vec{v}^2 \in \prod_{n+1 \leq i < N^{D_\mu}(\vec{v}^1)} A_i^{p^{n+1}}$$

$p^{n+1} \frown \vec{v}^1 \frown \vec{v}^2 \in D_\mu$. We note that this will guarantee $n^{D_\mu} = n + 1$ for $\lambda_n \leq \mu < \lambda_{n+1}$. p^{n+1} will be constructed from p^n in $(\lambda_n + 1)$ -many steps, using two sequences of condition parts, $\langle \vec{f}^i \mid i \leq \lambda_n \rangle$ and $\langle q^i \mid i \leq \lambda_n \rangle$ which satisfy the following requirements:

1. $\vec{f}_i = \langle f_{i,0}, f_{i,1}, \dots, f_{i,n} \rangle \in M$ is an $(n+1)$ -tuple, consisting of Cohen functions, $f_{i,k} : \lambda^+ \rightarrow \lambda_k$ of size at most λ . For $i = 0$, $\vec{f}_0 = \langle f_0^{p^n}, \dots, f_n^{p^n} \rangle$ is the tuple of the Cohen functions of the first $(n+1)$ Cohen components of p^{n+1} .
2. For each $k \leq n$, the sequence $f_{i,k}, i \leq \lambda_n$ is increasing in \subseteq , and $\text{dom}(f_{i,k}) \cap \text{dom}(a_k^{p^n}) = \emptyset$ for all $i \leq \lambda_n$.
3. $q^i = \langle q_m^i \mid n < m < \omega \rangle$ consists of tail segments of conditions in \mathbb{P} starting from the $(n+1)$ -th component, and $q^0 = p^n \setminus n + 1 = \langle p_m^n \mid n < m < \omega \rangle$.
4. the sequence $q^i, i \leq \lambda_n$ will be \leq^* -increasing in the obvious sense.

The construction of the two sequences will be internal to M , and definable from $p^n, \vec{D} \upharpoonright \lambda_{n+1}$, and using the fixed well-ordering \triangleleft of H_θ . Let $\delta \leq \lambda_n$ and suppose that $\langle q^i, \vec{f}_i \mid i < \delta \rangle$ has been defined and belongs to M . If δ is a limit ordinal, we define $\vec{f}_\delta = \langle f_{\delta,k} \mid k \leq n \rangle$ by $f_{\delta,k} = \bigcup_{i < \delta} f_{i,k}$. Similarly, q_δ is taken to be the supremum in the direct extension ordering of $q^i, i < \delta$, which is possible due to the fact that for each $k > n$, the direct extension ordering of the k -th components q_k^i , is κ_{k+1} -closed, and $\kappa_{k+1} > \lambda_n$. Therefore, for every $k > n$, we define $q_k^\delta = (f_k^{q^\delta}, a_k^{q^\delta}, A_k^{q^\delta})$ where

$$f_k^{q^\delta} = \bigcup_{i < \delta} f_k^{q^i}, a_k^{q^\delta} = \bigcup_{i < \delta} a_k^{q^i} \cup \{(\alpha, \gamma)\}, \text{ and } A_k^{q^\delta} = \bigcap_{i < \delta} (\pi_{\gamma, \text{mc}(a_k^{q^i})}^n)^{-1} A_k^{q^i},$$

where $\alpha = \sup \left(\bigcup_{i < \delta} \text{dom}(a_k^{q^i}) \right)$, and $\gamma = \sup \left(\bigcup_{i < \delta} \text{rng}(a_k^{q^i}) \right)$. Clearly, $q^\delta \in M$. Suppose now that $\delta = i + 1$ is a successor ordinal. We appeal to our fixed bijection $\psi_n : \lambda_n \rightarrow [\lambda_n]^{n+1} \times \lambda_n$, and consider $\psi_n(i) = (\vec{\nu}^i, \mu^i)$, where $\vec{\nu}^i \in [\lambda_n]^{n+1}$ and $\mu^i < \lambda_n$. We proceed as follows: If $\vec{\nu}^i \notin \prod_{i \leq n} A_i^{p^n}$ we make no change, setting $\vec{f}_\delta = \vec{f}_i$ and $q^\delta = q^i$. Otherwise, $\vec{\nu}^i \in \prod_{i \leq n} A_i^{p^n}$ and we consider the associated functions $\langle g_{i,0}, \dots, g_{i,n} \rangle$, defined by

$$g_{i,k} = \{ \langle \alpha, \pi_{\text{mc}(a_k^{p^n}), a_k^{p^n}}(\nu) \rangle \mid \alpha \in \text{dom}(a_k^{p^n}) \}.$$

We note that since $\text{dom}(g_{i,k}) = \text{dom}(a_k^{p^n})$, it is disjoint from $\text{dom}(f_{i,k})$, and we can therefore take their unions $f_{i,k}^* = f_{i,k} \cup g_{i,k}$, $k < n$ to define a sequence of functions $\vec{f}_i^* = \langle f_{i,0}^*, \dots, f_{i,n}^* \rangle$. By concatenating the $(n+1)$ -sequence \vec{f}_i^* with the tail q^i , we get a condition $q_*^i = \vec{f}_i^* \frown q^i \in \mathbb{P}$ with $\ell^{q_*^i} = n+1$, to which we apply the last clause of Lemma 21 (Strong Prikry Property) and find a direct extension $q_{**}^i \geq^* q_*^i$ and an integer N so that for every $\vec{\nu} \in \prod_{k \leq N} A_{n+1+k}^{q_{**}^i}$, $q_{**}^i \frown \vec{\nu}$ belongs to D_{μ_i} . Specifically, we choose $q_{**}^i \in M$ to be such a condition which is minimal according to the fixed well-ordering \triangleleft of H_θ , and define

- $N^{D_{\mu_i}}(\vec{\nu}^i) = N$,
- $\vec{f}_\delta = \langle f_{\delta,k} \mid k \leq n \rangle$ with $f_{\delta,k} = f_k^{q_{**}^i} \upharpoonright (\text{dom}(f_k^{q_{**}^i}) \setminus \text{dom}(g_{i,k}))$,⁸ and
- $q^\delta = \langle q_m^\delta \mid m \geq n+1 \rangle$ with $q_m^\delta = (q_{**}^i)_m$ for every $m \geq n+1$.

Finally, given $\vec{f}_{\lambda_n} = \langle f_{\lambda_n,k} \mid k \leq n \rangle$ and $q^{\lambda_n} = \langle q_m^{\delta_n} \mid m \geq n+1 \rangle$. we define $p^{n+1} \geq^* p^n$ by setting $a_m^{p^{n+1}} = a_m^{p^n}$ and $A_m^{p^{n+1}} = A_m^{p^n}$ and $f_m^{p^{n+1}} = f_{\lambda_n,m}$ for $m \leq n$, and $p_m^{n+1} = q_m^{\lambda_n}$ for $m \geq n+1$. Our use of the well-ordering \triangleleft throughout the construction guarantees that $p^{n+1} \in M$.

This concludes the construction of the sequence $\langle p^n \mid \ell^* \leq n < \omega \rangle$. We now define $p^* \geq^* p$ by $p_n^* = p_n^n$. It is straightforward to verify from the construction that $p^* \geq \ell^*$ satisfies the conclusion in the statement of the Lemma. \square

We now show that there are plenty of models M , satisfying the conclusion of Lemma 23.

Proposition 25. *Let $\vec{M} = \langle M_\alpha \mid \alpha < \lambda^+ \rangle$ be an internally approachable sequence,⁹ \subseteq -increasing and continuous sequence of elementary substructures $M_\alpha \prec (H_\theta; \in, \triangleleft)$ with $|M_\alpha| = \lambda$ and $M_\alpha \cap \lambda^+ \in \lambda^+$. For every limit ordinal $\alpha < \lambda^+$ of $\text{cof}(\alpha) = \omega$, satisfying $\alpha = M_\alpha \cap \lambda^+$, $\ell^* < \omega$, and $p \in M_\alpha$, there exists a direct extension $p^* \geq^{\ell^*} p$ satisfying the conclusion of Lemma 23 with respect to $M = M_\alpha$.*

Moreover, if the approachable ideal on λ^+ is trivial, i.e., $I[\lambda^+] = \lambda^+$, then the requirement of $\text{cof}(\alpha) = \omega$ can be removed.

⁸Therefore $\text{dom}(f_{\delta,k})$ is disjoint from $\text{dom}(g_{i,k}) = \text{dom}(a_k^{p^n})$.

⁹i.e., $\vec{M} \upharpoonright \beta \in M_{\beta+1}$ for every $\beta < \lambda^+$.

Proof. Suppose first that $\text{cof}(\alpha) = \omega$ and let $\langle \alpha_n \mid n < \omega \rangle$ be a cofinal sequence in α . Then $M_\alpha = \cup_n M_{\alpha_n}$, and for each $n < \omega$ since $M_{\alpha_n} \in M_\alpha$, there exists an enumeration $\vec{D}^n = \langle D_\mu^n \mid \mu < \lambda \rangle \in M_{\alpha_n}$ of all dense open subsets of \mathbb{P} in M_{α_n} . Using bijections from $\lambda_n \times n$ to λ_n , we can form a sequence $\vec{D} = \langle D_\mu \mid \mu < \lambda \rangle$ so that for every $n < \omega$, $\vec{D} \upharpoonright \lambda_n$ enumerates $\vec{D}^n \upharpoonright \lambda_n$ for each $i < n$. Therefore \vec{D} enumerates all dense open sets of \mathbb{P} in M_α and satisfies $\vec{D} \upharpoonright \beta \in M_\alpha$ for every $\beta < \lambda$. It follows from Lemma 23 that for every condition $p \in M_\alpha$ and $\ell^* < \omega$ there exists a direct extension $p^* \geq^{\ell^*} p$ as in the statement of the lemma. This concludes the first part of the statement.

Suppose now that $I[\lambda^+] = \lambda^+$. We proceed to prove by induction on limit ordinals $\alpha < \lambda^+$ with $\alpha = M_\alpha \cap \lambda^+$, that for every $\ell^* < \omega$ and $p \in M_\alpha$, there is $p^* \geq^{\ell^*} p$ satisfying the desirable property for M_α . Let α be such an ordinal and assume the statment holds for all $\beta < \alpha$. If $\text{cof}(\alpha) = \omega$ we are done by the first case above. Therefore, suppose that $\text{cof}(\alpha) = \rho$ is an uncountable regular cardinal. Since $I[\lambda^+] = \lambda^+$, there exists a closed and unbounded subset $X \subset \alpha$ of order-type $\text{otp}(X) = \rho$ so that $X \cap \beta \in M_{\alpha'}$ whenever $\beta < \alpha'$, $\alpha' \in \{\alpha\} \cup X$. Moreover, since $\vec{M} \upharpoonright \beta$ belongs to $M_{\alpha'}$, so does $\vec{M} \upharpoonright (X \cap \beta) = \langle M_\gamma \mid \gamma \in X \cap \beta \rangle$. Given $\ell^* < \omega$ as in the statement of the claim, we further increase it to assume that $\kappa_{\ell^*} > \rho$. Let $\langle \beta_i \mid i \leq \rho \rangle$ be an increasing enumeration of the limit points β in $X \cup \{\alpha\}$ which satisfy that $M_\beta \cap \lambda^+ = \beta$. Given $p \in M_\alpha$, we may assume that $p \in M_{\beta_0}$ and denote it by p^0 . Then, by applying the inductive assumption and using the well ordering \triangleleft , we form a sequence of conditions $\langle p^i \mid i \leq \rho \rangle$ which is increasing in \leq^{ℓ^*} , so that for each $i < \rho$, $p^i \in M_{\beta_{i+1}}$ and $p^{i+1} \geq^{\ell^*} p^i$ is the \triangleleft -minimal such extension, which satisfies the conclusion of Lemma 23 for $M_{\beta_{i+1}}$. Suppose now that $j \leq \rho$ is limit. Then every initial segment of $\langle p^i \mid i < j \rangle$ belongs to M_{β_j} , and the sequence has an upper bound in \leq^{ℓ^*} since this ordering is κ_{ℓ^*} -closed and $\kappa_{\ell^*} > \rho$. Defining the upper bound by p^j , it follows from the continuity of the sequence \vec{M} that p^j satisfies the desirable property for M_{β_j} . In particular, for $j = \rho$, we obtain a suitable condition $p^* = p^\rho$ for $M = M_\alpha$. \square

The following consequences of Lemma 23 and Proposition 25 will play a key role in our arguments concerning Approachable Bounded Subset Property in $V[G]$.

Lemma 26. *Let \dot{F} be a \mathbb{P} -name of a function from $\lambda^{<\omega}$ to ordinals, and $p \in \mathbb{P}$. There is a direct extension $p^* \geq^* p$ and a function $f^* : [\lambda]^{<\omega} \times [\lambda]^{<\omega} \rightarrow \text{On}$ which provide the following recipe for deciding the \mathbb{P} -names of ordinals $\dot{F}(\vec{\mu})$, $\vec{\mu} \in [\lambda]^{<\omega}$:*

For every $\vec{\mu} \in [\lambda]^{<\omega}$ there are $n^{\vec{\mu}} < \omega$ and a function $N^{\vec{\mu}} : [\lambda]^{<\omega} \rightarrow \omega$ such that for every $\vec{v}^1 \in \prod_{\ell^v \leq i < n^{\vec{\mu}}} A_i^{p^}$ and $\vec{v}^2 \in \prod_{n^{\vec{\mu}} \leq i < N^{\vec{\mu}}(\vec{v}^1)} A_i^{p^*}$,*

$$p^* \frown \vec{v}^1 \frown \vec{v}^2 \Vdash \dot{F}(\vec{\mu}) = f^*(\vec{\mu}, \vec{v}^1 \frown \vec{v}^2).$$

Proof. Let $M \prec (H_\theta; \in, \triangleleft)$ be a model of size which satisfies the assumption of Lemma 23 and has $\dot{F}, p \in M$ (the proof of Proposition 25 shows that such

structures exist). Since $\lambda \subseteq M$ then for every $\vec{\mu} \in [\lambda]^{<\omega}$, the dense open set

$$E_{\vec{\mu}} = \{q \in \mathbb{P} \mid \exists \xi \in \text{On}, q \Vdash \dot{F}(\vec{\mu}) = \check{\xi}\}$$

belongs to M . By taking $p^* \geq^* p$ as in the statement of Lemma 23 we obtain the desired extension of p . \square

Corollary 27. \mathbb{P} preserves λ^+ .

Proof. If $\dot{F} : \lambda \rightarrow \lambda^+$ is a \mathbb{P} -name of a function, then by Lemma 26 for every condition p there are $p^* \geq^* p$ and a function $f^* : [\lambda]^{<\omega} \times [\lambda]^{<\omega} \rightarrow \lambda^+$ in V , so that p^* forces $\text{rng}(\dot{F})$ is contained in $\text{rng}(f^*)$. \square

Let $G \subseteq \mathbb{P}$ be a generic filter. By a standard density argument, for every $\alpha < \lambda^+$ and $n < \omega$ there exists $p \in G$ so that $\ell^p > n$ and $\alpha \in \text{dom}(f_n^p)$. We define the generic scale $\langle t_\alpha \mid \alpha < \lambda^+ \rangle$ by $t_\alpha(n) = f_n^p(\alpha)$ for any such a condition $p \in G$.

Recalling that our setup includes that $a_n(0) = 0$ and $E_n(0)$ is a normal measure on λ_n , we get that the sequence $\langle \rho_n \mid n < \omega \rangle$, given by $\rho_n = t_0(n)$, is generic over V for the diagonal Prikry forcing with the sequence of normal measures $\langle E_n(0) \mid n < \omega \rangle$.

Recall that for every $n < \omega$, there exists a function $g_n : \lambda_n \rightarrow \lambda_n$ so that $j_{E_n}(g_n)(\lambda_n)$ is the supremum of the generators of E_n , and is inaccessible in M_{E_n} . It follows from a standard density argument that the sequence $\langle t_\alpha \mid \alpha < \lambda^+ \rangle$ is a scale in the product $\prod_n g_n(\rho_n)$, and that $g_n(\rho_n) < \lambda_n$ is regular for almost all $n < \omega$. Moreover, it is straightforward to verify that our assumption that the functions a_n in conditions $p \in \mathbb{P}$ are continuous and have closed domains, implies that the scale $\langle t_\alpha \mid \alpha < \lambda^+ \rangle$ is continuous.

Notation 28. In $V[G]$, we denote $g_n(\rho_n)$ by τ_n .

Theorem 29. *The Approachable Bounded Subset Property (ABSP) holds in $V[G]$ with respect to the sequence $\langle \tau_n \mid n < \omega \rangle$.*

Proof. Suppose otherwise, then there exists a stationary set $\mathcal{S} \subseteq \mathcal{P}_\lambda(H_\theta)$ of internally approachable structures $N \prec (H_\theta; \in)$ such that for every $N \in \mathcal{S}$ and $n < \omega$ there is a function $F_n^N : [\lambda]^{k_n^N} \rightarrow \lambda$ in N , of a finite arity $k_n^N < \omega$, and a finite sequence of distinct numbers $\vec{d}^{N,n} = \langle d_0^{N,n}, \dots, d_{k_n^N}^{N,n} \rangle \subseteq \omega \setminus n$, satisfying

$$\chi_N(\tau_{d_0^{N,n}}) \leq F_n^N \left(\chi_N(\tau_{d_1^{N,n}}), \dots, \chi_N(\tau_{d_{k_n^N}^{N,n}}) \right) < \tau_{d_0^{N,n}}.$$

By Lemma 13, applied to the assignments $N \mapsto \langle F_n^N \mid n < \omega \rangle$ and $N \mapsto \langle \vec{d}^{N,n} \mid n < \omega \rangle$, there exists a stationary set $S^* \subseteq \lambda^+$ and two fixed sequences $\langle F_n \mid n < \omega \rangle$, $\langle \vec{d}^n \mid n < \omega \rangle$, with $F_n : [\lambda]^{k_n} \rightarrow \lambda$ and $\vec{d}^n = \langle d_0^n, \dots, d_{k_n}^n \rangle$, such that for every $\delta \in S^*$ there exists $N \in \mathcal{S}$ so that $\delta = \chi_N(\lambda^+)$, $\langle F_n^N \mid n < \omega \rangle = \langle F_n \mid n < \omega \rangle$,

and $\langle \vec{d}^{N,n} \mid n < \omega \rangle = \langle \vec{d}^n \mid n < \omega \rangle$. For each $\delta \in S^*$ there are $m_\delta < \omega$ and $N \in S^*$ such that for every $n \geq m_\delta$, $t_\delta(n) = \chi_N(\tau_n)$ and thus,

$$t_\delta(d_0^n) \leq F_n(t_\delta(d_1^n), \dots, t_\delta(d_{k_n}^n)) < \tau_{d_0^n} \quad (1)$$

We move back to V to contradict the above, and complete the proof. Let p be a condition forcing the statement of (1) with respect to the \mathbb{P} -names \dot{S}^* , $\langle \dot{F}_n \mid n < \omega \rangle$, and $\langle \vec{d}^n \mid n < \omega \rangle$. By taking a direct extension if needed, we may assume p decides the integer values for $\vec{d}^n \subseteq \omega \setminus n$, for all $n < \omega$. Apply Lemma 26 repeatedly for each F_n , $n < \omega$, to form sequences, $\langle p^n \mid n < \omega \rangle$ of \leq^* -extensions of p , and $\langle f^n \mid n < \omega \rangle$ of functions, $f^n : [\lambda]^{<\omega} \times [\lambda]^{<\omega} \rightarrow \lambda$, so that for each $n < \omega$, $p^n \geq^* p^{n-1}$ and f^n are formed to satisfy the conclusion of Lemma 26 with respect to \dot{F}_n . We define

$$\alpha = \sup \left(\bigcup_{n,m} (\text{dom}(a_m^{p^n}) \cup \text{dom}(f_m^{p^n})) \right) + 1$$

and let p^* be a common direct extension of $\langle p^n \mid n < \omega \rangle$ with $\alpha = \max(\text{dom}(a_m^{p^*}))$ for all $m \geq \ell^{p^*}$. Next, let q be an extension of p^* which forces $\check{\delta} \in \dot{S}^*$ for some ordinal $\delta > \alpha$. Since q extends p^* , it is a direct extension of $p^* \frown \vec{v}^*$ for some $\vec{v}^* \in \prod_{\ell^{p^*} \leq i < \ell^*} A_i^{p^*}$. By taking a direct extension of q if needed, we may also assume that q decides the integer values m_δ from above and that $\delta \in \text{dom}(a_n^q)$ for some $n < \omega$.

Next, we pick $n < \omega$ satisfying $n \geq m_\delta$, ℓ^q and $\delta \in \text{dom}(a_n^q)$, and denote for ease of notation, k_n , $\langle d_0^n, \dots, d_{k_n}^n \rangle$ by k , $\langle d_0, \dots, d_k \rangle$ respectively. Our choice of $p^* \geq^* p^n$, and function f^n guarantee that for every $\vec{\mu} = \langle \mu_{d_1}, \dots, \mu_{d_k} \rangle \in [\lambda]^k$ there are $n_*^{\vec{\mu}} < \omega$ and a function

$$N_*^{\vec{\mu}} : \prod_{\ell^q \leq i < n_*^{\vec{\mu}}} A_i^q \rightarrow \omega,^{10}$$

such that for every

$$\vec{v}^1 \in \prod_{\ell^q \leq i < n_*^{\vec{\mu}}} A_i^q \text{ and } \vec{v}^2 \in \prod_{n_*^{\vec{\mu}} \leq i < N_*^{\vec{\mu}}(\vec{v}^1)} A_i^q,$$

denoting $\vec{v}^* \frown \pi_{\text{mc}(q), \alpha}^q(\vec{v}^1 \frown \vec{v}^2)$ by \vec{v} , we have that $p^* \frown \vec{v}$ forces $\dot{F}_n(\vec{\mu}) = \check{f}^n(\vec{\mu}, \vec{v})$.

Recalling that $q \geq^* p^* \frown \vec{v}^*$, we get that in particular, $q \frown \vec{v}^1 \frown \vec{v}^2$, which extends $p^* \frown \vec{v}$ forces the same value, which depends only on $\pi_{\text{mc}(q), \alpha}^q(\vec{v}^1 \frown \vec{v}^2)$.

To complete the argument, we will make use of the last fact, and the fact that $E_n(a_n^q(\delta))$ is strictly stronger than $E_n(a_n^q(\alpha))$ in the Rudin-Keisler ordering, to find many distinct choices of sequences $\vec{v}^1 \frown \vec{v}^2$, whose projections

¹⁰The vales $n_*^{\vec{\mu}}$ and $N_*^{\vec{\mu}}$ are the obvious shifts of $n^{\vec{\mu}}$ and $N^{\vec{\mu}}$ associated to p^n, f^n from Lemma 26, resulting from the fact that the values given by $n^{\vec{\mu}}$ and $N^{\vec{\mu}}$ apply to p^n and hence to p^* , while $q \geq^* p^* \frown \vec{v}^*$, already determined an initial segment \vec{v}^* of a possible end-extension of p^* which decides $\dot{F}_n(\vec{\mu})$ according to the recipe of the lemma.

$\pi_{\text{mc}(q),\delta}^p(\vec{\nu}^1 \frown \vec{\nu}^2)$ are fixed, as well as the values they force for $t_\delta(d_i)$, $1 \leq i \leq k$, yet they force many distinct values for $t_\delta(d_0)$. This will be used to find a condition which extends p but forces (1) to fail.

To this end, we fix first a sequence of $\langle d_1, \dots, d_k \rangle$ -indices,

$$\vec{\nu}_+ = \langle \nu_{d_1}, \dots, \nu_{d_k} \rangle \in \prod_{i \in \{d_1, \dots, d_k\}} A_i^q$$

(note that we omit choosing a d_0 -coordinate) and define $\vec{\mu} = \pi_{\text{mc}(q),\delta}^q(\vec{\nu}_+)$.

Let $j \leq k+1$ largest so that $d_i < n_*^{\vec{\mu}}$ for every $i < j$ and define $\vec{\nu}_+^1 = \langle \nu_{d_0}, \dots, \nu_{d_{j-1}} \rangle$. Therefore, in order to extend $\vec{\nu}_+^1$ to a relevant sequence $\vec{\nu}^1 \in \prod_{\ell^q \leq i < n_*^{\vec{\mu}}} A_i^q$, one needs to choose remaining coordinates

$$\vec{\nu}_-^1 \in \prod_{i \in [\ell^q, n_*^{\vec{\mu}}] \setminus \vec{d}} A_i^q.$$

In particular, to every such sequence $\vec{\nu}_-^1$ we can assign the integer $N_*^{\vec{\mu}}(\vec{\nu}^1)$, where $\vec{\nu}^1 = \vec{\nu}_-^1 \cup \vec{\nu}_+^1$, and by taking a direct extension of q if needed, we may assume that the numbers $N_*^{\vec{\mu}}(\vec{\nu}^1)$ take a constant value N for all $\vec{\nu}_-^1 \in \prod_{i \in [\ell^q, n_*^{\vec{\mu}}] \setminus \vec{d}} A_i^q$. Moreover, by increasing N if necessary, we may also assume that $N > d_k$.

With N being fixed, we conclude that every choice of a sequence

$$\vec{\nu}_-^2 \in \prod_{i \in [n_*^{\vec{\mu}}, N] \setminus \vec{d}} A_i^q$$

will allow us to extend the remaining portion of the fixed sequence $\vec{\nu}_+$ to $\vec{\nu}^2 \in \prod_{i \in [n_*^{\vec{\mu}}, N] \setminus \vec{d}} A_i^q$, which together with a choice of $\vec{\nu}^1$ will produce a suitable extension $q \frown \vec{\nu}^1 \frown \vec{\nu}^2$ forcing

$$\dot{F}_n(\vec{\mu}) = f^n \left(\vec{\mu}, \vec{\nu}^* \frown \pi_{\text{mc}(q),\alpha}^q(\vec{\nu}^1 \frown \vec{\nu}^2) \right).$$

Following this recipe, we extend our fixed choice of $\vec{\nu}_+$ to a choice of all relevant coordinates for $\vec{\nu}^1 \frown \vec{\nu}^2$, except for the coordinate of $i = d_0$. Namely, we extend $\vec{\nu}_+$ to a fixed sequence $\vec{\nu}_{++} \in \prod_{i \in [\ell^q, n_*^{\vec{\mu}} + N] \setminus \{d_0\}} A_i^q$. With the fixed choice $\vec{\nu}_{++}$, we derive a function $h : A_{d_0}^q \rightarrow \lambda_{d_0}$, defined by $h(\nu) = f^n(\vec{\mu}, \pi_{\text{mc}(q),\alpha}^q(\vec{\nu}_{++} \cup \{\nu\}))$ if the last ordinal value is below τ_{d_0} , and $h(\nu) = 0$ otherwise. The properties of the function f^n guarantee that $h(\nu)$ depends only on $\pi_{\text{mc}(q),\alpha}(\nu)$, and by the last clause from Fact 18, there is a subset $A^* \in E_{a_n(d_0)}$, $A^* \subseteq A_{d_0}^q$, so that $h(\nu) < \pi_{\text{mc}(q),\delta}^q(\nu)$ for all $\nu \in A^*$. Picking such an ordinal ν , and setting $\vec{\nu}^1 \frown \vec{\nu}^2 = \vec{\nu}_{++} \cup \{\nu\}$, we conclude that $p^* \frown \vec{\nu}^1 \frown \vec{\nu}^2 \geq p$ must force

$$h(\nu) = \dot{F}_n(t_\delta(d_1), \dots, t_\delta(d_k)) < t_\delta(d_0).$$

Contradicting the statement 1 forced by p . □

Corollary 30. *ABSP holds in $V[G]$ with respect to $\langle \tau_n \mid n < \omega \rangle$ and thus, by Lemma 15, AFSP holds and there are no continuous scales on $\prod_n \tau_n$ which are essentially tree-like.*

2.1 Down to \aleph_ω

We define a variant $\hat{\mathbb{P}}$ of the forcing \mathbb{P} from the previous section, to obtain the result of Theorem 29 in a model where $\langle \tau_n \mid n < \omega \rangle$ form a subsequence of the first uncountable cardinals.

Conditions $q \in \hat{\mathbb{P}}$ are pairs $q = \langle p, h \rangle$ of sequences, $p = \langle p_n \mid n < \omega \rangle$ and $h = \langle h_{-1}^{\text{high}} \rangle \frown \langle h_n \mid n \in \omega \rangle$ satisfying the following conditions:

1. $p \in \mathbb{P}$, i.e., p satisfies Definition 19 above
2. for every $n < \ell^p$, $h_n = \langle h_n^{\text{low}}, h_n^{\text{high}} \rangle$, is a pair of functions, which satisfy the following properties:
 - $h_n^{\text{low}} \in \text{Coll}(\rho_n^p, < \tau_n^p)$ where $\rho_n^p = f_n^p(0)$, and $\tau_n^p = g_n(\rho_n^p)$,¹¹
 - $h_n^{\text{high}} \in \text{Coll}((\tau_n^p)^+, < \rho_{n+1}^p)$ if $n < \ell^p - 1$, and $h_{\ell^p - 1}^{\text{high}} \in \text{Coll}((\tau_{\ell^p - 1}^p)^+, < \lambda_{\ell^p})$.
3. for every $n \geq \ell^p$, $h_n = \langle h_n^{\text{low}}, h_n^{\text{high}} \rangle$, is a pair of functions, which satisfy the following properties:
 - $\text{dom}(h_n^{\text{low}}) = \text{dom}(h_n^{\text{high}}) = A_n^p$,
 - for every $\nu \in A_n^p$,

$$h_n^{\text{low}}(\nu) \in \text{Coll}(\rho_n^\nu, < \tau_n^\nu)$$

where $\rho_n^\nu = \pi_{\text{mc}(a_n^p), 0}^n(\nu)$ and $\tau_n^\nu = g_n(\rho_n^\nu)$, and

$$h_n^{\text{high}}(\nu) \in \text{Coll}((\tau_n^\nu)^+, < \lambda_{n+1})$$

4. h_{-1}^{high} belongs to $\text{Coll}(\omega, < \rho_0^p)$ if $\ell^p \geq 1$, and to $\text{Coll}(\omega, < \lambda_0)$ otherwise
5. $h_{\ell^p - 1}^{\text{high}} \in V_{\rho_{\ell^p}^\nu}$ for every $\nu \in A_{\ell^p}^p$, and $h_{n-1}^{\text{high}}(\nu') \in V_{\rho_n^\nu}$ for every $n > \ell^p$, $\nu' \in A_{n-1}^p$, and $\nu \in A_n^p$.

A condition $q^* = \langle p^*, h^* \rangle$ is a direct extension of $p = \langle p, h \rangle$ if the following conditions hold:

1. $p^* \geq^* p$ in the sense of \mathbb{P} ,
2. for every $n < \ell^p$, $h_n^{\text{low}} \subseteq (h_n^*)^{\text{low}}$ and $h_n^{\text{high}} \subseteq (h_n^*)^{\text{high}}$,
3. for every $n \geq \ell^p$, and $\nu \in A_n^{p^*}$, $h_n^{\text{low}}(\pi_{\text{mc}(a_n^{p^*}), \text{mc}(a_n^p)}^n(\nu)) \subseteq (h_n^*)^{\text{low}}(\nu)$, and $h_n^{\text{high}}(\pi_{\text{mc}(a_n^{p^*}), \text{mc}(a_n^p)}^n(\nu)) \subseteq (h_n^*)^{\text{high}}(\nu)$.

Given a condition $q = \langle p, h \rangle$ and an ordinal $\nu \in A_{\ell^p}^p$, we define the one-point end-extension of q by ν , denoted $q \frown \langle \nu \rangle$ to be the condition $\langle p', h' \rangle$ given as follows:

¹¹By Definition 19 $\rho_n^p < \tau_n^p < \lambda_n$ are both inaccessible for $n \geq 0$.

- $p' = p \widehat{\langle \nu \rangle}$ in the sense of \mathbb{P} , in particular $\ell^{p'} = \ell^p + 1$,
- $h'_n = h_n$ for every $n \leq \ell^p$, and in addition, $(h'_{\ell^p})^{\text{high}}$ is now considered as a condition of the restricted collapse poset $\text{Coll}(\rho_{\ell^p-1}^p, < \tau_{\ell^p}')$ (replacing $\text{Coll}(\rho_{\ell^p-1}^p, < \lambda_{\ell^p})$).
- $(h'_{\ell^p})^{\text{low}} = h_{\ell^p}^{\text{low}}(\nu)$ and $(h'_{\ell^p})^{\text{high}} = h_{\ell^p}^{\text{high}}(\nu)$,
- $h'_n = h_n$ for every $n \geq \ell^p + 1$.

Given a condition $q = \langle p, h \rangle$ and a finite sequence $\vec{\nu} = \langle \nu_{\ell^p}, \dots, \nu_{n-1} \rangle \in \prod_{\ell^p \leq i < n} A_i^p$, the end-extension of q by $\vec{\nu}$ is defined by

$$q \widehat{\vec{\nu}} = q \widehat{\langle \nu_0 \rangle} \widehat{\langle \nu_1 \rangle} \widehat{\dots} \widehat{\langle \nu_{n-1} \rangle}$$

In general, a condition in $\widehat{\mathbb{P}}$ extends q if it is obtain from q by finitely many end-extensions and direct extensions. Equivalently, it is a direct extension of $q \widehat{\vec{\nu}}$ for some $n \geq \ell^p$ and $\vec{\nu} \in \prod_{\ell^p \leq i < n} A_i^p$.

Let $G \subseteq \widehat{\mathbb{P}}$ be a V -generic filter. Through its projection to \mathbb{P} , given by $q = \langle p, h \rangle \mapsto p$, $p \in \mathbb{P}$, it is clear that $V[G]$ adds a sequence of functions $\langle t_\alpha \mid \alpha < \lambda^+ \rangle$ in the product $\prod_n \tau_n$, where $\tau_n = g_n(\rho_n)$ is derived from the generic filter, as in the case of \mathbb{P} . In addition, it is clear that the cardinals in the intervals $(\tau_{n-1}^+, \rho_n) \cup (\rho_n, \tau_n)$, $n < \omega$, are all collapsed in $V[G]$.

A standard argument for diagonal Prikry-type forcings with collapses (e.g., see [12]) shows that no other cardinals are collapsed. It follows that $\tau_n = \aleph_{3n+2}^{V[G]}$ for every $n \in \omega$. Most relevant to us, is Lemma 31 below, which is the $\widehat{\mathbb{P}}$ analog of the Strong Prikry Property from 21.

We first introduce certain useful notations. For a sequence of regular cardinals $\vec{\rho} = \langle \rho_i \mid i < n \rangle$, satisfying $\lambda_{i-1} < \rho_i < g_i(\rho_i) < \lambda_i$, $\mathbb{Q}_{\vec{\rho}}$ to be the product of collapse forcings

$$\text{Coll}(\omega, < \rho_0) \times \left(\prod_{i < n-1} \text{Coll}(\rho_i, < g_i(\rho_i)) \times \text{Coll}(g_i(\rho_i)^+, < \rho_{i+1}) \right) \times \text{Coll}(g_{n-1}(\rho_{n-1})^+, < \lambda_n)$$

where for $i = 0$ we set $g_{-1}(\rho_{-1}) = \omega$. For a condition $q = \langle p, h \rangle \in \widehat{\mathbb{P}}$ we denote $\vec{\rho}^q = \langle f_n^p(0) \mid n < \ell^p \rangle$, $\mathbb{Q}_p = \mathbb{Q}_{\vec{\rho}^p}$, and define for every $q \geq p$, the collapse restriction $q \upharpoonright \mathbb{Q}_p$ to be $h \upharpoonright \ell^p + 1 = \langle h_0, \dots, h_{\ell^p} \rangle$.

Lemma 31. *Suppose that $D \subseteq \widehat{\mathbb{P}}$ is a dense open set and $q = \langle p, h \rangle \in \widehat{\mathbb{P}}$ a condition. Then there exists a direct extension $q^* \geq^* q$, $n < \omega$, such that for every $\vec{\nu} \in \prod_{\ell^p \leq i < n} A_i^{q^*}$, $q^* \widehat{\vec{\nu}}$ reduces meeting D to $\mathbb{Q}_{p^* \widehat{\vec{\nu}}}$, in the sense that there exists a dense open subset $D(\vec{\nu})$ of $\mathbb{Q}_{q^* \widehat{\vec{\nu}}}$ so that for every $q' \geq q$, if $q' \upharpoonright \mathbb{Q}_{q^* \widehat{\vec{\nu}}} \in D(\vec{\nu})$ then $q' \in D$.*

This version of the strong Prikry Property naturally extends to versions of Lemmas 23 and 26, in which in addition to n^D, N^D ($n^{\vec{\mu}}, N^{\vec{\mu}}$, respectively)

which were used to determine the length of sequences $\vec{v}^1 \frown \vec{v}^2$ for $q^* \frown \vec{v}^1 \frown \vec{v}^2$ to meet dense open sets D (or decide values of functions $\dot{F}(\vec{\mu})$), here an additional function \bar{D} mapping sequences $\vec{v}^1 \frown \vec{v}^2$ to dense open subsets $\bar{D}(\vec{v}^1 \frown \vec{v}^2)$ of $\mathbb{Q}_{q^* \frown \vec{v}^1 \frown \vec{v}^2}$, are added, and reduce the problem of finding $q' \geq q^* \frown \vec{v}^1 \frown \vec{v}^2$ inside D , (or deciding $\dot{F}(\vec{\mu})$) to $q' \upharpoonright \mathbb{Q}_{q^* \frown \vec{v}^1 \frown \vec{v}^2}$ being a member of $\bar{D}(\vec{v}^1 \frown \vec{v}^2)$.

We note that in particular, the conclusion of 24 applies to $\hat{\mathbb{P}}$, since $V_\lambda \subseteq M$ implies that the finite collapse products $\mathbb{Q}_{\vec{v}}$ are contained in M .

From this point, it is straightforward to verify that the rest of the argument, leading to an analogous proof of Theorem 29, remains essentially the same, with the additional key being that the identity of the newly introduced collapse products $\mathbb{Q}_{q^* \frown \vec{v}}$ and their dense sets $\bar{D}^{\vec{\mu}}(\vec{v})$ will be decided by the generic information of a bounded part of the scale, t_β , $\beta < \alpha = \sup(M \cap \lambda^+)$ for a suitable structure M of size λ . This information remains independent from higher generic scale functions t_δ , $\delta > \alpha$, which allows one to naturally modify the proof of Theorem 29, to conclude the same result.

Theorem 32. *Let $G \subseteq \hat{\mathbb{P}}$ be a generic filter over V . The Approachable Bounded Subset Property (ABSP) holds in $V[G]$ with respect to the sequence $\langle \aleph_{3n+2} \mid n < \omega \rangle$.*

3 Fine structure and the tree-like scale

3.1 Successor Cardinals

Let $\mathcal{M} \models \text{ZFC}^-$ be a premouse such that every countable hull of \mathcal{M} has an $(\omega_1 + 1)$ iteration strategy, $\lambda \in \mathcal{M}$ a limit cardinal (in \mathcal{M}) of V -cofinality ω (which need not agree with its cofinality in \mathcal{M}) such that \mathcal{M} has a successor cardinal to λ , denoted $(\lambda^+)^{\mathcal{M}}$.

Note if \mathcal{N} is a premouse and $\alpha \in \mathcal{N}$ is such that $\mathcal{N} \models \alpha$ is the largest cardinal, then we let $(\alpha^+)^{\mathcal{N}} = \text{On} \cap \mathcal{N}$.

Let $\vec{\kappa} := \langle \kappa_n : n < \omega \rangle$ be a sequence of \mathcal{M} -cardinals cofinal in λ . We do not assume $\vec{\kappa}$ is in \mathcal{M} . Let $\tau_n := (\kappa_n^+)^{\mathcal{M}}$. We will define a sequence in $\prod_{n < \omega} \tau_n$ that is increasing, tree-like, and continuous.

Let $C^{\lambda, \mathcal{M}} := \{\alpha < (\lambda^+)^{\mathcal{M}} \mid \mathcal{M} \models \alpha < \mathcal{M} \mid (\lambda^+)^{\mathcal{M}}\}$. For $\alpha \in C^{\lambda, \mathcal{M}}$ let \mathcal{M}_α be the collapsing level for α . Let n_α be minimal such that $\rho_{n_\alpha+1}^{\mathcal{M}_\alpha} = \lambda$, $p_\alpha := p_{n_\alpha+1}^{\mathcal{M}_\alpha}$, and $w_\alpha := w_{n_\alpha+1}^{\mathcal{M}_\alpha}$ is the solidity witness described Section 1.2.4. Let also F_α be the top predicate of \mathcal{M}_α .

By Lemma 17 there exists some $\mathcal{M}_\alpha^n \trianglelefteq \mathcal{M}$ such that $\mathcal{C}_0(\mathcal{M}_\alpha^n)$ is isomorphic to $\text{Hull}_{n_\alpha+1}^{\mathcal{M}_\alpha}(\kappa_n \cup \{p_\alpha\})$.

$$f_{\vec{\kappa}, \mathcal{M}}^\alpha(n) = \begin{cases} (\kappa_n^+)^{\mathcal{M}_\alpha^n} & \text{if } \{w_\alpha, \lambda\} \in \text{Hull}_{n_\alpha+1}^{\mathcal{M}_\alpha}(\kappa_n \cup \{p_\alpha\}) \\ 0 & \text{otherwise} \end{cases}$$

Note that the above function is non-zero almost everywhere, that is if $\lambda \in \mathcal{C}_0(\mathcal{M}_\alpha)$. This can fail if (and only if) \mathcal{M}_α is active and $\nu^{\mathcal{M}_\alpha} = \lambda$. For such α we define:

$$f_{\alpha}^{\vec{\kappa}, \mathcal{M}}(n) = \begin{cases} (\kappa_n^+)^{\text{Ult}(\mathcal{M}; F_\alpha \upharpoonright \kappa_n)} & \kappa_n > \mu^{\mathcal{M}_\alpha} \\ 0 & \text{otherwise} \end{cases}$$

By the initial segment condition there must be some $\gamma_\alpha^n < \lambda$ such that the trivial completion of $F_\alpha \upharpoonright \kappa_n$ is indexed at γ_α^n . We let $\mathcal{M}_\alpha^n := \mathcal{M} \upharpoonright \gamma_\alpha^n$. We note that it is impossible that the alternative case of the initial segment condition applies here as κ_n is a cardinal of \mathcal{M} and hence not an index¹²

Note that in either case \mathcal{M}_α^n is the least level of \mathcal{M} over which a surjection from κ_n on to the ordinal $f_{\alpha}^{\vec{\kappa}, \mathcal{M}}(n)$ is definable. Hence the ordinal defines the level and vice versa.

In cases where it is clear which mouse and which sequence of cardinals we are talking about, e.g. for the rest of this subsection, we will omit the superscripts.

Lemma 33. *Let $\alpha < \beta$ both in C . If m is such that $f_\alpha(m) = f_\beta(m)$ then $f_\alpha(n) = f_\beta(n)$ for all $n \leq m$.*

Proof. Note first that if $f_\beta(m) = 0$, then $f_\beta(n) = 0$ and the same holds for α . Let us then consider $f_\beta(m) \neq 0$, it follows that $\mathcal{M}_\alpha^m = \mathcal{M}_\beta^m$. We will start with the assumption that neither α nor β are anomalous. In that situation we must have that $w_\alpha \in \text{Hull}_{n_\alpha+1}^{\mathcal{M}_\alpha^m}(\kappa_n \cup \{p_\alpha\})$. This implies that p_α collapses down to $p_{n_\alpha+1}(\mathcal{M}_\alpha^m)$. The same, of course, holds for β . Note we must have $n_\alpha = n_\beta$. It follows that

$$\begin{aligned} \mathcal{C}_0(\mathcal{M}_\alpha^n) &\cong \text{Hull}_{n_\alpha+1}^{\mathcal{M}_\alpha^m}(\kappa_n \cup \{p_{n_\alpha+1}(\mathcal{M}_\alpha^m)\}) \\ &= \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta^m}(\kappa_n \cup \{p_{n_\beta+1}(\mathcal{M}_\beta^m)\}) \cong \mathcal{C}_0(\mathcal{M}_\beta^n). \end{aligned}$$

This implies $f_\beta(n) = f_\alpha(n)$. Note that $f_\beta(n) = 0$ if and only if $w_{n_\beta+1}(\mathcal{M}_\beta^m) \notin \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta^m}(\kappa_n \cup \{p_{n_\beta+1}(\mathcal{M}_\beta^m)\})$ and similarly for α .

Assume then that at least one of α and β is anomalous. Let us assume that α is anomalous, the proof for β is only notationally different. We will realize that, in fact, both must be anomalous. As types are preserved by taking hulls we must have that both are active type III. As at least one is anomalous we do know that the top extender of \mathcal{M}_α^m has no generators above κ_m . If then the other were not to be anomalous we must have that λ is an element of the appropriate hull. This implies that $\mathcal{C}_0(\mathcal{M}_\beta^m)$ has ordinals and hence generators above κ_n . Contradiction!

As then both are anomalous and $\mathcal{M}_\alpha^m = \mathcal{M}_\beta^m$, we have $F_\alpha \upharpoonright \kappa_m = F_\beta \upharpoonright \kappa_m$. From this follows $\mu^{\mathcal{M}_\alpha} = \mu^{\mathcal{M}_\beta}$ and $F_\alpha \upharpoonright \kappa_n = F_\beta \upharpoonright \kappa_n$. Therefore $f_\alpha(n) = f_\beta(n)$. \square

¹²It also cannot be type Z. Type Z extenders have a largest generator.

Lemma 34. *Let $\alpha < \beta$ both in C . Then $f_\alpha(n) < f_\beta(n)$ for all but finitely many n .*

Proof. Note that since $\alpha < \beta$ are in C then $\mathcal{M}_\alpha \neq \mathcal{M}_\beta$ and so $\mathcal{M}_\alpha \in \mathcal{M}_\beta$. Let us first assume that β is not anomalous.

Let n^* be such that $\mathcal{M}_\alpha \in \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\kappa_{n^*} \cup \{p_\beta\})$. The pre-image of \mathcal{M}_α in \mathcal{M}_β^n ($n \geq n^*$) can then compute \mathcal{M}_α^n and hence $f_\alpha(n)$ correctly.

If on the other hand β were anomalous, let n^* be such that \mathcal{M}_α is generated by some $a \in [\kappa_{n^*}]^{<\omega}$, i.e. $\mathcal{M}_\alpha = \iota_{F_\beta}(h)(a)$ for some $h \in \mu^{\mathcal{M}_\beta}(\mathcal{M} \parallel \mu^{\mathcal{M}_\beta})$. Then $f_\alpha(n)$ ($n \geq n^*$) can be computed from $\iota_{F_\beta \upharpoonright \kappa_n}(h)(a)$ inside $\text{Ult}(\mathcal{M}; F_\beta \upharpoonright \kappa_n)$ by Loś's Theorem. \square

Lemma 35. *Let $\beta \in C$ be of uncountable cofinality. Then β is a continuity point of the sequence (i.e. f_β is the exact upper bound of $\langle f_\alpha : \alpha \in C \cap \beta \rangle$).*

Proof. Let $\alpha_n < f_\beta(n)$. We shall find some $\alpha < \beta$ such that f_α dominates $\langle \alpha_n : n < \omega \rangle$ almost everywhere. Towards that end, we deal first with the case where β is not anomalous.

For almost all $n < \omega$ we have some surjection from κ_n onto α_n in \mathcal{M}_β^n , given by some parameter $a_n \in [\kappa_n]^{<\omega}$ and term τ_n . Let $\xi_n < \rho_{n_\beta}(\mathcal{M}_\beta)$ be such that the uncollapse image of such a surjection is (Σ_1) -definable over $\mathcal{M} \parallel \xi_n$ with $\text{Th}_{n_\beta}^{\mathcal{M}_\beta}(\xi_n, p_{n_\beta}(\mathcal{M}_\beta))$ as an additional predicate.

By Lemma 16, $\rho_{n_\beta}(\mathcal{M}_\beta)$ has uncountable cofinality. So $\xi := \sup_{n < \omega} \xi_n < \rho_{n_\beta}(\mathcal{M}_\beta)$. Take then some A that codes the Σ_1 theory of $\mathcal{M} \parallel \xi$ with $\text{Th}_{n_\beta}^{\mathcal{M}_\beta}(\xi, p_{n_\beta}(\mathcal{M}_\beta))$ as an additional predicate. Such an A exists in \mathcal{M}_β .

Pick some $\alpha < \beta$ such that $A \in \mathcal{M}_\alpha$. Let $n < \omega$ be such that A has a pre-image \bar{A} in \mathcal{M}_α^n . \mathcal{M}_α^n can then compute α_n as the ordertype of

$$\{(\gamma, \delta) \mid (k, a_n \hat{\ } \langle \gamma, \delta \rangle) \in \bar{A}\}$$

where k is the Gödel number of “ $\tau_n(a_n)(\gamma) < \tau_n(a_n)(\delta)$ ”. Hence $\alpha_n < f_\alpha(n)$. Similarly, if α were to be anomalous, we can pick n such that $A = \iota_{F_\alpha}(h)(a)$ for some $h \in \mu^{\mathcal{M}_\beta}(\mathcal{M} \parallel \mu^{\mathcal{M}_\beta})$ and $a \in [\kappa_n]^{<\omega}$. The rest of the argument remains the same.

Let us then assume that β is anomalous. Pick $h_n \in \mu^{\mathcal{M}_\beta}(\mathcal{M} \parallel \mu^{\mathcal{M}_\beta})$ such that $\iota_{F_\beta}(h_n)(a_n)$ is a surjection from κ_n onto α_n for some $a_n \in [\kappa_n]^{<\omega}$. We have that $\text{cof}((\mu^{\mathcal{M}_\beta, +})^\mathcal{M}) > \omega$.

Pick then some $\xi < (\mu^{\mathcal{M}_\beta, +})^\mathcal{M}$ such that $\langle h_n : n < \omega \rangle \subset \mathcal{M} \parallel \xi$. By weak amenability the extender fragment $\bar{F} := \{(a, X) \in F_\beta \mid X \in \mathcal{M} \parallel \xi, a \in [\lambda]^{<\omega}\}$ is in \mathcal{M}_β . Pick then $\alpha < \beta$ with $\bar{F} \in \mathcal{M}_\alpha$. Any \mathcal{M}_α^n containing \bar{F} a pre-image of \bar{F} can then compute α_n as the ordertype of $\{(\gamma, \delta) \mid B_n^{\gamma, \delta} \in \bar{F}\}$ where

$$B_n^{\gamma, \delta} = \{\bar{a} \in [\mu^{\mathcal{M}_\beta}]^{|\bar{b}_n^{\gamma, \delta}|} \mid h_n^{a_n, \bar{b}_n^{\gamma, \delta}}(\bar{a})(\text{id}^{\gamma, \bar{b}_n^{\gamma, \delta}}(\bar{a})) < h_n^{a_n, \bar{b}_n^{\gamma, \delta}}(\bar{a})(\text{id}^{\delta, \bar{b}_n^{\gamma, \delta}}(\bar{a}))\},$$

and $\bar{b}_n^{\gamma, \delta} := a \cup \{\gamma, \delta\}$. Hence $\alpha_n < f_\alpha(n)$. \square

Lemma 36. Assume $\langle \kappa_n : n < \omega \rangle \in \mathcal{M}$, then $\langle f_\alpha : \alpha \in C \rangle$ is a scale in $\prod_{n < \omega} \tau_n \cap \mathcal{M}$.

Proof. Let $f \in \prod_{n < \omega} (\tau_n / J_{bd}) \cap \mathcal{M}$. Pick $\alpha \in C$ such that $f \in \mathcal{M}_\alpha$. Then $f(n) < f_\alpha(n)$ for all but finitely many n . \square

Remark 37. We note that it is possible to associate a sequence in $\prod_{n < \omega} \tau_n$ to any initial segment of \mathcal{M} projecting to λ and it would obey the established rules.

In certain situations we will want to consider a variant construction. Let us consider an additional set of parameters $\vec{\alpha} := \langle \alpha_n : n < \omega \rangle \in \prod_{n < \omega} \tau_n$. Let $\beta \in C$. By the condensation lemma there exists some $\mathcal{M}_\beta^{n, \alpha_n}$ such that $\mathcal{C}_0(\mathcal{M}_\beta^{n, \alpha_n})$ is isomorphic to $\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\kappa_n \cup \{p_\beta \hat{\ } \langle \alpha_n \rangle\})$. We then define:

$$f_\beta^{\vec{\kappa}, \vec{\alpha}, \mathcal{M}}(n) = \begin{cases} (\kappa_n^+)^{\mathcal{M}_\beta^{n, \alpha_n}} & \{\lambda, w_\beta\} \in \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\kappa_n \cup \{p_\beta \hat{\ } \langle \alpha_n \rangle\}) \\ 0 & \text{otherwise} \end{cases}$$

If β is anomalous, then we use $F_\beta \upharpoonright (\alpha_n + 1)$ (instead of $F_\beta \upharpoonright \kappa_n$) to define the sequence.

This sequence will behave just like the previously defined sequence. The proofs are mostly the same. The only minor problem adapting these arguments lie in the preservation of standard parameters. Let p_β^n be the image of p_β under the collapse map in $\mathcal{M}_\beta^{n, \alpha_n}$. Then p_β^n might fail to be the standard parameter of $\mathcal{M}_\beta^{n, \alpha_n}$ as it can fail to be a good parameter.

Though certainly we do know that $p_\beta^n \hat{\ } \langle \alpha_n \rangle$ is a parameter so the standard parameter is below it in the lexicographic order. As we do have a preimage of the solidity witness in $\mathcal{M}_\beta^{n, \alpha_n}$, its standard parameter can only be lesser on that last component, i.e. $p_{n_\beta+1}(\mathcal{M}_\beta^{n, \alpha_n}) = p_\beta^n \hat{\ } \alpha'$ with $\alpha' \leq \alpha_n$.

Then $\mathcal{M}_\beta^{m, \alpha_m}$ can always compute $\mathcal{M}_\beta^{n, \alpha_n}$ from its standard parameter and the ordinal α_n in a consistent matter, guaranteeing tree-likeness of the sequence. Everything else goes through with minor changes.

3.2 Limit cardinals

Let now each of the κ_n be an inaccessible cardinal in \mathcal{M} . We want to extract from $\mathcal{M}_\beta, \beta \in C$, a sequence of structures that singularize some $g_\beta(n) < \kappa_n$. For a singularizing structure the uncollapse embedding ought to move such $g_\beta(n)$ to some regular cardinal in \mathcal{M} . Ideally, we would like this cardinal to be κ_n . Unfortunately, this is unrealistic in general, e.g. if the sequence $\langle \kappa_n : n < \omega \rangle$ is a Prikry sequence for a normal measure on λ . For $\beta \in C$ we define $(\kappa_n^*)_\beta$ to be the minimal ordinal in $\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\kappa_n \cup \{p_\beta\})$ that is $\geq \kappa_n$ if such exists, otherwise $(\kappa_n^*)_\beta = 0$. If β is anomalous, then $(\kappa_n^*)_\beta$ is the least ordinal that can be represented as $\iota_{F_\beta}(f)(a)$ where $f \in \mu^{\mathcal{M}_\beta}(\mu^{\mathcal{M}_\beta})$ and $a \in [\kappa_n]^{<\omega}$ if such exists and zero otherwise.

Note that $(\kappa_n^*)_\beta$ is regular as otherwise it could not be the least ordinal in the hull above κ_n .

In general, it seems likely that the sequences $\langle (\kappa_n^*)_\beta : n < \omega \rangle$ depend on β . For our construction we would like to know that they stabilize (modulo finite) on a tail of C .

Lemma 38. *Let $\alpha < \beta$ both in C . Then $(\kappa_n^*)_\beta \leq (\kappa_n^*)_\alpha$ for all but finitely many $n < \omega$.*

Proof. We have $\mathcal{M}_\alpha \in \mathcal{M}_\beta$. Therefore for some $n^* < \omega$ we have $\mathcal{M}_\alpha \in \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\kappa_n^* \cup \{p_\beta\})$. Hence $(\kappa_m^*)_\alpha \in \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\kappa_m \cup \{p_\beta\})$ for $m \geq n^*$. The anomalous case is similar. \square

If we knew that $\text{cof}((\lambda^+)^\mathcal{M}) > 2^{\aleph_0}$, then we could use the Erdős-Rado Theorem to show that the sequences stabilize. While this should be a reasonable assumption in most use cases, we will present a different result here that uses merely $\text{cof}((\lambda^+)^\mathcal{M}) > \aleph_1$.

Lemma 39. *Let $\beta \in \lim(C)$ such that $\text{cof}(\beta) > \aleph_0$. Then there exists some $\alpha < \beta$ such that $(\kappa_n^*)_\alpha = (\kappa_n^*)_\beta$ for all but finitely many n .*

Proof. Let $\xi_n < \rho_{n_\beta}(\mathcal{M}_\beta)$ be such that $(\kappa_n^*)_\beta$ is (Σ_1) -definable over $(\mathcal{M} \upharpoonright \xi_n \text{Th}_{n_\beta}^{\mathcal{M}_\beta}(\xi_n, p_{n_\beta}(\mathcal{M}_\beta)))$.

As in the proof of Lemma 35, we can find some $A \subset \lambda$ coding the Σ_1 theory of $(\mathcal{M} \upharpoonright \xi_n; \text{Th}_{n_\beta}^{\mathcal{M}_\beta}(\xi_n, p_{n_\beta}(\mathcal{M}_\beta)))$ with parameters from λ in \mathcal{M}_β .

Any $\alpha \in C$ with $A \in \mathcal{M}_\alpha$ will then be as wanted. The anomalous case is similar. \square

Note that the two preceding lemmas taken together imply that $(\kappa_n^*)_\gamma = (\kappa_n^*)_\beta$ for all but finitely many $n < \omega$ and all $\alpha < \gamma < \beta$. The next lemma is then immediate.

Lemma 40. *If $\text{cof}((\lambda^+)^\mathcal{M}) > \aleph_1$, then there exists some β^{**} such that $(\kappa_n^*)_\alpha = (\kappa_n^*)_\beta$ for all but finitely many $n < \omega$ for all $\beta^{**} < \alpha < \beta$.*

To finally define the scale we now need an additional vector of parameters $\vec{\alpha} = \langle \alpha_n : n < \omega \rangle$ where $\alpha_n < \kappa_n$. We also do require that $\sup_{n < \omega} \alpha_n = \lambda$. When do these parameters give rise to the right structure? This will depend on whether β is anomalous or not. We begin with listing three key factors for the case β is not anomalous:

$$(1)_n^\beta \sup(\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\alpha_n \cup \{p_\beta\}) \cap \kappa_n) > \alpha_n;$$

$$(2)_n^\beta (\kappa_n^*)_\beta \in \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\alpha_n \cup \{p_\beta\})$$

$$(3)_n^\beta \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\alpha_n \cup \{p_\beta\}) \text{ is cofinal in } \rho_{n_\beta}(\mathcal{M}_\beta).$$

If β is anomalous, we have the following two considerations:

$$(4)_n^\beta \sup(\{\iota_{F_\beta}(h)(a) \mid h \in \mu^{\mathcal{M}_\beta}(\mu^{\mathcal{M}_\beta}), a \in [\alpha_n]^{<\omega}\} \cap \kappa_n) > \alpha_n;$$

(5) $_n^\beta$ $(\kappa_n^*)_\beta = \iota_{F_\beta}(h)(a)$ for some $h \in \mu^{\mathcal{M}_\beta}(\mu^{\mathcal{M}_\beta})$ and $a \in [\alpha_n]^{<\omega}$.

We say β is adequate iff $\beta > \beta^{**}$,¹³ and (1) $_n^\beta + (2)_n^\beta + (3)_n^\beta$ or (4) $_n^\beta + (5)_n^\beta$ (depending on type) are met for all but finitely many n . If β is adequate and not anomalous then

$$g_\beta^{\vec{\kappa}, \vec{\alpha}, \mathcal{M}}(n) := \begin{cases} \sup(\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\alpha_n \cup \{p_\beta\}) \cap \kappa_n) & (1)_m^\beta + (2)_m^\beta + (3)_m^\beta \forall m \geq n \\ 0 & \text{otherwise} \end{cases}$$

We then let \mathcal{M}_β^n be the unique initial segment of \mathcal{M} such that $\mathcal{C}_0(\mathcal{M}_\beta^n)$ is isomorphic to $\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(g_\beta(n) \cup \{p_\beta\})$. Note that by the the second case in the condensation lemma cannot hold as by the proof of the next lemma $g_\beta(n)$ is the critical point of the natural map from \mathcal{M}_β^n to \mathcal{M}_β and since $g_\beta(n)$ is a limit of cardinals and hence a cardinal itself. This follows by elementarity, trivially so when $n_\beta > 0$ otherwise by (3) $_n^\beta$.)

If on the other hand β is anomalous then

$$g_\beta^{\vec{\kappa}, \vec{\alpha}, \mathcal{M}}(n) := \begin{cases} \sup(\{\iota_{F_\beta}(h)(a) \mid h \in \mu^{\mathcal{M}_\beta}, a \in [\alpha_n]^{<\omega}\} \cap \kappa_n) & (4)_m^\beta + (5)_m^\beta \forall m \geq n \\ 0 & \text{otherwise} \end{cases}$$

\mathcal{M}_β^n will be the unique initial segment of \mathcal{M} with the trivial completion of $F_\beta \upharpoonright g_\beta(n)$ as its top extender. As in the previous section, we will omit superscripts for the remainder of this section.

To ensure tree-likeness for this sequence we need a strong interdependence between the ordinal $g_\beta(n)$ and structure \mathcal{M}_β^n . Towards that end notice that $g_\beta(n)$ is definably singularized over \mathcal{M}_β^n . The next lemma will show that \mathcal{M}_β^n is the least level of \mathcal{M} with this property.

Lemma 41. *$g_\beta(n)$ is regular in \mathcal{M}_β^n for all n such that (1) $_n^\beta + (2)_n^\beta + (3)_n^\beta$ or (4) $_n^\beta + (5)_n^\beta$ holds.*

Proof. First we will consider β that is not anomalous. Since $(\kappa_n^*)_\beta$ is regular in \mathcal{M}_β , it will then be enough to show that $\sup(\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(g_\beta(n) \cup \{p_\beta\}) \cap \kappa_n) = g_\beta(n)$.

Let $\xi < \kappa_n$ be such that $\xi \in \text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(g_\beta(n) \cup \{p_\beta\})$. We can then take $\gamma < g_\beta(n)$ and $\delta < \rho_{n_\beta}(\mathcal{M}_\beta)$ such that $\xi \in \text{Hull}_1^{\mathcal{M}_\beta^\delta}(\gamma)$ where \mathcal{N}_δ is $\mathcal{M} \upharpoonright \delta$ together with $\text{Th}_{n_\beta}^{\mathcal{M}_\beta}(\delta, p_{n_\beta}(\mathcal{M}_\beta))$ as an additional predicate. We can take γ and δ to be in $\text{Hull}_{n_\beta+1}^{\mathcal{M}_\beta}(\alpha_n \cup \{p_\beta\})$ (by definition of $g_\beta(n)$ and (3) $_n$ respectively).

Then $\eta := \sup(\text{Hull}_1^{\mathcal{M}_\beta^\delta}(\gamma) \cap \kappa_n)$ is also in that hull (uses (2) $_n$) and thus $\xi < \eta < g_\beta(n)$.

Now consider an anomalous β . We will show that $g_\beta(n)$ is regular in $\text{Ult}(\mathcal{M}; F_\beta \upharpoonright g_\beta(n))$. We have some $h \in \mu^{\mathcal{M}_\beta}_{\mu^{\mathcal{M}_\beta}}$ and $a \in [\alpha_n]^{<\omega}$ such that $(\kappa_n^*)_\beta = \iota_{F_\beta}(h)(a)$. We will show that $g_\beta(n) = \iota_{F_\beta \upharpoonright g_\beta(n)}(h)(a)$. As this pair represents a regular cardinal in the larger ultrapower this will suffice.

¹³ β^{**} from Lemma 40 above.

Pick then some h_0 and b such that $b \in [g_\beta(n)]^{<\omega}$ and $\iota_{F_\alpha \upharpoonright g_\beta(n)}(h_0)(b) < \iota_{F_\beta \upharpoonright g_\beta(n)}(h)(a)$. Pick some $c \in [\alpha_n]^{<\omega}$ (w.l.o.g. $a \subset c$) and h_1 such that $b \subset \iota_{F_\beta}(h_1)(c)$. Define $h_2 : [\mu^{\mathcal{M}_\beta}]^{|c|} \rightarrow \mu^{\mathcal{M}_\beta}$ by $d \mapsto \sup\{h_0(e) \mid e \in [h_1(d)]^{|b|}, h_0(e) < h^{a,c}(d)\}$.

We then have $\iota_{F_\beta \upharpoonright g_\beta(n)}(h_0)(b) \leq \iota_{F_\beta \upharpoonright g_\beta(n)}(h_2)(c) < g_\beta(n)$ as required. \square

Thus $g_\beta(n)$ is regular in \mathcal{M}_β^n but is definably singular over it. Thus it is uniquely determined as a level of \mathcal{M} by $g_\beta(n)$.

The following is a straightforward corollary of the proof of the previous Lemma.

Corollary 42. *Let $\alpha \in C$ be such that $g_\beta(n)$ is defined for all but finitely many n . Let $\vec{\alpha}^* := \langle \alpha_n^* : n < \omega \rangle$ be such that $\alpha_n \leq \alpha_n^* < g_\alpha(n)$ for all but finitely many n . Then $g_\beta^{\vec{\kappa}, \mathcal{M}, \vec{\alpha}^*}$ is defined and agrees with g_β almost everywhere.*

Lemma 43. *Say β^* is adequate, then every $\beta > \beta^*$ in C of uncountable cofinality is also adequate.*

Proof. Let us assume for simplicity's sake that $(1)_n^{\beta^*} + (2)_n^{\beta^*} + (3)_n^{\beta^*}$ holds for all n . Let then n^* be such that $\beta^* \in \text{Hull}_{n, \beta^*+1}^{\mathcal{M}_{\beta^*}}(\alpha_m \cup \{p_\beta\})$ and $(\kappa_m^*)_{\beta^*} = (\kappa_m^*)_\beta$ for all $m \geq n^*$. Then that hull can compute $\text{Hull}_{n, \beta^*+1}^{\mathcal{M}_{\beta^*}}(\alpha_m \cup \{p_{\beta^*}\})$ for all $m \geq n^*$. $(1)_m^\beta + (2)_m^\beta$ then follows.

That is if β^* is not anomalous. If it is anomalous note that $\text{Hull}_{n, \beta^*+1}^{\mathcal{M}_{\beta^*}}(\alpha_m \cup \{p_\beta\})$ has access to the extender F_{β^*} and can compute $(\kappa_m^*)_\beta$ from it assuming $\alpha_m > \mu^{\mathcal{M}_{\beta^*}}$. $(1)_m^\beta$ follows for similar reasons.

$(3)_m^\beta$ almost everywhere follows for cofinality reasons.

If β is anomalous then we take some h and $a \in [\alpha_m]^{<\omega}$ such that $\beta^* = \iota_{F_\beta}(h)(a)$ and let τ be some $r\Sigma_{n, \beta^*+1}$ -term such that $(\kappa_m^*)_{\beta^*} = \tau_m^{\mathcal{M}_{\beta^*}}(b_m, p_{\beta^*})$ for $b_m \in [\alpha_m]^{<\omega}$. Define then $h_0^m : [\mu^{\mathcal{M}_\beta}]^{|a \cup b_m|} \rightarrow \mu^{\mathcal{M}_\beta}$ by

$$c \mapsto \tau_m^{\mathcal{M}_{h^a, a \cup b_m(c)}}(\text{id}^{b_m, a \cup b_m}(c), p_{h^a, a \cup b_m(c)}).$$

We then have $\iota_{F_\beta}(h_0^m)(a \cup b_m) = (\kappa_m^*)_{\beta^*}$. $(4)_m^\beta$ follows for similar reasons.

The idea is similar if β^* and β are both anomalous. (Pick h, a representing F_{β^*} etc.) We skip further details. \square

Assuming the existence of an adequate ordinal β^* we can then show that $\langle g_\beta : \beta \in C \setminus \beta^* \cap \text{cof}(> \omega) \rangle$ is increasing (mod finite), tree-like, and continuous as before.

Lemma 44. *Let $\langle \kappa_n : n < \omega \rangle$ and $\langle \alpha_n : n < \omega \rangle$ both in \mathcal{M} then there exists an adequate β^* and $\langle g_\beta : \beta \in C \setminus \beta^* \cap \text{cof}(> \omega) \rangle$ is a scale in $\prod_{n < \omega} \kappa_n \cap \mathcal{M}$.*

Proof. Any β^* of uncountable cofinality (in \mathcal{M}) such that $\mathcal{M} \parallel \beta^*$ contains both sequences will be adequate. The rest is then as before. \square

Remark 45. Note that while we have only considered sequences $\langle \kappa_n \mid n < \omega \rangle$ of uniform type (i.e., all successors or all inaccessible) we could easily deal with sequences $\langle \kappa_n : n < \omega \rangle$ of regular cardinals with both successor cardinals and inaccessible cardinals by mixing both constructions using parameters where needed. Usually, such a procedure would jeopardize tree-likeness, but our case is special. Tree-likeness is certified by the fact that the level \mathcal{M}_α^n can compute the structures \mathcal{M}_α^m for $m < n$. For this the type of structure is irrelevant, a collapsing structure can compute singularizing structures and vice versa. For example if $\kappa_0 < \kappa_1 < \lambda$ with κ_1 inaccessible and M_α^1 is the singularizing structure for $g_\alpha(\kappa_1)$ isomorphic to $\text{Hull}_{n_{\alpha+1}}^{\mathcal{M}_\alpha}(g_\alpha(\kappa_1) \cup \{p_\alpha\})$ then $\text{Hull}_{n_{\alpha+1}}^{\mathcal{M}_\alpha}(\kappa_0 \cup \{p_\alpha\})$ is isomorphic to $\text{Hull}_{n_{\alpha+1}}^{\mathcal{M}_\alpha^1}(\kappa_0 \cup \{p_\alpha\})$.

Remark 46. Assuming that λ is not subcompact in \mathcal{M} the sequences we defined should be very good, but we have yet to check this in detail. The proof would presumably proceed along similar lines as in [7].

4 Core models and the tree-like scale

We now want to consider the question when the sequences constructed in the previous section are scales in V . For this we need to consider the right mouse. The natural candidate is, of course, the core model. But even core model sequences are not always scales.

To keep the following as accessible as possible we are going to operate under a smallness assumption. This will allow us to cover all known anti tree-like scale results while greatly simplify the following arguments.

This assumption is:

$$\begin{aligned} \text{There is no inner model } W \text{ and } F \in W \text{ a total extender} \\ \text{such that } \text{gen}(F) \geq (\text{crit}(F)^{++})^W \end{aligned} \quad (2)$$

Corollary 47. *There is no ω_1 -iterable premouse $(\mathcal{M}, \in, \vec{E}, F)$ such that $\text{gen}(F) > (\text{crit}(F)^{++})^{\mathcal{M}}$.*

Proof. Assume towards a contradiction that $(\mathcal{M}; \in, \vec{E}, F)$ is a counterexample. Then we can generate an inner model W by iterating the top extender out of the universe. Note that by a standard reflection argument, ω_1 -iterability is enough to ensure that this model is wellfounded. By the initial segment condition $F \upharpoonright (\text{crit}(F)^{++})^W$ is then in W contradicting (2). \square

The reader should be aware, though, that our main results will hold under much weaker anti-Large Cardinals assumptions (up to one Woodin cardinal and beyond). Neither should the choice of indexing scheme affect their validity (though we have yet to check this in detail).

The most immediate payoff of (2) will be that all iterations we are going to consider are linear (This is one instance in which ms-indexing will make things simpler for us).

Proposition 48. *Let \mathcal{M} be a ω_1 -iterable premouse, and \mathcal{T} be a normal iteration tree on \mathcal{M} . Then no $\alpha < \beta \leq \text{lh}(\mathcal{T})$ is such that $\text{crit}(E_\beta^\mathcal{T}) < \text{gen}(E_\alpha^\mathcal{T})$.*

Proof. Let $\alpha < \beta$ such that $\text{crit}(E_\beta^\mathcal{T}) < \text{gen}(E_\alpha^\mathcal{T})$. There are three cases:

Case 1: $\text{crit}(E_\alpha^\mathcal{T}) < \text{crit}(E_\beta^\mathcal{T})$

By agreement between models in an iteration we have that $\text{crit}(E_\beta^\mathcal{T})$ is inaccessible in $\mathcal{M}_\alpha^\mathcal{T} \parallel \text{lh}(E_\alpha^\mathcal{T})$ and thus $(\text{crit}(E_\alpha^\mathcal{T})^{++})^{\mathcal{M}_\alpha^\mathcal{T} \parallel \text{lh}(E_\alpha^\mathcal{T})} < \text{crit}(E_\beta^\mathcal{T})$. As $E_\alpha^\mathcal{T}$ has generators above $\text{crit}(E_\beta^\mathcal{T})$, $\mathcal{M}_\alpha^\mathcal{T} \parallel \text{lh}(E_\alpha^\mathcal{T})$ is a counterexample to Corollary 47.

Case 2: $\text{crit}(E_\alpha^\mathcal{T}) > \text{crit}(E_\beta^\mathcal{T})$

In $\mathcal{M}_\beta^\mathcal{T}$ due to the agreement between models in an iteration, $\text{lh}(E_\alpha^\mathcal{T}) > \text{crit}(E_\beta^\mathcal{T})$ is a cardinal in $\mathcal{M}_\beta^\mathcal{T}$ so by strong acceptability $\text{crit}(E_\alpha^\mathcal{T})$ is inaccessible in $\mathcal{M}_\beta^\mathcal{T}$ and thus above $(\text{crit}(E_\beta^\mathcal{T})^{++})^{\mathcal{M}_\beta^\mathcal{T}}$. As \mathcal{T} is a normal iteration $\text{lh}(E_\beta^\mathcal{T}) > \text{lh}(E_\alpha^\mathcal{T}) > (\text{crit}(E_\alpha^\mathcal{T})^+)^{\mathcal{M}_\beta^\mathcal{T}}$ and so $\text{gen}(E_\beta^\mathcal{T}) > \text{crit}(E_\alpha^\mathcal{T})$ but then $\mathcal{M}_\beta^\mathcal{T} \parallel \text{lh}(E_\beta^\mathcal{T})$ is a counterexample to Corollary 47.

Case 3: $\text{crit}(E_\alpha^\mathcal{T}) = \text{crit}(E_\beta^\mathcal{T})$

Because \mathcal{T} is normal we have $\text{lh}(E_\alpha^\mathcal{T}) < \text{lh}(E_\beta^\mathcal{T})$ but this means that $E_\beta^\mathcal{T}$ must have generators cofinal in $(\text{crit}(E_\beta^\mathcal{T})^{++})^{\mathcal{M}_\beta^\mathcal{T}}$. Now, let γ be the last drop in the interval $(\alpha, \beta]$ if it exists or $\alpha + 1$ otherwise. We can assume that $\text{crit}(E_{\gamma-1}^\mathcal{T}) \geq \text{crit}(E_\beta^\mathcal{T})$. $\iota_{\gamma-1, \beta}^\mathcal{T}(\text{crit}(E_{\gamma-1}^\mathcal{T}))$ is then the critical point of an extender on the $\mathcal{M}_\beta^\mathcal{T}$ sequence and greater than $\text{crit}(E_\beta^\mathcal{T})$. As $E_\beta^\mathcal{T}$ must be total over $\mathcal{M}_\beta^\mathcal{T}$. Thus we can produce a class size model W containing $E_\beta^\mathcal{T}$ and agreeing with $\mathcal{M}_\beta^\mathcal{T}$ past $(\text{crit}(E_\beta^\mathcal{T})^{++})^W$ which contradicts (2). □

Another consequence of (2) is that the Jensen-Steel core model K exists by [17]. Note that by the smallness assumption there can be no anomalous ordinals in K . For the following results we will follow the general framework of the proof of weak covering for that model. Before going into the proofs we shall take quick note of the involved objects.

Let λ be a singular cardinal of countable cofinality. Let $\vec{\kappa} := \langle \kappa_n : n < \omega \rangle$ be a sequence cofinal in λ . Let $\tau_n := (\kappa_n^+)^K$. Consider some $X \prec H_\theta$ ($\theta \gg \lambda$) and let $\sigma^X : H_X \rightarrow X$ be the inverse of the transitive collapse map. X will need to satisfy certain properties:

- certain phalanxes “lift” through σ^X
- $\text{card}(X) < \lambda$,
- X is tight on $\vec{\kappa}$ (and $\langle \tau_n : n < \omega \rangle$), i.e. $X \cap \prod_{n < \omega} \kappa_n$ is cofinal in $\prod_{n < \omega} (X \cap \kappa_n) / J_{bd}$,
- the collection of $X \prec H_\theta$ with the above three properties is stationary.

The first point is quite vague, and we will provide more details where needed in the course of the argument. By [20] ω -closed X do satisfy the first property, but it seems possible that there are not enough, i.e. stationary many, X with all properties available. In such cases, by [21] we do know that for every internally approachable chain $\vec{Y} := \langle Y_i : i < \kappa \rangle$ in H_θ there exists some $i < \kappa$ of uncountable cofinality such that Y_i satisfies the first property. That it satisfies the other properties should be easy to see.

Let then from now on X be some such set with the required properties. Let $\sigma_X : H_X \rightarrow X$ be an isomorphism where H_X is transitive. Write $K_X := \sigma_X^{-1} \upharpoonright [K]$, $\lambda_X := \sigma_X^{-1}(\lambda)$, $\kappa_n^X := \sigma_X^{-1}(\kappa_n)$, etc.

As is standard we will compare K_X with K , we should have (for our choice of X) that the iteration tree on K_X is trivial (this is $(1)_\alpha$ from [20] or $(1)_\alpha^i$ from [21] respectively). Let then \mathcal{I}_X be the iteration tree on K that arises from the co-iteration. We will simplify notation by writing \mathcal{M}_α^X for $\mathcal{M}_\alpha^{\mathcal{I}_X}$ etc. Let $\zeta_X := \text{lh}(\mathcal{I}_X)$ be the length of the iteration.

Lemma 49. $(\text{crit}(\sigma_X)^+)^{K_X} < (\text{crit}(\sigma_X)^+)^K$ and if E_0^X is defined then it is not total over K .

Note that K_X and K agree up to $(\text{crit}(\sigma_X)^+)^{K_X}$ as a result of the condensation lemma.

Proof. Assume towards a contradiction that $(\text{crit}(\sigma_X)^+)^{K_X} = (\text{crit}(\sigma_X)^+)^K$. Then E_{σ_X} the $(\text{crit}(\sigma_X), \sigma_X(\text{crit}(\sigma_X)))$ -extender derived from σ_X measures all subsets of its critical point that are in K . It also coheres with K by the elementarity of σ_X . (This is a little bit of a lie. We would actually need to know that all Mitchell-Steel initial segments of E_{σ_X} are on the K -sequence. But if this fails we could simply apply the argument we are about to give to the least missing segment instead.)

We do know that the phalanx $\langle \langle K, \text{Ult}(K; E_{\sigma_X}) \rangle, \sigma_X(\text{crit}(\sigma_X)) \rangle$ is iterable. This is $(2)_\alpha$ from [20] or [21] where $\text{crit}(\sigma_X) = (\aleph_\alpha)^{K_X}$. (Once again this is something of a lie. We actually have to replace K with an appropriate soundness witness in the above statement, but we can choose W such that it agrees with K past the level we actually care about. Thus this will not make a difference here.)

But then by [27, 8.6] we have that E_{σ_X} is on the K -sequence. It should be obvious that $\text{gen}(E_{\sigma_X}) = \sigma_X(\text{crit}(\sigma_X))$ and thus $K \upharpoonright \text{lh}(E_{\sigma_X})$ contradicts Corollary 47.

As for the second part, assume E_0^X is applied to K . By the first part, if $\text{crit}(E_0^X) \geq \text{crit}(\sigma_X)$, then we must truncate. If $(\text{crit}(E_0^X)^+)^{K_X} = \text{crit}(\sigma_X)$, then by elementarity $(\text{crit}(E_0^X)^+)^K = \sigma_X(\text{crit}(\sigma_X))$ so we must truncate.

If $\text{crit}(\sigma_X) \geq (\text{crit}(E_0^X)^{++})^{K_X}$ then its generators must be cofinal in $\text{crit}(\sigma_X)$. So, if the strict inequality holds then E_0^X contradicts Corollary 47. A similar argument applies if $\text{lh}(E_0^X) > (\text{crit}(\sigma_X)^+)^{K_X}$.

So, we must have that $\text{crit}(\sigma_X) = (\text{crit}(E_0^X)^{++})^{K_X}$. Consider \mathcal{M}_1^X . It agrees with K_X up to $(\text{crit}(\sigma_X)^+)^{K_X}$ and that ordinal is a cardinal there. Thus we

can apply the extender E_{σ_X} to it. The properties of X will guarantee that $\tilde{K} := \text{Ult}(\mathcal{M}_1^X; E_{\sigma_X})$ is iterable (similar to the proof of [20, Lemma 3.13]).

We have that K and \tilde{K} agree up to $\text{sup}(\sigma_X)'' [(\text{crit}(\sigma_X)^+)^{K_X}]$ which lies past $\sigma_X(\text{crit}(\sigma_X))$, but on the other hand

$$(\text{crit}(E_0^X)^{+++})^{\tilde{K}} = \text{sup}(\sigma_X)'' [(\text{crit}(\sigma_X)^+)^{K_X}] < \sigma_X((\text{crit}(\sigma_X)^+)^{K_X}) = (\text{crit}(E_0^X)^{+++})^K$$

as a result of weak covering. Consider then \tilde{E} the first extender applied on the K side in the co-iteration of K and \tilde{K} . Its index must be above $\sigma_X(\text{crit}(\sigma_X))$ but its critical point cannot be larger than $\text{crit}(E_0^X)$. \tilde{E} on the K -sequence then contradicts (2). □

Remember now the sequence $\langle \kappa_n : n < \omega \rangle$ and the sequence of successors $\langle \tau_n : n < \omega \rangle$. The general idea for the following proofs is to find some ordinal $\alpha_X < \lambda^+$ such that the natural scales of the core model at α_X align with the characteristic function of X .

From now on we shall assume that κ_n is a cutpoint of (the extender sequence of) K and hence κ_n^X is a cutpoint of K_X . ($\alpha \in (\mathcal{M}; \in, \vec{E})$ is a cutpoint (of \vec{E}) iff whenever $\text{crit}(E_\beta) < \alpha$, then $\text{lh}(E_\alpha) < \alpha$ for all $\beta \in \text{dom}(\vec{E})$.)

Lemma 50. *There exist some $n_X, k_X < \omega$, a sequence of models $\langle \mathcal{N}_n^X : n_X \leq n < \omega \rangle$, and maps $\langle v_{n,m}^X : n_X \leq n \leq m < \omega \rangle$ such that:*

- $((\kappa_n^X)^+)^{\mathcal{N}_n^X} = \tau_n^X$, and \mathcal{N}_n^X agrees with K_X up to τ_n^X for all $n \geq n_X$;
- \mathcal{N}_n^X is $(k_X + 1)$ -sound above κ_n^X for all $n \geq n_X$;
- $v_{n,m}^X : \mathcal{C}_0(\mathcal{N}_n^X) \rightarrow \mathcal{C}_0(\mathcal{N}_m^X)$ is $r\Sigma_{k_X+1}$ -elementary for all $m \geq n \geq n_X$;
- $\text{crit}(v_{n,m}^X) \geq \kappa_n^X$ for all $m \geq n \geq n_X$.

For our purposes the critical point of the identity will be defined as the ordinals of its domain.

Proof. There are a couple of cases.

Case 1: \mathcal{I}_X has no indices below λ .

In that case, we have $K_X | (\lambda^+)^{K_X} \trianglelefteq K$. By Lemma 49 we do know that some $\mathcal{N}' \trianglelefteq K$ exists with $(\text{crit}(\sigma_X)^+)^{\mathcal{N}'} = (\text{crit}(\sigma_X)^+)^{K_X}$ and $\rho_\omega(\mathcal{N}') \leq \text{crit}(\sigma_X)$. By assumption we must have $K_X | (\lambda^+)^{K_X} \trianglelefteq \mathcal{N}'$.

Take then \mathcal{N}^* to be the smallest initial segment of K that end-extends $K_X | (\lambda^+)^{K_X}$ such that $\rho_\omega(\mathcal{N}^*) < \lambda_X$.

Let k_X be minimal such that $\rho_{k_X+1}(\mathcal{N}^*) < \lambda_X$. Let n_X be minimal such that $\kappa_{n_X}^X \geq \rho_{k_X+1}(\mathcal{N}^*)$. We then let $\mathcal{N}_n^X := \mathcal{N}^*$ for all $n \geq n_X$, and let $v_{n,m}^X$ be the identity for all $m \geq n \geq n_X$. As an initial segment of K , \mathcal{N}^* is sound so this works.

Case 2: The set $\{\text{lh}(E_\beta^X) \mid \beta < \zeta_X\}$ is bounded below λ_X .

Let $\eta_X < \zeta_X$ be minimal such that $E_{\eta_X}^X$ has length $> \lambda_X$, if it exists. If there is no such ordinal, let $\eta_X = \zeta_X$. We must then have that $\mathcal{M}_{\eta_X}^X$ agrees with K_X past λ_X . If $\mathcal{M}_{\eta_X}^X$ has some proper initial segment of length greater than λ_X projecting below λ_X , then this is no different from the previous case.

So let us assume that this is not the case. Let n_X be minimal such that $\kappa_{n_X}^X$ is above the set $\{\text{lh}(E_\beta^X) \mid \beta < \eta_X\}$. By Lemma 49, $\mathcal{M}_{\eta_X}^X$ is not a weasel and is $(k_X + 1)$ -sound above $\kappa_{n_X}^X$ for some unique k_X .

We then let $\mathcal{N}_n^X := \mathcal{M}_{\eta_X}^X$ for all $n \geq n_X$, and $v_{n,m}^X$ the identity for all $m \geq n \geq n_X$.

Case 3: The set $\{\text{lh}(E_\beta^X) \mid \beta < \zeta_X\}$ is cofinal below λ_X .

Let $\eta_X := \sup(\{\beta < \zeta_X \mid \text{lh}(E_\beta^X) < \lambda_X\})$. By assumption and Lemma 49 there is some drop in the interval $(0, \eta_X)$. Let then $\gamma + 1$ be the last such.

Let k_X be minimal such that $\rho_{k_X+1}((\mathcal{M}_{\gamma+1}^X)^*) \leq \text{crit}(E_\gamma^X)$. Let n_X be minimal such that $\kappa_{n_X}^X \geq \text{lh}(E_\gamma^X)$. Let $\eta_n^X < \eta_X$ be minimal such that $\text{crit}(E_{\eta_n^X}^X) \geq \kappa_n^X$ for $n \geq n_X$.

Let then $\mathcal{N}_n^X := \mathcal{M}_{\eta_n^X}^X$ and $v_{n,m}^X := \iota_{\eta_n^X, \eta_m^X}^X$ for $m \geq n \geq n_X$. It is easy to see that the maps are as wanted, but it remains to check that \mathcal{N}_n^X is $(k_X + 1)$ -sound above κ_n^X . This is going to be the one critical use of the assumption that κ_n^X is a cutpoint.

We have to show that the generators of the iteration up to η_n^X are bounded by κ_n^X . If η_n^X is a limit this is obvious as by choice of η_n^X all previous critical points are less than κ_n^X . So assume that $\eta_n^X = \delta + 1$ and E_δ^X has a generator $\geq \kappa_n^X$. By the initial segment condition we then have that the trivial completion G of $E_\delta^X \upharpoonright \kappa_n^X$ is on the sequence of K_X . But we have $\text{crit}(G) = \text{crit}(E_\delta^X) < \kappa_n^X$ and $\text{lh}(G) > \kappa_n^X$, contradicting that κ_n^X is a cutpoint. \square

The covering argument goes through three cases. Thanks to Lemma 49 we can eliminate one of these cases, we will now see that we can also eliminate the other less than convenient case.

Lemma 51. *If \mathcal{N}_n^X for $n \geq n_X$ has a top extender, then μ_n^X , its critical point, is $\geq \kappa_n^X$.*

Proof. Let us first assume that \mathcal{N}_n^X has been constructed according to Case 1 or Case 2. Then λ_X is a limit cardinal in \mathcal{N}_n^X and thus by (2) μ_n^X cannot be smaller than λ_X .

If \mathcal{N}_n^X is constructed according to Case 3, then some ordinal $\geq \kappa_n^X$ has to be the critical point of an extender on the \mathcal{N}_n^X -sequence. As no overlaps can exist on the \mathcal{N}_n^X -sequence, $\mu_n^X \geq \kappa_n^X$ follows. \square

Remark 52. Note that \mathcal{N}_n^X in the notation of [20] is the mouse \mathcal{P}_γ where $\kappa_n = \aleph_\gamma^{K_X}$. Recall that \mathcal{P}_γ is the least initial segment (if it exists) of \mathcal{M}_δ^X that defines a subset of κ_n not in K_X where $\delta < \zeta_X$ is least such that $\text{gen}(E_\delta^X) > \kappa_n$. In addition, by the preceding lemma $\mathcal{P}_\gamma = \mathcal{Q}_\gamma$, i.e. we are avoiding protomice in this construction.

Let then $\mathcal{N}_X := \text{dirlim}(\langle \mathcal{N}_n^X, v_{n,m}^X : n_X \leq n \leq m < \omega \rangle)$ and $v_n^X : \mathcal{C}_0(\mathcal{N}_n^X) \rightarrow \mathcal{C}_0(\mathcal{N}_X)$ be the direct limit map. It should be easy to see that \mathcal{N}_X is wellfounded and that the direct limit maps are $r\Sigma_{k_X+1}$ -elementary as they are generated by an iteration on K . But more is true:

Lemma 53. *The phalanx $((K_X, \mathcal{N}_X), \lambda_X)$ is iterable.*

Proof. We cannot quote [20] here as it seems a priori possible that the mouse (or weasel) \mathcal{P}_β , where $\lambda_X = \aleph_\beta^{K_X}$, from that proof is not equal to \mathcal{N}_X . (This would happen if $(\lambda_X^+)^{K_X}$ is not equal to $(\lambda_X^+)^{\mathcal{N}_X}$.)

Nevertheless, the proof presented in [20] works just as well with \mathcal{N}_X substituted for \mathcal{P}_β .

For those readers not content with this answer, we want to point out that there is an easy cheat available to us in this case as (2) implies that λ_X must be a cutpoint in \mathcal{N}_X , and hence the iterability of the phalanx reduces to the iterability of \mathcal{N}_X . The latter holds as \mathcal{N}_X is an iterate of the core model. \square

Theorem 54. *Let λ be a singular cardinal of countable cofinality. Let $\vec{\kappa} := \langle \kappa_n : n < \omega \rangle$ be a sequence of K -cut points cofinal in λ . Let $\tau_n := (\kappa_n^+)^K$, then $\prod_{n < \omega} \tau_n$ carries a continuous, tree-like scale.*

Proof. We will show that $\vec{f} := \langle f_\alpha^{\vec{\kappa}, K} : \alpha \in C^{\lambda, K} \rangle$ as defined in the last section is that scale. Towards that purpose we need to show that this sequence is cofinal in $\prod_{n < \omega} \tau_n / J_{bd}$. Let $g \in \prod_{n < \omega} \tau_n / J_{bd}$ be arbitrary. Let $X \prec (H_\theta; \in, K || \theta, \vec{f})$ be of good type, as explained at the beginning of this section, with $g \in X$. It will suffice to show that there is some α_X such that $f_{\alpha_X}^{\vec{\kappa}, K}(n) = \sup(X \cap \tau_n)$ for all but finitely many n .

Let $\langle \mathcal{N}_n^X, v_{n,m}^X : n_X \leq n \leq m < \omega \rangle$ and $\langle \mathcal{N}_X, v_n^X : n_X \leq n < \omega \rangle$ be as previously discussed appropriate to our choice of $\vec{\kappa}$ and X .

The first step will be to show that we can realize the least level of K to define a surjection onto $\sup(X \cap \tau_n)$ by taking an ultrapower of \mathcal{N}_n^X for $n \geq n_X$. Let $\mathcal{O}_n^X := \text{Ult}_{k_X}(\mathcal{N}_n^X; \sigma_X \upharpoonright K_X | \tau_n^X)$ and $\tilde{\sigma}_n^X$ be the ultrapower map for $n \geq n_X$. (This ultrapower is formed using equivalence classes $[f, a]_{\sigma_X}$ where $a \in [\kappa_n]^{<\omega}$ and f is a function with domain $[\kappa_n^X]^{|a|}$ that is $r\Sigma_{k_X}$ -definable over \mathcal{N}_n^X .)

We do know that these models are wellfounded, in fact, the phalanx $((K, \mathcal{O}_n^X), \kappa_n)$ must be iterable. (This is $(2)_\beta$ from [20] or [21], where $\kappa_n^X = \aleph_\beta^{K_X}$.) This means that \mathcal{O}_n^X is an initial segment of K . Furthermore, \mathcal{O}_n^X is sound above κ_n , and $\tilde{\sigma}_n^X(\tau_n^X) = \sup(X \cap \tau_n)$ is a cardinal there by the choice of \mathcal{N}_n^X . This means that \mathcal{O}_n^X is the level of K we are looking for.

The next step must be to tie the sequence $\langle \mathcal{O}_n^X : n_X \leq n < \omega \rangle$ to some level of K projecting to λ . Our candidate is $\mathcal{O}_X := \text{Ult}_{k_X}(\mathcal{N}_X; \sigma_X \upharpoonright K_X | \lambda_X)$. Let $\tilde{\sigma}_X$ be the ultrapower map. By Lemma 53 and the lifting properties of our X not only is \mathcal{O}_X wellfounded, but it is an initial segment of the core model. Let $\alpha_X := (\lambda^+)^{\mathcal{O}_X}$.

The last thing we need are appropriate embeddings from \mathcal{O}_n^X into \mathcal{O}_X for $n \geq n_X$. Define $\pi_n^X : \mathcal{C}_0(\mathcal{O}_n^X) \rightarrow \mathcal{C}_0(\mathcal{O}_X)$: let $\pi_n^X([f, a]_{\sigma_X}) = [v_n^X(f) \upharpoonright [\kappa_n^X]^{<\omega}, a]_{\sigma_X}$. It is to be understood here that if f is not an element of $\mathcal{C}_0(\mathcal{N}_n^X)$ but merely definable over it, then $v_n^X(f)$ is the function over $\mathcal{C}_0(\mathcal{N}_X)$ with the same definition and parameters moved according to v_n^X .

Let now f an $r\Sigma_{k_X}$ definable function over $\mathcal{C}_0(\mathcal{N}_n^X)$, ϕ an $r\Sigma_{k_X}$ -formula, and $a \in [\kappa_n^X]^{<\omega}$.

$$\begin{aligned} \mathcal{C}_0(\mathcal{O}_n^X) \models \phi([f, a]_{\sigma_X}) &\Leftrightarrow a \in \sigma_X(\{b \in [\kappa_n^X]^{<\omega} \mid \mathcal{N}_n^X \models \phi(f(b))\}) \\ &\Leftrightarrow a \in \sigma_X(\{b \in [\kappa_n^X]^{<\omega} \mid \mathcal{N}_X \models \phi(v_n^X(f)(b))\}) \\ &\Leftrightarrow \mathcal{C}_0(\mathcal{O}_X) \models \phi([v_n^X(f) \upharpoonright [\kappa_n^X]^{<\omega}, a]_{\sigma_X}) \end{aligned}$$

This shows that π_n^X is $r\Sigma_{k_X}$ -elementary. Consider then the following diagram:

$$\begin{array}{ccc} & & \mathcal{C}_0(\mathcal{O}_X) \\ & \nearrow \bar{\sigma}_X & \uparrow \pi_n^X \\ \mathcal{C}_0(\mathcal{N}_X) & & \mathcal{C}_0(\mathcal{O}_n^X) \\ \uparrow v_n^X & \nearrow \tilde{\sigma}_n^X & \\ \mathcal{C}_0(\mathcal{N}_n^X) & & \end{array}$$

The diagram commutes, and all of $v_n^X, \bar{\sigma}_X, \tilde{\sigma}_n^X$ are cofinal (in $\rho_{k_X}(\cdot)$). Thus so is π_n^X which shows that it is $r\Sigma_{k_X+1}$ -elementary. Also note that the critical point of π_n^X is $\geq \kappa_n$.

It then follows that $\mathcal{C}_0(\mathcal{O}_n^X)$ is isomorphic to $\text{Hull}_{k_X+1}^{\mathcal{O}_X}(\kappa_n^X \cup \{p_{k_X+1}(\mathcal{O}_X)\})$, so $\text{sup}(X \cap \tau_n) = f_{\alpha_X}^{\bar{\kappa}, K}(n)$ for $n \geq n_X$. \square

Remark 55. The last line is inaccurate, as it seems possible that $\alpha_X \notin C^{\lambda, K}$ meaning $f_{\alpha_X}^{\bar{\kappa}, K}$ might not be defined. Nevertheless the structure \mathcal{O}_n^X is definable from α_X and κ_n in K which implies that the sequence $\langle \text{sup}(X \cap \tau_n) : n_X \leq n < \omega \rangle$ is dominated by some $f_{\beta}^{\bar{\kappa}, K}$ for $\beta \in C^{\lambda, K}$.

Corollary 56. *In the above situation $\alpha_X = \text{sup}(X \cap \lambda^+)$.*

Proof. By continuity $f_{\text{sup}(X \cap \lambda^+)}^{\bar{\kappa}, K}$ is the exact upper bound of $\langle f_{\beta}^{\bar{\kappa}, K} : \beta < \text{sup}(X \cap \lambda^+) \rangle$. On the other hand, as we know that \vec{f} is a scale, by the tightness of X we also know that $\langle \text{sup}(X \cap \tau_n) : n < \omega \rangle$ is also an exact upper bound for

this sequence. This implies that both agree almost everywhere, but the latter equals $f_{\alpha_X}^{\vec{\kappa}, K}$ almost everywhere. The desired equality then follows. \square

Let us now move on to the second theorem. This one concerns scales on products that concentrate on ordinals that are inaccessible in K . We will see that scales on such ordinals are significantly more restricted.

Theorem 57. *Let λ be a singular cardinal of countable cofinality. Let $\langle \kappa_n : n < \omega \rangle$ be a cofinal sequence such that each κ_n is an inaccessible limit of cutpoints of K . Assume there is some $\delta < \lambda$ such that ordinals β with $\text{o}^K(\beta) \geq \delta$ are bounded in each of the κ_n . Then $\prod_{n < \omega} \kappa_n$ admits a continuous, tree-like scale.*

Let from now on $\vec{\kappa} := \langle \kappa_n : n < \omega \rangle$ and $\delta < \lambda$ be as in the statement of the theorem. As this theorem deals with scales on ordinals which are inaccessible in K we will have need of a theorem that provides information about the possible cofinalities of such ordinals. In general, we cannot expect these cofinalities to be high because of the existence of Prikry forcing. The next theorem essentially states that this is the only real obstacle. Versions of this theorem for different forms of the core model have existed for some time, but its newest form appropriate for the Jensen-Steel core model is due to Mitchell and Schimmerling.

Theorem 58 (Mitchell-Schimmerling). *Assume there is no inner model with a Woodin cardinal, and let K be the Jensen-Steel core model. Let $\alpha \geq \aleph_2$ be such that α is regular in K , but $\text{cof}(\alpha) < \text{card}(\alpha)$. Then $\text{o}^K(\alpha) \geq \nu$ where $\text{cof}(\alpha) = \omega \cdot \nu$.*

See [19]. Alternatively, as we only deal with linear iterations here it should be plausible that the results from [4] even though not directly applicable can be mimicked here to achieve a similar end.

Let us now again consider some $X \prec (H_\theta; \in, \dots)$ containing relevant objects. In addition to its previous properties we will require that $\text{cof}(\text{sup}(X \cap \kappa_n)) > \delta$. Note then that by our assumption and 58, and this fact will be crucial, $\text{sup}(X \cap \kappa_n)$ is a singular cardinal in K .

We will once again have need of the directed system $\langle \mathcal{N}_{n,m}^X, v_{n,m}^X : n_X \leq n \leq m < \omega \rangle$ and its limit $\langle \mathcal{N}_X, v_n^X : n_X \leq n < \omega \rangle$, but we will require some additional properties.

Lemma 59. *There exist $\tilde{\alpha}_n^X < \kappa_n^X$ such that $\text{Hull}_{\kappa_X+1}^{\mathcal{N}_n^X}(\tilde{\alpha}_n^X \cup \{p_{\kappa_X+1}(\mathcal{N}_n^X)\})$ is cofinal in κ_n^X .*

Proof. If the system is constructed as in Case 1 and 2 then there is a single α such that \mathcal{N}_n^X (which is independent of n) is sound above α so the conclusion follows.

Consider then that the system is constructed as in Case 3. Pick $n \geq n_X$. Recall that $\mathcal{N}_n^X = \mathcal{M}_{\eta_n^X}^X$ and $\gamma + 1$ is the last drop below η_n^X . Note that by definition of η_n^X all critical points before that point are less than κ_n^X . There are two cases.

Case 3.1: $\eta_n^X = \bar{\eta} + 1$.

In that case as κ_n^X is a limit cardinal we must have that $\text{lh}(E_{\bar{\eta}}^X) < \kappa_n^X$. Let $\tilde{\alpha}_n^X < \kappa_n^X$ be such that $\mathcal{M}_{\eta_n^X}^X$ is sound above $\tilde{\alpha}_n^X$.

Case 3.2: η_n^X is a limit ordinal.

Let $\gamma < \beta < \eta_n^X$ be such that $\iota_{\beta, \eta_n^X}^X(\bar{\kappa}) = \kappa_n^X$ for some $\bar{\kappa} \in \mathcal{M}_{\beta}^X$. We must have that $\iota_{\beta, \xi}^X(\bar{\kappa}) \geq \text{crit}(\iota_{\xi, \eta_n^X}^X)$ for all $\xi \in [\beta, \eta_n^X)$. The key is to consider when equality holds in the above equation.

Let us assume towards a contradiction that $\iota_{\beta, \xi}^X(\bar{\kappa}) = \text{crit}(\iota_{\xi, \eta_n^X}^X)$ for an unbounded in η_n^X set A . For $\nu \in \lim(A) \cap \eta_n^X$ we have

$$\text{crit}(\iota_{\nu, \eta_n^X}^X) \geq \sup_{\xi \in A \cap \nu} \text{crit}(\iota_{\xi, \eta_n^X}^X) = \sup_{\xi \in A \cap \nu} \iota_{\beta, \xi}^X(\bar{\kappa}) = \iota_{\beta, \nu}^X(\bar{\kappa})$$

and hence $\nu \in A$. But then $B := \{\iota_{\beta, \xi}^X(\bar{\kappa}) \mid \xi \in A\}$ is a club of indiscernibles in κ_n^X . As σ_X is continuous at points of cofinality ω , $C := \sigma_X''[B]$ is an ω -club in $\sup(X \cap \kappa_n)$. As the latter is singular in K there must exist a club $D \subset \sup(X \cap \kappa_n)$ consisting of K -singulars. But $C \cap D \neq \emptyset$, and C consists of K -regulars. Contradiction!

We conclude that $\text{crit}(\iota_{\xi, \eta_n^X}^X) < \iota_{\beta, \xi}^X(\bar{\kappa})$ for all $\xi \geq \nu$ for some $\nu \in [\beta, \eta_n^X)$. This means that $\iota_{\nu, \eta_n^X}^X$ is continuous at $\iota_{\beta, \nu}^X(\bar{\kappa})$. We then finish the argument by

noticing that \mathcal{M}_{ν}^X is (k_X+1) -sound above $\text{crit}(\iota_{\nu, \eta_n^X}^X)$, and thus $\text{Hull}_{k_X+1}^{\mathcal{M}_{\eta_n^X}^X}(\text{crit}(\iota_{\nu, \eta_n^X}^X) \cup \{p_{k_X+1}(\mathcal{M}_{\eta_n^X}^X)\})$ is cofinal in κ_n^X . \square

Proof of Theorem 57. We want to show that for some $\vec{\alpha}_X := \langle \alpha_n^X : n_X \leq n < \omega \rangle \in X$ the sequence $\langle \sup(X \cap \kappa_n) : n < \omega \rangle$ agrees almost everywhere with $g_{\alpha_X}^{\bar{\kappa}, K, \vec{\alpha}_X}$. (Implicit here is that α_X will be adequate.)

Recall the structures \mathcal{O}_n^X from the proof of the preceding theorem. We will need a slightly different structure here. Let $(\mathcal{O}_n^X)^* := \text{Ult}_{k_X}(\mathcal{N}_n^X; \sigma_X \upharpoonright K_X \upharpoonright \kappa_n^X)$. (This ultrapower is formed using equivalence classes $[f, a]_{\sigma_X}$ where $a \in [\sup(X \cap \kappa_n)]^{<\omega}$ and f is a function with domain $[\gamma]^{|a|}$ where $\gamma < \kappa_n$ is a cardinal with $a \subset \sigma_X(\gamma)$ and f is $r\Sigma_{k_X}$ -definable over \mathcal{N}_n^X . Note that functions with different domains can be compared by adding dummy arguments.)

Let $\bar{\sigma}_n^X$ be the ultrapower map. Note that $\bar{\sigma}_n^X$ maps κ_n^X cofinally into $\sup(X \cap \kappa_n)$ so we have $\text{Hull}_{k_X+1}^{(\mathcal{O}_n^X)^*}(\alpha_n^X \cup \{p_{k_X+1}((\mathcal{O}_n^X)^*)\})$ is cofinal in $\sup(X \cap \kappa_n)$ where $\alpha_n^X := \bar{\sigma}_n^X(\tilde{\alpha}_n^X)$.

The phalanx $((K, (\mathcal{O}_n^X)^*), \sup(X \cap \kappa_n))$ is iterable as $\mathcal{C}_0((\mathcal{O}_n^X)^*)$ can be mapped into $\mathcal{C}_0(\mathcal{O}_n^X)$ by a map with critical point $\sup(X \cap \kappa_n)$, so $(\mathcal{O}_n^X)^*$ is an initial segment of K , in fact, the least one to define a witness to the singularity of $\sup(X \cap \kappa_n^X)$.

Just as before we can map $\mathcal{C}_0((\mathcal{O}_n^X)^*)$ into $\mathcal{C}_0(\mathcal{O}_X)$, so $g_{\alpha_X}^{\bar{\kappa}, K, \vec{\alpha}_X}(n) = \sup(X \cap \kappa_n)$ for all $n \geq n_X$. Note that α_X is adequate relative to the sequence $\vec{\alpha}_X$. The condition (1) $_{\alpha_X}^n$ can be easily fulfilled by taking α_n^X to be a successor ordinal; the condition (3) $_{\alpha_X}^n$ for all but finitely many $n < \omega$ follows for cofinality reasons; the condition (2) $_{\alpha_X}^n$ is fulfilled because we took $\tilde{\alpha}_n^X$ to be such

that $\kappa_n^X \in \text{Hull}_{k_X+1}^{\mathcal{N}_n^X}(\tilde{\alpha}_n^X \cup \{p_{k_X+1}(\mathcal{N}_n^X)\})$ and therefore $\kappa_n \in \text{Hull}_{k_X+1}^{\mathcal{O}_n^X}(\alpha_n^X \cup \{p_{k_X+1}((\mathcal{O}_n^X)^*)\})$, we are done once we notice that $\pi_n^X(\kappa_n) = (\kappa_n^*)_{\alpha_X}$.

We would like to have $\tilde{\alpha}_X \in X$. This is obvious if X is ω -closed. If X is merely internally approachable then we can still find some $\tilde{\alpha}' \in X \cap \prod_{n < \omega} \text{sup}(X \cap \kappa_n)$ that dominates $\tilde{\alpha}_X$ almost everywhere. Then $g_{\alpha_X}^{\tilde{\kappa}, K, \tilde{\alpha}_X}$ and $g_{\alpha_X}^{\tilde{\kappa}, K, \tilde{\alpha}'}$ agree almost everywhere by Corollary 42, so we can replace $\tilde{\alpha}_X$ with $\tilde{\alpha}'$.

By Fodor's Lemma we then have a stationary set of X and a single $\tilde{\alpha}$ such that $g_{\alpha_X}^{\tilde{\kappa}, K, \tilde{\alpha}}$ agrees with $\langle \text{sup}(X \cap \kappa_n : n < \omega) \rangle$ almost everywhere. This then shows that $\langle g_{\alpha}^{\tilde{\kappa}, K, \tilde{\alpha}} : \alpha \in C \cap \text{cof}(>\omega) \rangle$ is a scale. \square

We are going to finish by showing how to weaken the assumption of Theorem 54 yet achieving the same result. It is here that we will make use of the sequence $\langle f_{\alpha}^{\tilde{\kappa}, K, \tilde{\alpha}} : \alpha \in C^{\lambda, K} \rangle$.

We say a cardinal $\kappa \in K$ is a weak cutpoint if $\text{crit}(E) < \kappa$ implies $\text{lh}(E) < (\kappa^+)^K$ for all extenders E on the K -sequence.

Theorem 60. *Let λ be a singular cardinal of countable cofinality. Let $\langle \kappa_n : n < \omega \rangle$ be a sequence of weak cutpoints cofinal in λ . Let $\tau_n := (\kappa_n^+)^K$. Then $\prod_{n < \omega} \tau_n$ carries a continuous, tree-like scale.*

Lemma 61. *There exist some $n_X, k_X < \omega$, a sequence of ordinals $\langle \tilde{\alpha}_n^X : n_X \leq n < \omega \rangle$, a sequence of models $\langle \mathcal{N}_n^X : n_X \leq n < \omega \rangle$, and maps $\langle v_{n,m}^X : n_X \leq n \leq m < \omega \rangle$ such that:*

- $((\kappa_n^X)^+)^{\mathcal{N}_n^X} = \tau_n^X$ and \mathcal{N}_n^X agrees with K_X up to τ_n^X for all $n \geq n_X$;
- \mathcal{N}_n^X is $(k_X + 1)$ -sound above κ_n^X relative to $p_{k_X+1}(\mathcal{N}_n^X) \cap \tilde{\alpha}_n^X$ for all $n \geq n_X$;
- $v_{n,m}^X : \mathcal{C}_0(\mathcal{N}_n^X) \rightarrow \mathcal{C}_0(\mathcal{N}_m^X)$ is $r\Sigma_{k_X+1}$ -elementary for all $m \geq n \geq n_X$;
- $\text{crit}(v_{n,m}^X) \geq \max\{\kappa_n^X, \tilde{\alpha}_n^X + 1\}$ for all $m \geq n \geq n_X$.

Proof. This proof goes through the same cases as the proof of Lemma 50, in fact, many of the cases will be the same. (In those cases we can take $\tilde{\alpha}_n^X$ to be 0.) In the interest of time we shall only deal with the case that is unique to this situation.

Let us assume that $\eta_X := \text{sup}(\{\beta < \zeta_X \mid \text{lh}(E_{\beta}^X) < \lambda^X\})$ is a limit ordinal. Let $\gamma + 1$ be the last drop in the interval $(0, \eta_X)$. Let k_X be minimal such that $\rho_{k_X+1}((\mathcal{M}_{\gamma_1}^X)^*) \leq \text{crit}(E_{\gamma}^X)$. Let n_X be minimal such that $\kappa_{n_X}^X \geq \text{lh}(E_{\gamma}^X)$. Let $\eta_n^X < \eta_X$ be minimal such that $\text{crit}(E_{\eta_n^X}^X) \geq \kappa_n^X$ for $n \geq n_X$.

Let then $\mathcal{N}_n^X := \mathcal{M}_{\eta_n^X}^X$ and $v_{n,m}^X := \iota_{\eta_n^X, \eta_m^X}^X$ for $m \geq n \geq n_X$. Let $n \geq n_X$ be such that $\eta_n^X = \tilde{\eta}_n^X + 1$ and $\text{gen}(E_{\tilde{\eta}_n^X}^X) \geq \kappa_n$. Otherwise the argument will proceed just as in the proof of Lemma 50 (and $\tilde{\alpha}_n^X = 0$).

First note then $\text{gen}(E_{\tilde{\eta}_n^X}^X) < \tau_n^X$ as otherwise κ_n^X could not be a weak cutpoint by the initial segment condition. Moreover, it then follows that $E_{\tilde{\eta}_n^X}^X$ has a largest generator as otherwise $\text{gen}(E_{\tilde{\eta}_n^X}^X) \in (\kappa_n^X, \tau_n^X)$ must be a cardinal in \mathcal{N}_n^X .

Let $\tilde{\alpha}_n^X$ be that largest generator. We will be done if we can show that $\kappa_n^X \cup \{\tilde{\alpha}_n^X\}$ generates the whole ultrapower. Let then $\tilde{\mathcal{M}} := \text{Ult}(\mathcal{M}_{\tilde{\eta}_n^X}^X; E_{\tilde{\eta}_n^X}^X \upharpoonright \kappa_n^X \cup \{\tilde{\alpha}_n^X\})$ and $\tilde{\iota} : \mathcal{C}_0(\tilde{\mathcal{N}}) \rightarrow \mathcal{C}_0(\mathcal{N}_n^X)$ be the canonical embedding.

We have that $\tilde{\alpha}_n^X \in (\kappa_n^X, \tau_n^X)$ is in the range of $\tilde{\iota}$, thus so is $\kappa_n^X = \text{card}^{\mathcal{N}_n^X}(\tilde{\alpha}_n^X)$ and some surjection from κ_n^X on to $\tilde{\alpha}_n^X$. Then $\tilde{\alpha}_n^X \subset \text{ran } \tilde{\iota}$ and thus so are all of the other generators of $E_{\tilde{\eta}_n^X}^X$. \square

Remark 62. Note that in the “special” case of Lemma 61 unlike Remark 52 \mathcal{N}_n^X is not equal to the mouse \mathcal{P}_γ (where $\kappa_n = \aleph_\gamma^{K_X}$) from [20], but it is equal to \mathcal{Q}_γ . To see this we must first realize that \mathcal{P}_γ must be an initial segment of $\mathcal{M}_{\tilde{\eta}_n^X}^X$ as in this case $E_{\tilde{\eta}_n^X}^X$ has generators $\geq \kappa_n$.

As the Dodd projectum of $E_{\tilde{\eta}_n^X}^X$ is below κ_n we have, in fact, $\mathcal{P}_\gamma = \mathcal{M}_{\tilde{\eta}_n^X}^X \upharpoonright \text{lh}(E_{\tilde{\eta}_n^X}^X)$. Note though that lifting this mouse by σ_X would create a proto mouse. Hence we must move to the mouse \mathcal{Q}_γ which is formed by applying the extender $E_{\tilde{\eta}_n^X}^X$ using the usual iteration tree rules. Hence the resulting mouse must be equal to $\mathcal{M}_{\tilde{\eta}_n^X+1}^X = \mathcal{N}_n^X$.

Proof of Theorem 60. Let $\langle \mathcal{N}_n^X : n_X \leq n < \omega \rangle, \langle v_{n,m}^X : n_X \leq n \leq m < \omega \rangle$ and $\langle \tilde{\alpha}_n^X : n_X \leq n < \omega \rangle$ as in the lemma. We will find some α_X and $\vec{\alpha}_X := \langle \alpha_n^X : n_X \leq n < \omega \rangle$ such that $\text{sup}(X \cap \tau_n) = f_{\alpha_X}^{\vec{\kappa}, K, \vec{\alpha}}(n)$ for all $n \geq n_X$. In fact, $\alpha_n^X = \sigma_X(\tilde{\alpha}_n^X)$ will do. A priori $\vec{\alpha}_X$ will depend on X but we will be able to deal with that by pressing down just as in the proof of Theorem 57.

We can mostly proceed as in the proof of Theorem 54. We will form \mathcal{O}_n^X and \mathcal{O}_X as before, and generate embeddings between them by lifting v_n^X . Note that v_n^X will not move $\tilde{\alpha}_n^X$ as iteration maps do not move generators. So neither will its lift move α_n^X . Thus $\mathcal{C}_0(\mathcal{O}_n^X)$ will be isomorphic to $\text{Hull}_{k_X+1}^{\mathcal{O}_X}(\kappa_n \cup \{p_{k_X+1}(\mathcal{O}_X) \cap \alpha_n^X\})$ as required.

As before the fact that the phalanx $\langle \langle K, \mathcal{O}_n^X \rangle, \kappa_n \rangle$ is iterable follows from the covering lemma, noticing that in this case we might have to consider the mouse \mathcal{Q}_β not \mathcal{P}_β as explained above. Fortunately, this does not change anything about the rest of the argument. We skip further detail. \square

Proof of Theorem 5. We have different cases depending on if the κ_i are limit cardinals or successor cardinals in the core model. Let us first assume that all κ_i share a type. If that shared type is limit cardinals, then we can use Theorem 57 to finish. If that type is successor cardinals we have two cases: if $\bar{\kappa}_i$ is the K -predecessor of κ_i is measurable, then it must be a cutpoint by the smallness assumption therefore we can use Theorem 54 to finish; if it is not, then it must be a weak cutpoint thus we can use Theorem 60 to finish.

In cases of mixed types divide the sequence into three parts of pure type. Each of these parts do have a scale with indices in the club $C \subseteq \lambda^+$ by the

above construction. These individual scales can then be integrated. This works as individual elements of the different scales can be tied to some common ordinal $< \lambda^+$. The preservation of tree-likeness is guaranteed by all constructions using the same base mechanic of the condensation lemma. See also Remark 45. \square

5 Open questions

We conclude this work with a discussion on further possible developments, and open questions.

1. Consider the following natural strengthening of the ABSP with respect sequence of regular cardinals $\vec{\tau} = \langle \tau_n \mid n < \omega \rangle$ with $\lambda = \cup_n \tau_n$: For every sufficiently large regular cardinal θ and internally approachable structure $N \prec (H_\theta, \in, \vec{\tau})$, there is some $m < \omega$, so that for every strictly increasing sequence $d_0, \dots, d_k \in \omega \setminus m$ and $F \in N$, $F : [\lambda]^k \rightarrow \lambda$, if

$$F(\chi_N(\tau_{d_1}), \dots, \chi_N(\tau_{d_k})) < \tau_{d_0}$$

then

$$F(\chi_N(\tau_{d_1}), \dots, \chi_N(\tau_{d_k})) \in N.$$

Is it consistent?

2. We saw in Section 2.1 that from the same large cardinal assumptions of Theorem 29, it is consistent that ABSP holds with respect to a sequence $\langle \tau_n \mid n < \omega \rangle$ so that $\tau_n = \aleph_{2n}$ for all $n < \omega$. Is ABSP consistent with respect to cofinite sequence of the \aleph_n 's?
3. The definitions of Tree-like scales, Essentially Tree-like scales, ASFP, and ABFP naturally extend to uncountable sequences of cardinals $\langle \tau_i \mid i < \rho \rangle$, $\rho > \aleph_0$ regular. Are those principles consistent? If so, what is their consistency strength?
4. Another natural extension of the principles AFSP and ABFP, is to require the appropriate principle to hold for any elementary substructure $N \prec (H_\theta \in \vec{\tau})$. Is it consistent?
5. Is there a version of Theorem 6 for Neeman-Steel long extender mice?
6. Pereira showed in [24] that it consistent relative to the existence of a supercompact cardinal that there exist products $\prod_{n < \omega} \tau_n$ carrying a continuous tree-like scale of length greater than $\sup_{n < \omega} (\langle \tau_n \rangle_n)^+$. Can the same be achieved from a weaker large cardinal assumptions at the level of strong cardinals?

References

- [1] Omer Ben-Neria. On singular stationarity ii. *Journal of Symbolic Logic*, 84(1):320–342, 2019.
- [2] Omer Ben-Neria, Moti Gitik, Itay Neeman, and Spencer Unger. On the powersets of singular cardinals in hod. *Proceedings of the American Mathematical Society*, 148:1777 – 1789, 2020.
- [3] Omer Ben-Neria and Spencer Unger. Homogeneous changes in cofinalities with applications to hod. *Journal of Mathematical Logic*, 17(2):1750007, 2017.
- [4] Sean D. Cox. Covering theorems for the core model, with an application to stationary set reflection. *Annals of Pure and Applied Logic*, 161:66 – 93, 2009.
- [5] James Cummings. Continuous tree like scales. *Central European Journal of Mathematics*, 8:314 – 318, 2010.
- [6] James Cummings, Matthew Foreman, and Menachem Magidor. Squares, scales and stationary reflection. *Journal of Mathematical Logic*, 01(01):35–98, 2001.
- [7] Hans-Dieter Donder, R. B. Jensen, and L. Stanley. Condensation-coherent global square systems. In Anil Nerode and Richard A. Shore, editors, *Recursion Theory*, 1985.
- [8] Matthew Foreman and Menachem Magidor. Large cardinals and definable counterexamples to the continuum hypothesis. *Annals of Pure and Applied Logic*, 76(1):47 – 97, 1995.
- [9] Matthew Foreman, Menachem Magidor, and Saharon Shelah. Martin’s maximum, saturated ideals, and non-regular ultrafilters. part i. *Annals of Mathematics*, 127(1):1–47, 1988.
- [10] Moti Gitik. Changing cofinalities and the nonstationary ideal. *Israel Journal of Mathematics*, 56:280–314, 1986.
- [11] Moti Gitik. On a question of pereira. *Archive for Mathematical Logic*, 47:53 – 64, 2008.
- [12] Moti Gitik. Prikry-type forcings. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, pages 1351–1447. Springer Netherlands, Dordrecht, 2010.
- [13] Moti Gitik. Short extenders forcing i. *Journal of Mathematical Logic*, 12(2), 2013.

- [14] Moti Gitik and Menachem Magidor. The singular cardinal hypothesis revisited. In Haim Judah, Winfried Just, and Hugh Woodin, editors, *Set Theory of the Continuum*, pages 243 – 279, 1992.
- [15] Thomas Jech. *Set Theory*. Springer Verlag, 3rd edition, 2002.
- [16] Ronald Jensen. A new fine structure. handwritten notes, online at <https://www.mathematik.hu-berlin.de/~raesch/org/jensen.html>.
- [17] Ronald Jensen and John Steel. \mathcal{K} without the measurable. *The Journal of Symbolic Logic*, 78(3):708 – 734, 2013.
- [18] Peter Koepke. The consistency strength of the free subset problem for ω_ω . *The Journal of Symbolic Logic*, 49(4):1198 – 1204, 1984.
- [19] William Mitchell and Ernest Schimmerling. Covering at limit cardinals of \mathcal{K} . submitted, online preprint at <http://www.math.cmu.edu/~eschimme/Measurables-in-K.pdf>.
- [20] William Mitchell, Ernest Schimmerling, and J Steel. The covering lemma up to a woodin cardinal. *Annals of Pure and Applied Logic*, 84(2):219 – 255, 1997.
- [21] William Mithcell and Ernest Schimmerling. Weak covering without countable closure. *Mathematical Research Letters*, 2:595 – 609, 1995.
- [22] Luis Pereira. *Combinatoire des cardinaux singuliers et structures PCF*. PhD thesis, University of Paris VII, 2007.
- [23] Luis Pereira. The pcf conjecture and large cardinals. *The Journal of Symbolic Logic*, 73(2):674 – 688, 2008.
- [24] Luis Pereira. Morasses, semimorasses and supercompact ultrafilters. *Acta Mathematica Hungarica*, 152:257 – 268, 2017.
- [25] Saharon Shelah. *Cardinal Arithmetic*. Oxford Science Publications, 1994.
- [26] Saharon Shelah. Pcf and infinite free subsets in an algebra. *Archive for Mathematical Logic*, 41:321–359, 2002.
- [27] John Steel. *The Core Model Iterability Problem*, volume 8 of *Lecture Notes in Logic*. Springer Verlag, 1996.
- [28] John Steel. An outline of inner model theory. In M. Foreman and A. Kanamori, editors, *Handbook of Set Theory*, pages 1595 – 1684. Springer Netherlands, 2010.
- [29] John Steel and William Mitchell. *Fine Structure and Iteration Trees*, volume 3 of *Lecture Notes in Logic*. Springer Verlag, 1994.
- [30] Philip Welch. A question on free subsets in internally approachable models. preprint.
- [31] Martin Zeman. *Inner Models and Large Cardinals*. De Gruyter, 2002.